



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>





6000309001



e. 39

27 *M. Haeon*
CATALOGVS
PLANTARVM
HORTI ACADEMICI ET AGRI
GOTTINGENSIS
CONSCRIPTVS

A
JOHANN GOTTFRIED ZINN
M. D. MED. ET BOT. P. P. O. ACAD. REG.
SC. BEROL. SODALI ET SOC. REG.
SC. GOTTING. MEMBRO EXTRAOR-
DINARIO.



GOTTINGAE
CVNTIVS VIDVAE ABR. VANDENHOECK
1757.



VIRIS ILLVSTRISSIMIS
REGIAE MAIESTATIS
BRITANNICAE
IN
REGIONIBVS GERMANICIS
STATVS MINISTRIS
ET
CONSILIARIIS INTIMIS
DOMINIS SVIS CLEMENTISSIMIS

DEMISSO ANIMO

D. D. D.

A V C T O R.

[illegible]

the 1990s, the number of people in the world who are under 15 years of age is expected to increase from 1.1 billion to 1.5 billion. The number of people aged 65 and over is expected to increase from 200 million to 400 million. The number of people aged 15 and over is expected to increase from 3.5 billion to 4.5 billion. The number of people aged 15 and over is expected to increase from 3.5 billion to 4.5 billion. The number of people aged 15 and over is expected to increase from 3.5 billion to 4.5 billion.

VIRI ILLVSTRISSIMI

DOMINI CLEMENTISSIMI.

Summa, qua Academiam Georgiam
Augustam fovetis, VESTRA indul-
gentia, VIRI ILLVSTRISSIMI, ani-
mum & fiduciam mihi addit, vt hunc
Horti Academici, cuius cura clementissi-
me mihi demandata est, plantarum catalo-
gum, VESTRO nomini inscribam. Muni-
ficientia VESTRA factum est, vt sumti-
bus non spernendis fabrica caldarium
emendata, & ad alendas servandasque
plantas rariores aptior facta, sed & ipse
hortus vario modo ornatus, & intra paucis-
simos

simos annos tot plantis ex omni terrarum
ambitu adlatis ita instructus fuerit, vt ca-
talogus aliquot earum centenis auctior
nunc prodire possit. Tantam tamque
insignem benevolentiam, qua semper pre-
cibus meis in commodum horti factis in-
dulistis, demisse veneror, & publico hoc
monumento gratissimam mentem profi-
teor. Vt vero porro benignitate VE-
STRA res nostras adiuuare, hortumque
Academicum ornatiorem reddere curae-
que VESTRAE commendatum habere
velitis, summa, qua par est, veneratione
obsecro. Ita Deus in reipublicae, in A-
cademiae, &, quae pars eius non exigua
est, in horti commodum saluos & inco-
lumes VOS seruet, & ad vltimos sene-
ctutis terminos VOS, & ILLV-
STRISSIMAS VESTRAS FAMI-
LIAS perpetua felicitate ornet. Gottin-
gae d. 14. Apr. MDCCLVII.

PRAE-



PRAEFATIO.

*D*um catalogum plantarum horti & agri Gottin-
gensis edendi consilium sumsi, nonnulla prae-
mittere visum est, quae rationem operis exponant.
Munificentia Illustrissimorum Virorum Regi Magnae
Britanniae & Electoris Brunsvico-Luneburgensis a
Consiliis intimis, factum est, ut non solum sumtibus
non spernendis Horti Academici vaporarium melius
instructum, & ad alendas servandasque plantas ra-
riores aptius redditum, sustentaculisque, in quae
ollae disponantur, horto non exiguum ornamentum
& commodum paratum fuerit, sed eidem quae de-
bemus, si catalogus intra trium annorum spatium
aliquot centenis plantarum auctior prodire possit.
Auditores autem libellum optabant, qui, dum plan-

P R A E F A T I O.

tas recenter decerptas demonstramus, memoriam adiuuare, illosque regere possit, ut in studio Botanico proprio Marte ulterius progredi queant. Omnem ergo operam dedi, ut, commodo illorum inservirem, viamque facilem & tutam, qua ad vberiorem plantarum cognitionem pervenire possint, illos docerem, animo in id quoque intento, ut libellum, quo & aliis in locis & diversis a nostra Florae regionibus vti possent, illis traderem. Methodum elegi, quae, quantum ipsa conditio studii Botanici finit, ad naturalem proxime accedat, & hanc ob rationem maxima ex parte illam a Perillustri HALLERO in Enumeratione stirpium Helveticarum adhibitam servavi, ita tamen, ut classes quasdam naturales prioribus adderem, variisque in locis plantas alium in ordinem disponerem. Classis cuiuslibet definitionem ubique addidi, succinctam, quae tamen scopo nostro sufficiat. Generibus variis, aut propter minutiam partium obscurioribus, aut quorum limites confluere videbantur, praemissi characteres breves & essentielles, experientia edoctus, nimiam difficultatem in investigandis generibus tedium saepe tyronibus creare posse, ut a suavissimo studio facile deterreretur se patiantur. In pluribus

gene-

P R A E F A T I O.

generibus eiusmodi characteres essentialia addere animus erat, nisi moles libelli nimis increfcens obtuiffet. Inter nomina specifica & fynonima plantarum tam ea felegi, quae descriptionem fere brevem, proximasque discriminis notas exprimentem, contineant, quam illa, quae in officinis aut pinacibus v. g. Bauhinianis vel Tournefortianis recepta effe videbam. Descriptiones vero vberiores Specierum & observationes parcius adfperfi, illasque nonnifi breves paucioribus speciebus, quae aut plane novae, aut nonnifi obscure notae minusque definitae videbantur, addidi. Animus autem eft, proxime variarum plantarum, quarum hiftoria nondam fatis extricata eft, aut discriminis notae minus plenae datae effe videntur, vberiore descriptionem & fufiorem Synonimiam proprio libello exhibere; uti fupplemento novos cives, de quibus, etfi in horto iam adfint, quum nondum tamen floruerint, certi quidquam vix fatis tuto determinari poffe videtur, & plantulas illas ad cryptogamiam pertinentes, fpatio hic exclusas, quamprimum fere occafio, subiungam. Pauciffimas fpecies prioris Enumerationis omifi, easque tantum, quae in regionibus Germaniae ab agro Gottingenfi remotioribus


PRAEFATIO.

bus crescunt, nec inter cives Florae nostrae referri possunt. Nullas, si quae forte hoc anno in horto nostro desiderantur, nota peculiari signavi, cum copia plantarum in omnibus hortis res sit mutationi & vicissitudini summe obnoxia, & satis multae, quae in priori enumeratione periisse dicuntur, horum nostrum iterum ornent. Loca natalia forte non satis plena me dedisse nonnullis videbor. Sed mole libelli impeditus, in ipsis demonstrationibus plura imprimis circa plantas indigenas fusiis addi posse credideram. Iconem addidi unicam plantae, quae tam propter novitatem quam ob singularem fabricam peculiari attentione digna videbatur.

Gottingae. d. 20. Apr. MDCCLVII.









APETALAE.

Plantae petalis nullis vel obscurissimis, semine, flore, & veris staminibus donatae.

CLASSIS. I.

AMENTACEAE.

Flores mares & flores feminas a se invicem remotos, aut in eadem, aut in distinctis plantis, ex receptaculo, sive axi communi filiformi gerunt, squamis inter flores interpositis.

ORDO. I.

AMENTACEAE CONIFERAE.

Flos *masf.* Squamae circa axin communem in conum congestae, quae ad basin internam antheras fovent, filamentis aut nullis aut coactis.

Flos *femin.* Squamae similes, ut in mare, quae ad basin internam femina fovent; in non
A nullis

APETALAE

nullis fructus baccam aemulatur. Omnibus praererea indoles resinosa, & angusta sempervirentia folia.

EPHEDRA. LINN. 1007.

Planta anomala, & media inter Equifeta & Coniferas. Diphyta. *Fl. Masc.* Ad squamas singulas coni stamina septem; *Fem.* Semina duo calyce baccato tecta.

H. *Ephedra* pedunculis oppositis: amentis geminis. LINN. *Spec.* 1. *distachya*. p. 1040. *Hort. Cliff.* 465. ROY. *Prodr. Fl. Lugd. Bat.* p. 87. HALLER *Enum. Helv.* p. 145. *Ephedra maritima minor*. I. R. H. HALL. *Enum. Gott.* p. 9. *Vua marina Monspeliensium*. LOBEL. *icon.* p. 796. In Vallesia. Gallia Narbon. Hispania. Fr.

TAXUS LINN. 1006. HALL.

Helv. 146.

Diphyta. *Flos. masc.* Ad squamas coni antherae peltatae octo. *Flos fem.* Calyx in baccam mutatus semen unicum fovet.

1 *Taxus*. omnium auctorum. LINN. *Cliff.* 464. ROY. *Lugd. bat.* 87. HALL. *Helv.* 146. *Enum. Gott.* p. 9. C. BAVH. *Pin.* 505. *Taxus* foliis approximatis. *baccata*. LINN. *Sp.* 1. p. 1040. *Taxbaum: Eibenbaum.* In sylvis circa *Pless* abunde. Fr.

JVNIPERVS. LINN. 1005. HALL. 146.

Diphyta. *Masc.* Ad squamas coni tres antherae connatae. *Fem.* Bacca trisperma, tribus tubis.

Juni-

AMENTACEAE CONIFERAE.

3

Juniperus foliis patentibus. LINN. *Cliff.* 465. ROY. *Lugd. bat.* 89. HALL. *Helv.* p. 146. *Enum. Gott.* p. 9. JUNIPERUS vulgaris α. fruticosa. β. arborea. C. BAVH. *Pin.* 488. *Wachbolder-Staude. Krametsbeeren-Staude.* Gottin-
gae rarius in declivibus montis *Heimberg* ad sini-
stra viae *Robringensfr.* Fruticosa tamen in aliis
regionibus Germaniae. v. g. circa *Jenam* & *Nori-
bergam* copiose; arborea rarius reperitur. Fr.

h. *Juniperus* foliis inferne adnatis; oppo-
sitionibus concatenatis. LINN. *Cliff.* 464. ROY.
Lugd. bat. p. 90. BOEHMER. *Fl. Lips.* p. 283.
HALL. *Enum. Helv.* p. 9. *Juniperus Sabina*
LINN. *Spec.* 6. BLACKW. *T.* 244. *Mas.* *Sabina*
sterilis. I. B. *Femina.* *Sabina baccifera.* I. B. *Hist.*
plant. I. b. 288. *Sebenbaum. Segelbaum.* In Hel-
vetia. Italia. Sibiria. Fr.

h. *Juniperus* foliis inferioribus ternis, supe-
rioribus binis decurrentibus subulatis patulis acu-
tis. *Bermudiana.* LINN. *Sp.* 5. ROY. *Lugdb.* 90.
Juniperus Bermudiana. HERM. *Lugdb.* 345. *T.*
347. In America. Fr.

Juniperus barbadensis & *Juniperus virginiana*

PINUS. LINN. 956. Abies. Larix.

TOURN.

Monophyta. *Masc.* Ad basin anteriorem squa-
marum conii duo loculi antheriferi. *Fem.* Ad ba-
sin squamarum duae nuculae alatae.

Pinus. *Abies.* LINN. *Spec.* 10. *Abies* foliis
solitariis apice acuminatis. LINN. *Cliff.* 449.

A 2

HALL.

HALL. *Helv.* 147. *Enum. Gott.* p. 9. Picea major prima f. Abies rubra BAVH. *Pin.* 493. Picea varior. auctor. Rothe Tanne. Coni penduli longiores, squamæ coni fem. apice subrotundæ. Folia solitaria. Vbique in sylva Hercynica & reliquis borealibus Europae. Fr.

Pinus. *Picea*. LINN. *Spec.* 8. Abies foliis folitariis apice emarginatis. LINN. *Cliff.* 449. HALL. *Enum. Gott.* p. 9. Abies foliorum apice bifido. HALL. *Helv.* p. 148. BLACKW. *T.* 203. Abies Taxi folio, fructu sursum spectante. TOURN. 585. *Weistanne*, Folia inferne albida. In montosis Germaniae. Circa Noribergam. Fr.

Pinus. *Larix*. LINN. *Spec.* 7. Abies foliis fasciculatis obtusis. LINN. *Cliff.* 450. Larix folio deciduo. HALL. *Enum. Helv.* p. 149. Larix. C. BAVH. & reliquorum auctorum. Lerchenbaum. In alpinis Helvetiae, Sibiriae. In sylvis Hercynicis sata laete provenit. Fr.

Pinus. *Sylvestris*. LINN. *Spec.* 1. Pinus foliis geminis, cono pendulo, nucleis strigosis. HALL. *Enum. Helv.* 149. Pinus sylvestris C. BAVH. *Pin.* 491. *Fiechte*. *Kibnbaum*. Folia bina. Coni femmini squamæ apice quadrangulo. Vbique in arenosis Germaniae. Fr.

Pinus foliis ternis. *Taeda*. LINN. *Sp.* 3. Pinus foliis longissimis ex vna theca ternis COLD. *Novch.* 230. In Virginiae, Canadæ paludosis. Fr.

AMENTACEAE CONIFERAE.

THUJA. LINN. 975.

Monophyta. *Mast.* Ad basin squamarum amen-
ti ovati stamina quatuor. *Fl. fem.* Squamae biflo-
rae, styli subulati, semina oblonga alata.

H. *Thuja* strobilis laevibus : squamis obtusis.
occidentalis. LINN. *Spec.* 1. BLACKW. *T.* 210.
Thuja Theophrasti. C. BAVH. *Pin.* 486. Arbor
vitae GLVS. *Hist.* I. p. 36. *Lebensbaum*. In Po-
lonia. Canada. Sibiria. Fr.

H. *Thuja* strobilis squarrosis ; squamis acu-
minatis reflexis. *orientalis*. LINN. *Spec.* 2. HALL.
Enum. Gott. p. 10. In China. Fr.

CUPRESSVS. LINN. 976.

Monophyta. *Thuiae* maxime adfinis. *Fl. masc.*
Ad singulas squamas peltatas antherae quatuor.
Fl. fem. Ad basin squamarum puncta numerosa
truncata loco styli. Nux angulata.

H. *Cupressus* foliis imbricatis erectis. LINN.
Cliff. 449. ROY. *Lugd. bat.* 88. *Cupressus sem-*
peruirens. LINN. *Spec.* 1. p. 1002. BLACKW.
T. 127. *Cupressus* meta in fastigium convolu-
ta, quae femina Plinii. I. R. H. HALL. *Enum.*
Gott. p. 7. 587. *Cupressus* BAVH. *Pin.* 488. &
officin. *Cypresse*. Sponte in Creta. Fr.

ORDO. II.

AMENTACEAE JVLIFERAE.

Aliae monophytae, aliae diphytae. *Fl. masc.*
Amentum, *Julus* dictum, pendulum constans :

fr. Populus foliis subrotundis, dentato-angulatis, subtus tomentosis. LINN. *Cliff.* 400. HALL. *Helv.* n. 1. p. 156. Populus alba. LINN. *Spec.* 1. *Pappel-Weide. Weisser Pappelbaum. An der Unstrut.* In Rheni infulis, Circa Berolinum.

ALNVS. HALLER. 157. Betulae species
LINN.

Monophyta. *Fl. masc.* Ad singulas squamas tres flores tetrastrémones. *Fl. fem.* Semina inter squamas apice crassiores.

Alnus rotundifolia, glutinosa viridis. c. BAVH. *Pin.* 428. HALL. *Enum. Gott.* p. 10. Alnus auctorum: Betula alnus. LINN. *Spec.* 5. *Er-len.* In paludosis & ad rivos. *Fr.*

BETVLA. HALLER. 158.
LINN. 933.

Monophyta Alno similis. *Fr.* Inter squamas coni semina alata.

Betula foliis acuminatis serratis. *Cliff.* 447. HALL. *Helv.* 158. Betula BLACKW. *T.* 240. BAVH. *Pin.* 447, & auctor. Betula alba. LINN. *Spec.* 1. p. 982. *Bircke.* Vbique in dumis sylvisque. *Fr.*

Betula foliis orbiculatis crenatis. *nana* LINN. *Fl. Lapp.* 266. *T. 6. f. 4. Amoen. Acad.* 1. p. 4. *T. 1.* HALL. *Helv.* 158. *Enum. Gotting.* p. 10. *Moor-Bircke.* In paludosis, circa montem *Brüller.* v. g. *Auf der Heinrichs-Höhe.* *Fr.*

AMENTACEAE JVLIFERAE.

CARPINVS. LINN. 951.

Monophyta. *Masc.* Julus ex squamis polystemonibus. *Fem.* Inter squamas foliaceas nuces monospermae.

Carpinus squamis strobilorum planis. *Betula*. LINN. *Spec.* 1. p. 998. *Carpinus* HALL. *Helv.* 158. & *auctorum.* *Charma.* Gall. *Haynbüchen.* *Steinbüchen.* In sylvis. Fr.

QUERCVS. LINN. 949.

Monophyta. *Masc.* Julus ex Floribus disitis. *Fem.* Aliquot scuta plana loco tubae. *Fr.* Nux oblonga in calyce haemispherico, scabro, persistente.

Quercus foliis annuis oblongis superne latioribus; sinubus acutioribus; angulis obtusis *Cliff.* 448. HALL. *Helv.* 159. *Quercus Robur.* LINN. *Spec.* 12. p. 996. *Quercus vulg.* BLACKW. T. 487. & *auctor.* *Eiche.* Ubique in sylvis. Fr.

JUGLANS. LINN. 950.

Monophyta. *Masc.* Ad squamas singulas stamina plurima. *Fem.* Stigmata maxima clavata, lacera. *Fr.* Sub cortice carnosio nux, in quo nucleus quadrilobus.

H. *Juglans* foliis ovalibus glabris subserretis, subaequalibus. *Regia.* LINN. *Sp.* 1. p. 997. Nux *Juglans.* s. regia vulgaris. BAVH. *Pin.* 417. Nux *juglans.* BLACKW. T. 247. *varior.* *auctor.* *Welsche Nüsse.* Colitur in hortis exotica. Fr.

CORYLVS. LINN. 953.

Monophyta. *Masc.* Ad squamas singulas stamina octo ad decem. *Fem.* Fructus s. nuces congestae singulae in calyce lacero.

CORYLVS stipulis ovatis obtusis. *Avellana.* LINN. *Spec.* I. p. 998. HALL. *Helv.* 159. Corylus sylvestris. BLACKW. T. 293. BAVH. *Pin.* 418. & *officin.* *Häselstauden, Häselnüsse.* In sylvis frequens. Fr.

FAGVS. LINN. 951. Castanea. HALL. TOVRN.

Monoph. *Masc.* Flores campanulati quinquefidi axi communi affixi. *Fem.* Fr. Nuces duae in capsula quadrivalvi echinata.

H. *Fagus Castanea.* LINN. *Spec.* I. *Fagus* foliis lanceolatis acuminato ferratis subtus nudis LINN. *Cliff.* 447. *Castanea* α sylvestris β sativa. C. BAVH. *Castanea Helv.* 160. BLACKW. T. 330. & *officinarum.* *Chataigner Gall. Castanien.* Flores mares in cylindrum dispositi. Variis in locis colitur non indigena. Fr.

1 FAGVS foliis ovatis obsolete ferratis *Cliff.* 447. *Fagus officin.* & *omnium fere auctorum.* *Herre.* Gall. *Büchen. Roth-Büchen.* Flores mares in globum digesti. Vbique in sylvis. Fr.

PLATANVS. LINN. 954.

Monoph. *Mares* in amentum globosum digesti. *Fem.* Fructus plures in globum dispositi.

AMENTACEAE JVLIFERAE.

11

H. *Platanus* foliis palmatis. LINN. *Cliff.* 447.
Platanus orientalis. LINN. *Spec.* p. 199.
Platanus BAVH. *Pin. & auctor.* Colitur in hor-
 tis principum. v. g. in horto *Wilhelmsbat* pro-
 pe *Cassel*. Fr.

LIQVIDAMBAR. LINN. 955.

Monoph. *Mares* polystemones in amentum,
 conicum, laxum. *Feminae* ad basin spirae mascu-
 lae, in globum digestae.

H. *Liquidambar* foliis palmato angulatis.
Syraciflua. LINN. *Spec.* 1. p. 999. *Liquidam-*
bar. *Enum.* Gott. p. 12. LINN. *Mat. med.* 432.
 C. BAVH. *Pin.* 502. *offic. & Auctor.* Nota ar-
 bor ob resinam officinalem *Storacis liquidae*. In
 Virginia. Fr.

PISTACIA. LINN. 982. Terebinthus.

Lentiscus.

Diphyta. *Mares* quinquefidi, pentastemones
 in amento laxo. *Fem.* Amentum nullum. *Styli*
tres. Fr. Nux ovata intra drupam.

H. *Pistacia* foliis abrupte pinnatis, folio-
 lis lanceolatis. *Lentiscus*. LINN. *Spec.* 5. *Mat. med.*
 453. *Lentiscus vulgaris* BAVH. *Pin.* 399. & *offi-*
cinarum. *Mastixbaum*. In calidiori Europa. In
 Palaestina. Fr.

H. *Pistacia* *Terebinthus*.

MYRICA. LINN. 981. Gale.

TOVRN.

Diphyta. *Masc.* Ad singulas squamas stamina
 quatuor. *Fem.* *Styli* duo. Fr. bacca vni-
 cularis.

H.

H. *Myrica* foliis lanceolatis, fructu sicco.
 LINN. *Mat. med.* 451. HALL. *Enum. Germ.* p. 112.
 12. *Myrica*. *Gale*. LINN. *Spes.* 1. *Myrica* bra-
 bantica *offic.* In ducatu Cellensi. Fr. inter Calce-
 oclaplan.

H. *Myrica* foliis oblongis opposite sinuatis
Quercifolia. LINN. *Spes.* 4. *Laurus* Africana
 minor, folio *Quercus*. COMMELE H. 2. p. 161.
 T. 81. In Aethiopia. Fr.

HIPPOPHAE. LINN. 980.

Diphyta. *Masc.* In calyce bipartito, staminibus
 quatuor. *Fem.* In calyce tubuloso, bifido, stylo
 simplex. Fr. bacca unilocularis.

H. *Hippophae* foliis lanceolatis. *Rhamnoides*.
 LINN. *Sp.* 1. HALL. *Helv.* 161. *Hippophae*.
 Cliff. 454. ROYEN. 207. In littoribus arenosis
 fluviorum. Fr.

CLASS. II.

NVCAMENTACEAE.

Flores masculi & feminini distincti aut in eadem
 planta, aut in eodem disco. Mares multi, tubu-
 losi, monopetali, squamis interpositis, recepta-
 culo communi affixi, antheris in singulis conni-
 ventibus. Flores feminae aut ad basin marium,
 aut in eodem disco cum maribus. Fructus in
 aliis nux, in aliis semina ex placenta communi
 plura, squamis interpositis.

XANTHIVM. LINN. 937.

Flores *mares* in capitulum collecti. *Fem.* Capsulae infra masculos congestae siccae, aculeatae, apice bifido, biloculares.

Xanthium caule inermi. *Strumarium*. LINN. *Spec.* 1. p. 987. *Mat. Med.* 424. HALL. *Helv.* 151. *Xanthium*. BLACKW. *T.* 444. *Xanthium*. Dioscoridis f. *Lappa minor* C. BAVH. *Pin.* 198. *Spitzkletten*. Circa Salz der Helden. & Allendorf. *Berolini*, in ipsa vrbe prope sterquilina. Ann.

H. *Xanthium* spinis ternatis. *spinosum*. LINN. *Sp.* 2. *Xanthium* Lusitanicum spinosum. HERM. *Parad.* *T.* 246. HALL. *Enum. Gott.* p. 11. In Lusitania. Ann.

AMBROSIA. LINN. 938.

Flores *mares* congesti, calyce communi monophyllo. *Fem.* ad basin marium, tuba bifida. *Fr.* Nux ovata, aculeata, monosperma.

H. *Ambrosia* foliis multifidis, racemis foliariis pilosis. *maritima*. LINN. *Sp.* 4. p. 988. *Ambrosia* *maritima*. BAVH. *Pin.* 138. Ad ripas maris mediterranei. A.

H. *Ambrosia* foliis trilobis ferratis. *trifida*. LINN. *Sp.* 1. *Ambrosia* Canadensis altissima, hirsuta Platani folio. HALL. *Enum. Gott.* p. 11. vbi characterem emendatum huius plantae dedit. In Virginia. Canada. A.

IVA, LINN. 940. Tarchonanthus.

VAILL.

Flores *mares & fem.* in discum congesti, calyce communi. *Mares* in disco, tubulosi, staminibus quinque conniventibus. *Feminae* in ambitu pauciores, ex apice capsulae superne crassioris cordatae, in qua semen exiguum latet; styli duo capillares, vagina quinquedentata inferius cincti.

H. *Iva* foliis lanceolato ovatis, caule herbaceo. *annua*. LINN. *Sp.* 1. p. 988. *Hort. Vps.* 285. Tarchonanthos foliis cordatis serratis trinerviis. ROYEN. 538. In America meridionali. A.

H. *Iva* foliis lanceolatis, caule fruticoso. *frutescens*. LINN. *Spec.* 2. *Gen. nov.* 1106. Parthenium foliis lanceolatis serratis. *Cliff.* 443. ROYEN. 86. In Virginia, Peru. Fr.

PARTHENIVM, LINN. 939. Hysterophorus.

VAILL.

Flores *feminini & androgyni* calyce communi excepti. *Androg.* plures in disco tubulosi. *Fem.* pauciores in radio lingulati, stylo bifido. Semina turbinato cordata.

H. *Parthenium* foliis composito-multifidis. *Hysterophorus*. LINN. *Sp.* 1. Partheniastrum Americanum Ambrosiae folio. NISSOL. *Act. Par.* 1711. p. 423. *T.* 13. f. 2. In Jamaica. A.

MICROPVS. LINN. 892. Gnaphalodes.

TOVRN.

Flores *androgyni & feminini* congesti ad alas foliorum. *Androg.* steriles tubulosi, plures in disco. *Fem.* in radio nudi, quorum germina latent intra singulas calycis squamas galeatas, dorso echinatas. Omnes partes plantae disruptae fila trahunt.

H. *Micropus supinus*. LINN. *Spec.* 1. p. 1. 927. ROY. 145. Gnaphalodes Lusitanica TOVRN. I. R. H. 439. HALL. *Enum. Gott.* p. 12. Ad ripas maris mediterranei. A.

CLASS. III.

Calyxiflorae

Sexus varius. Calyx quadri- aut quinquepartitus, plerumque interne coloratus. Corolla nulla aut petala minima. Stamina tot, quot segmenta calycis, ex calyce orta, paucioribus exceptis, ubi stamina bis tot. Tubae duae vel tres. Fructus plerumque bacca, tot seminibus foeta, quot stigmata adfunt.

ORDO I.

VEPRECVLAE LINN.

EMPETRVM LINN. 977.

Diphyt. *Masc.* Calyx trifidus. Stamina tria. *Fem.* Calyx similis. Styli novem, bacca totidem seminibus foeta.

Empe-

Empetrum procumbens, nigrum, LINN. *Sp.* 2. p. 1022. HALL. *Helv.* 162. *Erica baccifera*. CLUSII & *aliorum*. In monte *Bructero*. Fr.

VISCVM. LINN. 979. HALLER.
Helv. 162.

Monoph. HALLER. *Diphyta* LINN. *Masc.* Stamina sessilia quatuor *Fem.* Calyx quadrifolius, bacca monosperma, glutine plenissima. Flores ad ortum foliorum conglomerati. Planta parasitica.

Viscum. album. LINN. *Sp.* 1. HALL. *Helv.* 162. *Viscum baccis albis*. C. BAVH. *Pin.* 423. *Viscum* BLACKW. T. 184. & *officin.* *Mistel*. In variis arboribus, in primis in *Pyro*. Fr.

BVXVS. LINN. 934. HALLER. 163.

Monoph. Flores utriusque sexus misti in gemmis plantae. *Masc.* Calyx quadrifolius. *Fem.* Calyx similis. Capsula trirostris, trilocularis. Folia sempervirentia.

H. *Buxus. sempervirens*. LINN. *Spec.* 1. HALLER. 163. BLACKW. T. 196. & *offic.* *Buxbaum*. Sponte in calidiori Europa, in hortis culta & tonsa. Fr.

RHAMNVS. LINN. 235. *Frangala*. *Paliurus*. *Alaternus*. *Ziziphus*. *Cervi spina*.

TOVRN.

Calyx quadri- & quinque-partitus. Ad singulas divisiones segmentorum squamulae introrsum conniventes, sub quibus stamina.

Rhamnus

Rhamnus spinosus, foliis acuminatis, serratis. HALL. *Helv.* 163. *Enum. Gott.* p. 13. *Rhamnus catharticus*. LINN. *Sp.* 1. *Rhamnus catharticus* BAVH. *Pin.* 478. BLACKW. *T.* 135. Spina cœvina *Offic.* *Creutzbeer.* *Wegdorn.* Planta diphyta. Calyx quadrifidus. Stigma quadrifidum. In adlescentia M. Hengst. circa Plesz, & alibi in sepibus. Fr.

H. *Rhamnus inermis*, foliis glabris, crenatis. HALL. *Comment. Soc. Sc. Gott.* p. 349. *T.* 16. *En. Gott.* p. 13. *Rhamnus alpinus*. LINN. *Sp.* 2. In horto tantum adest femina, cuius calyx quadrifidus, stigma tri- & quadrifidum. Sponte in alpibus Helveticis. Fr.

Rhamnus inermis, floribus monogynis hermaphroditis, foliis integerrimis. *Frangula*. LINN. *Spec.* 3. ROY. 224. *Frangula* pra folii integra. HALL. *Enum. Helv.* 164. *Frangula*. *Alnus nigra*. BLACKW. *T.* 152. *Offic.* *Faulbaum*. Calyx quinquefidus. Stigma emarginatum. Bacca disperma. In humidis v. g. circa *Landolfshausen*. Ad sepes super *Weende*. Fr.

H. *Rhamnus inermis* floribus dioicis, stigmate triplici. *Alaternus*. LINN. *Spec.* 4. *Phytica elatior*. C. BAVH. *Pin.* 427. *Alaternus* GLVS. & *varior. auctor.* Calyx quinquefidus. Squamae calycis deficiunt. In calidiori Europa. v. g. Hispania. Fr.

H. *Rhamnus aculeis geminatis rectis*, floribus digynis, foliis ovato-oblongis. *Zizyphus* LINN. *Sp.* 9. *Juiuba sylvestris*. BAVH. *Pin.* 446. *Zi-*

zyphus. DOD. *Pempt.* 807. *Juiuba Offic. Rotbe Brust-Beeren.* Calyx quinquefidus. Baccae nucleus bilocularis. In Europa Australi, Africa & Syria. Fr.

H. *Rhamnus aculeis geminatis*, inferiore reflexo, floribus trigynis. *Paliurus.* LINN. *Sp.* 5. *Paliurus.* TOURN. 387 & Auctor. Nucleus trilocularis, calyx quinquefidus. In Europa australi. Fr.

THESIVM. LINN. 258. HALLER. 183.
Linofyris. Linophyllum.

Calyx quinquepartitus. Tuba stigmatibus globosa. Semen in fundo calycis.

Thesium panicula foliata, foliis linearibus. *Linophyllum.* LINN. *Sp.* 1. *Thesium Cliff.* 41. HALL. *Helv.* 183. In montosis siccis Germaniae, v. g. M. *Bructero, Weisner.* Fr.

Quae sequuntur duo genera, stamina bis tot, quot segmenta calycis habent.

ORD. II.

SCLERANTHVS. LINN. 497. Knawel. T.

Calyx semiquinquefidus, segmentis acutis oris albis. In fundo calycis capsula disperma.

Scleranthus calycibus fructus patulis. *annuus.* LINN. *Sp.* 1. p. 406. Knawel annum maius, flore minori. HALL. *En. Gott.* p. 32. Polygonum minus polycarpon. TABERN. p. 834. Vbique in agris. A. Scle-

Scleranthus calycibus fructus clausis. *perennis*. LINN. *Sp.* 2. Kewel incanum, flore maiore, perenne. HALL. *Helv.* 186. Polygonum Polonicum, cocciferum. 1. BAVH. III. p. 378. Celebris olim planta propter gallam, radici adhaerentem, enatam ex sic dicto Progallo-insecto, quae ad rem tinctoriam & Confectionem *Alkermes* in usu fuit. In arenosis *Cellensibus*. P.

DAPHNE. LINN. 436. Thymelaea. HALL. 188. TOURN. 366.

Calyx petalo similis, caducus, tubulosus, limbo quadrifido. Bacca unilocularis.

Daphne floribus sessilibus infra folia elliptico-lanceolata. LINN. *Cliff.* 147. *Mat. med.* 179. *Daphne Mezereum*. LINN. *Spec.* 1. p. 356. Thymelaea spica foliis levibus, mollibus lanceolatis subiecta. Enum. *Helv.* 166. Enum. *Gött.* p. 166. Laureola folio deciduo flore purpureo officinis laureola femina. BAVH. *Pin.* 462. *Kettersbals*. Vbiq. in sylvis primo vere. Fr.

Daphne Laureola. LINN. *Sp.* 5. Thymelaea foliis duris levibusque, floribus intermissis. HALL. Enum. *Gött.* p. 166. Laureola sempervirens, flore viridi, quibusdam Laureola mas. c. BAVH. *Pin.* 462. BLACKW. T. 62. In Helvetiae montosis circa *Heidelberg*. Fr.

CHRYSOPLINUM. LINN. 493.

Calyx quadri- & quinquepartitus, coloratus, patulus. Stamina ex receptaculo orta. Capsula unilocularis, birostris, calyce cincta.

B 2

Chry.

Chrysosplenium foliis alternis. LINN. *Spec.* 1.
 HALL. *Helv.* 189. Chryf. foliis pedunculis ob-
 longis insidentibus. 1. R. H. 146. In vdis opacis.
 Flores racemosi. A.

ORD. III.

Prioribus omnino adsimiles. Styli autem duo
 ex fructu sicco monospermo.

VLMVS. LINN. 281.

Stamina calyce colorato longiora. Fructus la-
 tus. Flosculi ex eadem gemma plures.

Vlmus fructu membranaceo. LINN. *Cliff.* 83.
 HALL. *Helv.* 167. Vlmus *campestris*. LINN. *Sp.*
 1. p. 225. Vlmus *campestris* & Theophrasti.
 BAVH. *Pin.* 246. *Vlmenbaum*. In sylvis & ad
 pagos. Fr.

BOSEA. LINN. 280.

Calycis segmenta lanceolata. bacca globosa,
 Flores racemosi.

H. Bosea. *Yervamora*. *Spec. pl.* 1. p. 225.
 ROY. *Lugdb.* 223. Bosea arbor. HALL. *Gott.* p.
 14. Frutex peregrinus horto Bosiano Yervamo-
 ra dictus. WALTH. *Hort.* 24. T. 10. In insulis
 Canariis. Fr.

CELTIS. LINN. 1012. LVDW. 782.

Calycis segmenta patula. Flores *androg.* &
masc. Stamina breviora. Styli pubescentes. Fr. ro-
 tundus nuce foetus. Flores in alis foliorum soli-
 tarii.

CALYCIFLORAE.

22

H. Celtis foliis ovato-lanceolatis: *australis*. LINN. *Sp.* 1. Celtis fructu nigricante. HALL. *Gott. p.* 14. Lotus arbor. LOS, *ic.* 186. In Europa australi. Fr.

ORD. IV.

Plantae variae arboreae, prioribus multum adsimiles, vario tamen modo anomalae.

MORVS. LINN. 936.

Pl. Monophyta. Flores fere amentacei. *Mas.* Stamina calyce longiora quatuor. *Fem.* Bacculae multae pulposae, baccam compositam faciunt.

H. Morus foliis oblique cordatis laevibus. *alba*. LINN. *Spec.* 1. Morus fructu albo. BAVH. *Pin.* 459. Morus *alba*. HALL. *Gott. p.* 14. *Weisser Maulbeerbaum*. Celebris propter Bombycem, quae folia illius pascitur, Sponte in China, nunc vbique fere in Europa colitur, coeli nostri satis patiens. Fr.

H. Morus foliis cordatis scabris. *nigra*. LINN. *Sp.* 2. *Mat. med.* 422 Morus *nigra*. BLACKW. *T.* 126. BAVH. *Pin.* 459. *Schwartzer Maulbeerbaum*. Sponte in Italia. Nobis in hortis colitur. Fr.

LAVRVS. LINN. 452. LVDW. 133.

Stamina novem, quorum tria inferiora ad basin glandulifera. Ex tuberculis filamenta circa germen. Tuba simplex. Sub drupa nux. Sexus & corolla variat. Flores racemosi. Fr.

H. Laurus foliis lanceolatis venosis perennantibus, floribus quadrifidis dioicis. *nobilis*. LINN. *Sp.*

4. *Matt. med.* 194. *Laurus vulgaris*. C. B. *Pin.* 460. HALL. *Gott.* 15. BLACKW. *T.* 175. *Lorbeerbaum*. In Italia. Graecia. Fr.

H. *Laurus* foliis lanceolatis perennantibus venosis planis, ramulis tuberculatis cicatricibus, floribus racemosis. *Indica*. LINN. *Sp.* 5. *Laurus Indica*. PLUK. HALL. L. cu ubi vide characterem. In Virginia. Fr.

FICVS. LINN. 1032.

In interna superficie calycis communis carnosus flosculi mares & feminae. *Mas.* Calyx tripartitus, stamina tria. *Fem.* Calyx quinquepartitus. Tuba bifida.

15. *Ficus* foliis palmatis. *Carica*. LINN. *Spec.* 1. p. 1059. *Matt. med.* 478. *Ficus communis*. BAVM. *Pin.* 457. BLACKW. *T.* 125. *Feigenbaum*. In Europa australi. Asia. Fr.

H. *Ficus* foliis ovatis integerrimis obtusis, caule inferne radicato. *Benghalensis*. LINN. *Sp.* 4. *Ficus Benghalensis*. &c. H. *Gott.* p. 14. In India. Fr.

CLASSIS IV.

SARMENTACEAE. LINN.

Omnes isostemones; stamina quatuor ad sex, tuba una ad tres. Fructus ficus aut bacca. Caudes plerumque volubiles aut sarmentosi. Flores racemosi.

CANNABIS. LINN. 988.

Diphyta. *Mas.* calyx quinquepartitus. *Fem.* Calyx monophyllus altero latere hians. Styli duo. *Nux bivalvis*. H.

SARMENTACEAE.

23

H. Cannabis, foliis digitatis. LINN. *Mat. med.* 457. fativa. LINN. *Sp.* 1. Cannabis mas & femina. *Aust.* HALL. *Gott.* 14. BLACKW. *T.* 223. a. b. *Hanf.* Sponte in India: nobis in agris colitur. A.

HVMVLVS. LINN. 989. Lupulus.
HALL. 166.

Pl. Diphyta. *Mares* pentastemonones racemosi. Flores *fem.* in amentis. Ad squamas maiores singulas styli duo. Semen vnicum. Caulis volubilis.

Humulus. LINN. *Sp.* 1. p. 1028. *Mat. med.* 458. Lupulus. HALL. *Helv.* & *offic.* *Hopsen.* Ad sepes. P.

VITIS. LINN. 250. HALL. 166.

Petala quinque emarcida, minima. Tuba obtusa. Bacca unilocularis. Flores racemosi. Caulis caepreolis scandens.

H. Vitis foliis palmato angulatis. LINN. *Mat. med.* 97. vinifera. LINN. *Sp.* 1. HALL. *Helv.* 166. BLACKW. *T.* 153. *Weinstock.* Sponte in Europa calidiori. ad sepes. Fr.

HEDERA. LINN. 249. HALL. 165.

Petala quinque oblonga. Stylus simplex. Bacca pentasperma. Flosculi umbellati.

Hedera foliis ovatis & lobatis. *Helix*. LINN. *Sp.* 1. *Mat. med.* 98. HALL. *Helv.* 164. BLACKW. 188. α. sterilis, β. arborea. BAVH. *Pin.* 305. *Ephēn.* *Wintergrün.* Circhis arboribus & muris adhaeret, &

scandit parasitica. Prior α . ubique ad arbores. β . in calidioribus floret, sed & circa *Plesi*. Fr.

H. *Hedera* foliis quinatis ovatis serratis, *quinquefolia*, LINN. *Sp.* 2. *Vitis* quinquefolia Canadensis, CORNVT. *Canad.* T. 100. HALL. *Gott.* 179. In Canada. Fr.

MENISPERMVM, LINN. 413.

Calyx sexfolius. Petala sex. Styli tres. Baccæ tres monospermae. Caulis volubilis.

H. *Menispermum* foliis peltatis subrotundis angulatis, *Canadense*. LINN. *Sp.* 1. *Menispermum* Canadense scandens, umbilicato folio. TOVRN. *Art. Par.* 1705. p. 311. In Virginia. Canada. P.

SMILAX, LINN. 992.

Diphyta. Calyx sexfolius in utroque sexu. *Fem.* Styli tres. Bacca trilocularis. Planta capreolis scandens.

H. *Smilax* caule aculeato angulato, foliis dentato aculeatis cordatis. *aspera*. LINN. *Sp.* 1. *Smilax aspera* fructu rubente. BAVH. *Pin.* 296. HALL. *Gott.* 78. In Hispania. Italia. Fr.

H. *Smilax* caule aculeato angulato, foliis inermibus sagittatis, *excelsa*. LINN. *Sp.* 2. *Smilax* orientalis, aculeatis farmentis, foliis non spinosis. TOVRN. *Cor.* 45. HALL. *Gott.* 79. In Oriente. Syria. Fr.

RVSCVS, LINN. 1005.

Pl. Diphyta. Calyx sexfolius, & nectarium ovatum, monophyllum, in utroque sexu. *Mas.* Stamina tria.

ria. Fem. Stylus vnicus. Bacca globosa. Caulis geniculatus.

H. *Ruscus foliis supra floriferis nudis. aculeatus* LINN. *Sp. 1. Mat. med.* 768. *Bruscus* vel *Ruscus*. BLACKW. *T.* 155. *Ruscus myrtifolius aculeatus*. I. R. H. 79. HALL. *Gott.* 78. In Anglia. Gallia. Italia. Fr.

H. *Ruscus foliis subtus floriferis sub foliolo. Hypoglossum* LINN. *Sp. 3. Laurus Alexandrina* fructu pedunculo insidente. BAVH. *Pin.* 304. *Hypoglossum Bislingua*. BLACKW. *T.* 128. In Italia. Hispania. Fr.

H. *Ruscus racemo terminali hermaphroditico. racemosus*. LINN. *Sp. 5. Laurus Alexandrina* ramosa, fructu e summitate caulium prodeunte. HERMANN. *Lugd. bot.* 379. *T.* 681, P.

Antonia elegans ... 438.

CLASSIS. V.

OLERACEAE LINN.

Calyx quadri- aut quinquepartitus, segmentis concavis. Petala nulla. Sexus variat. Stamina tot, quot partes calycis, ex thalamo orta, in plerisque elastica. Styli duo vel tres, in paucioribus quatuor. Ovarium calyce plerumque persistente comprehensum, siccum, monospermum. Pleraeque herbae. Flores in plurimis in artissimos glomeres, ad alas foliorum, aut in racemos dispositos, congesti.

BETA. LINN. 274.

Calyx quinquefolius. Capsula semine reniformi. Flores in alis foliorum.

H. *Beta vulgaris*. LINN. *Spec.* 1. *Mat. med.* 113.
BLACKW. *T.* 235. α . *Beta sylvestris maritima*. C.B.
118. β . varietates ex colore caulis & radicis. *Mangold. Rotbe Rüben*. In Angliae & Belgii littoribus
marinis. A. ~~rubra~~ *B. Cruda*.

SPINACHIA. LINN. 986.

Diphyta. *Mas.* Calyx quinquefidus. *Fem.* Calyx quadrifidus, laciniis duabus minimis. Styli quatuor. Flores in alis.

H. *Spinachia oleracea*. LINN. *Sp.* 1. *Mat. med.* 556. BLACKW. *T.* 49. α . semine aculeato. β semine non aculeato. Num varietates? *Spinat.* A.

H. *Spinachia* foliis angulosis, caule ramosissimo. HALL. *Gott.* p. 22. Seminis capsula foliolo terminata. A.

Spinacia alba v. pag 429.

ATRIPLEX. LINN. 1021. HALL. 173.

Flores in eadem planta androgyni, mares & feminae. *Androg.* Calyx quinquefolius. Stamina elastica. Ovarium compressum. *Masc.* Calyx & stamina similia. *Fem.* Calyx maior bivalvis, cuius valvulae compressae, magnae, semen androgyni simile includunt.

H. *Atriplex* caule erecto herbaceo, foliis triangularibus. *hortensis*. LINN. *Sp.* 4. *Mat. med.* 473. HALL. *Gott.* p. 18. α . *Atriplex hortensis alba*. β . *Atriplex hortensis rubra*. BAVH. *Pin.* 119. BLACKW. *T.* 99. *Melte*. Sponte in Tataria. A.

Atriplex

Atriplex foliis incanis, ferratis, capsulis quadrangularibus, dentatis. HALL. *Gott.* 19. *Atriplex sylvestris*, fructu roseo, compresso. C. B. *Prodr.* 58. In Thuringia. A.

Atriplex foliis triangularis, rariter dentatis. HALL. *Gott.* 1. c. *Atriplex basifolia*. LINN. *Sp.* 6. *Atriplex sylvestris* folio hastato, f. deltoide. RAII *histor.* 192. Circa loca falsa *Harsle*, & alibi in vicina. A.

Atriplex foliis imis sagittatis, superioribus lingulatis, valvis feminis dentatis. HALL. *Gott.* 19. *Atriplex putula*. LINN. *Sp.* 7. *Atriplex angustifolia*, oblongo folio. BAVH. *Pin.* 219. In hortis sponte, ad fossas & vias. A.

H. *Atriplex fruticosa*, foliis ovato-triangularibus, incanis. HALL. *Gott.* 1. c. *Atriplex Halimus*. LINN. *Sp.* 1. *Halimus latifolius* f. fruticosus. BAVH. *Pin.* 120. Ad sepes maritimas Hispaniae, Lusitaniae & Virginiae. Fr.

H. *Atriplex fruticosa*, foliis lingulatis incanis. HALL. *Gott.* p. 19. *Atriplex Portulacoides*. LINN. *Sp.* 2. *Portulaca marina*. DOD. *Pempt.* 771. Calyx feminae quinquefolius. Ad littora marina Sueciae, & ad mare mediterraneum. Fr.

CHENOPODIVM. LINN. 273. HALL. 175.

Flores omnes androgyni, androgynis prioris generis simillimi. Stylus bipartitus. Semen orbiculatum.

H. *Chenopodium* foliis sinuatis, subhirsutis. HALL. *Gott.* 20. *Chenopodium* foliis oblongis sinuatis, racemis nudis multifidis. *Botrys*. LINN. *Sp.*

8. *Mat. med.* 109. Botrys. BLACKW. 314. In Europae australis arenosis. A.

H. *Chenopodium foliis subhirsutis, oblongis dentatis, incanis.* HALL. *Gott.* 20. *Chenopodium foliis lanceolatis dentatis, racemis foliaceis simplicibus.* *Ambr. sioides* LINN. *Sp.* 9. Botrys ambrosioides Mexicana. BAVH. *Pin.* 138. In Lusitania. Mexico. A.

H. *Chenopodium foliis sinuatis, subtus incanis.* HALL. *l. c.* *Chenopodium glaucum.* LINN. *Sp.* 12. *Chenopodium angustifolium laciniatum minus.* TOURN. *I. R. H.* 506. Ad Europae finemeta & aquas salias. A.

Chenopodium foliis levibus, difformiter dentatis, calycibus rubris. HALL. *l. c.* 4. *Chenopodium rubrum.* LINN. *Sp.* 3. *Chenopodium n.* 3. HALL. *Helv.* 174. *Atriplex sylvestris latifolia.* BAVH. *Pin.* 119. & *offic.* *Pes anserinus.* TABERN. *ic.* 427. *Gänse-Fuß. wilde Melde.* (Inter segetes & locis ruderalis.) A. *Chenopodium foliis levibus, viridibus, septangulis.* HALL. *Gott.* 20. *Chenopodium hybridum.* LINN. *Sp.* 7. *Chenopodium stramonii folio.* VAILL. *Par.* 36. *T. 7. f. 3.* *Neun-Spitzen.* Sautodt. In locis cultis. A.

Chenopodium foliis subtus farinosis, rhomboideis, dentatis. HALL. *l. c.* *Helv.* 175. *Chenopodium viride.* LINN. *Sp.* 6. *Chenopodium Opuli folio.* VAILL. *T. 7. f. 1. a.* *Chenopodium foliis inferioribus ovatis, acutis, summis lineari-lanceolatis.* HALL.

HALL. l. c. *Chenopodium folio sinuato, candicante.*

1. R. H. Locis similibus. A.

Chenopodium album L. *Chenopodium album* L. *Chenopodium album* L. *Chenopodium album* L. *Chenopodium album* L.

Chenopodium foliis triangulari-sagittatis integerrimis. Bonus Henricus. LINN. *Spec.* 1. *Mat. med.*

106. HALL. *Helv.* 174. *Lapathum vinctuosum.*

BAVH. *Pin.* 115. *Bonus Henricus offic.* Guter Hein-

rich. Ad pagos & ruderosa. A.

Chenopodium album L. *Chenopodium album* L. *Chenopodium album* L. *Chenopodium album* L. *Chenopodium album* L.

Chenopodium ex alis racemosum, foliis glabris rhomboideis, serratis. HALL. *Gott.* p. 21. *Chenopodium*

Atriplicis sativae facie, spicis surrectis racemosis. RUPP. *Fl. Jen. Hall.* p. 338. Ad lacum

salum, aliisque locis Saxoniae. A.

Chenopodium foetidum, foliis farinosis, glaucis, obtuse rhomboideis. HALL. l. c. *Helv.* 175. n.

6. *Chenopodium. Vulvaria.* LINN. *Spec.* 13. *Mat.*

med. 108. *Atriplex olida.* BLACKW. *T.* 100. *Atriplex*

foetida. C. B. *Pin.* 119. *Vulvaria* TABERN. *ic.*

428. *Stinckende Melze.* In horto sponte, in cultis

& ruderosis. A.

Chenopodium foliis ovato-lanceolatis integerrimis. HALL. l. c. *Helv.* n. 7. p. 176. *Chenopodium*

polypermum. LINN. *Sp.* 14. *Chenopodium*

Betae folio. I. R. H. 506. In agris & hortis

sponte. A.

H. *Chenopodium foliis lineari-lanceolatis, ramis recurvis, dichotome ramosis, setis terminatis.* HALL,

omm. Soc. Sc. Gott. 1752. p. 347. *T.* 15. f. 3.

Chenopodium aristatum. LINN. *Sp.* 21. In Sibi-

a. A.

H. *Cheno-*

H. *Chenopodium foliis subulatis superne planis, subtus convexis.* LINN. *Cliff.* 86. HALL. *Gott.* p. 28. Chen: *maritimum.* LINN. *Sp.* 16. Kali minus album, semine splendente. BAVH. *Pin.* 289. Ad marina littora Europae. A.

H. *Chenopodium foliis linearibus, hirsutis, ascendentibus.* HALL. *l. c.* *Chenopodium.* *Scoparia.* LINN. *Spec.* 15. *Chenopodium Lini folio* I. R. H. *Revedere* hortulanorum. In Graecia, China, Japonia. A.

LERCHIA. HALL. *Comm. Soc. Sc. Gott.*

1751. p. 223. *ol. supine nulla*

Priori generi valde simile genus. Sed tubae tres ad quinque. Fructus oblongus, sursum gracilior, in cuius fundo semen lenticulare. Flores in summitatibus inter folia.

H. *Lerchia foliis obtusis.* HALL. *Gott.* 21. *Comm. Gott.* T. 1. p. 223. *Chenopodium fruticosum.* LINN. *Sp.* 17. In Persia. Ad mare Britannicum. Fr.

H. *Lerchia foliis filiformibus lacutis.* HALL. *Gott.* 22. *Comm. Gott.* l. c. p. 224. *Chenop. altissimum.* LINN. *Sp.* 18. Ad *Astracanthum.* Fr.

PARIETARIA. LINN. 1020. HALL. 177.

In eadem planta flores androgyni, mares & feminae. *Androg.* Calyx profunde quadrifidus, stamina elastica. Tuba plumosa. *Mas.* Stamina maiora. *Fem.* Calyx & fructus, ut in androgyno. Glomeruli florum densissimi.

Parie

Parietaria foliis ellipticis. HALL. *Gott. p.* 27. *Parietaria officinalis.* LINN. *Sp.* 1. *p.* 1052. *Mat. med.* 472. *Parietaria* BLACKW. *T.* 156. *Parietaria officinarum & Dioscoridis.* C. B. *Pin.* 121. *Glass-Kraut.* Circa *Northeim* & alibi in ruderatis. A.

Parietaria foliis ovatis. HALL. *Gott. l. c.* *Parietaria Lufitanica.* LINN. *Spec.* 2. *Parietaria ocimi folio, l. minor.* HALL. *Fl. Jen. Rupp.* 340. In muris Germaniae inferioris. Olim *Gottinae* in area horti. Nunc evanuit.

VRTICA. LINN. 935. HALL. *p.* 177.

Monophyta praeter unicam speciem. *Mat.* *Parietariae* mari similis. *Fem.* Calyx bivalvis, tuba plumosus, semen ovatum. Flores in racemis.

α. *Vrtica foliis oppositis, amentis fructiferis globosis.* *vilulifera.* LINN. *Sp.* 1. *p.* 983. *Mat. med.* 430. *β.* *Vrtica vilulifera, foliis cordatis, circumscissatis.* HALL. *Gott. p.* 27. *Vrtica Romana.* BLACKW. *T.* 321. *γ.* *Vrtica vilulifera, foliis ovatis integris.* *Gott. l. c.* Num distinctae species? In Europa australi. A.

Vrtica foliis ovatis, amentis cylindraceis. HALL. *Helv.* 178. *Vrtica vrens.* LINN. *Sp.* 3. *Vrtica vrens minor.* BAVH. *Pin.* 232. *Hetter-Nessel.* Ubique ad sepes. A.

Vrtica foliis oblongo-cordatis. LINN. *Mat. med.* 419. HALL. *Helv.* 177. *Vrtica dioica.* LINN. *Spec.* 4. *Vrtica vrens maxima.* BAVH. *Pin.* 232. BLACKW. *T.* 12. *Brennnessel.* Ubique in ruderosis, A.

H. Vrtica foliis oppositis, tripartitis, incis. *canabina*. LINN. *Sp.* 5. Vrtica foliis profunde lacinatis, semine lini. AMMANN. *Ruth.* 249. *T.* 25. HALL. *Gott.* 27. In Sibiria. P.

H. Vrtica foliis alternis orbiculato vtrinque acutis, subtus tomentosis. *nivea*. LINN. *Sp.* 12. Vrtica racemifera maxima Sinarum, foliis subtus lanuginosis. HALL. *Gott.* p. 27. In China. P.

AMARANTHVS. LINN. 941.

Glomeres florum in racemos dispositi. Pl. Monophyta. *Mas.* Calyx persistens plerumque coloratus tri- aut quinquefolius, foliis lanceolatis. Stamina tria aut quinque. *Fem.* Calyx, vt in mare. Stylis tres. Capsula ovata, monosperma.

H. Amaranthus capitulis subrotundis, lateralibus, foliis lanceolatis acuminatis. LINN. *Cliff.* 444. HALL. *Gott.* p. 23. Amaranthus *tricolor*. LINN. *Sp.* 1. BLACKW. *T.* 317. In India. A.

H. Amaranthus erectus, foliis emarginatis. HALL. *l. c.* Amaranthus *lividus*. LINN. *Sp.* 4. Blitum pulchrum, magnum, rectum, rubrum. I. B. *Hist.* 2. p. 966. In Virginia. A.

H. Amaranthus foliis ellipticis, racemis in aliis sessilibus. HALL. *Gott.* p. 24. Amaranthus floribus conglomeratis axillaribus, foliis lanceolatis obrufis. *Gracizans*. LINN. *Sp.* 5. ? Racemi virides. In Virginia. A.

H. Amaranthus procumbens foliis ovatis emarginatis. LINN. *Cliff.* 444. HALL. *Helv.* 176. Amaranthus

ranthus floribus glomeratis lateralibus, foliis ovatis retusis, caule diffuso. *Blitum*. LINN. *Spec.* 6. Glomeres virides. In Europa temperatori. A.

H. *Amaranthus* racemis triandris cylindricis, interruptis, axillaribus, terminalibusque, foliis rhombis obtusis, petiolo longioribus. An *Amaranthus* spicis triandris, glomeratis sessilibus, axillaribus, terminalibusque, foliis rhombis obtusis? LINN. *Cent.* I. *Plant. n.* 95. Videtur, etsi in nostro neque spicae aristatae, neque folia latiora quam longa sint. A.

H. *Amaranthus* racemis cylindricis, pendulis, longissimis. LINN. *Cliff.* 443. HALL. *Gott.* 23. *Amaranthus caudatus*. LINN. *Spec.* 7. *Amaranthus maximus*. C. BAVH. *Pin.* 120. α . viridis. β . maculatus. γ . ruber. &c. In Peru. Persia. Zeylona. A.

H. *Amaranthus* spicis cylindricis lateralibus conjunctis LINN. *Cliff.* HALL. *Gott.* 24. *Amaranthus hybridus*. LINN. *Spec.* 8. *Amaranthus sylvestris maximus* novae Angliae, spicis viridibus. RAIL *Hist.* 101. In Virginia. A.

H. *Amaranthus* racemis cylindricis erectis caule inermi. *Hypochondriacus*. LINN. *Sp.* 9. *Amaranthus sylvestris maximus* novae Angliae, spicis purpureis. TOURN. I. R. H. 238. *Amaranthus latifolius*, spicis purpureis tenuibus. HALL. *Gott.* 24. In Virginia. A.

H. *Amaranthus* racemis cylindricis erectis, axillis spinosis. *spinosus*. LINN. *Sp.* 11. ROYEN. 419. *Amaranthus indicus spinosus*, spica herbacea. HERM. *Lugd. bat.* 31. T. 33. In India. A.

CELOSIA. HALL. Gott. 25.

Achyranthes. LINN. 255.

Androgyna. Calyx Amaranthio similis quinquefolius cum squamis ad basin accessoriis tribus. Stamina connata in annulum. Stylus simplex. Fructus vnilocularis.

H. *Celosia* foliis lanceolatis, pedunculis angulatis, spica ovato-oblonga. *argentea*. LINN. *Spec.* 1. *Amaranthus* spica albescente habitiori. MART. *Cent.* 7. T. 7. Flores magni, iuniores rubri, adulescentes argentei. In America. A.

H. *Celosia* foliis lanceolato-ovatis. LINN. *Cliff.* 43. HALL. *Gott.* 25. *Celosia cristata*. LINN. *Sp.* 2. *Amaranthus* α . spicatus. β . cristatus. Flores rubri & flavi. In India. A.

H. *Celosia* floribus rigidis retroversis. HALL. *Gott.* 25. *Achyranthes* caule erecto, calycibus reflexis spicae adpressis. *aspera*. LINN. *Sp.* 1. *Amaranthus* Sicularis spicatus. BOCC. *Sic.* 16. T. 9. Flores maturi contra caulem reflexi. Circa germen valvulae quinque plumosae. In Sicilia. Ceylona. Jamaica. Fr.

GOMPHRENA. LINN. 279.

Calyx Celosiae similis. In medio tubus cylindraceus ore quinquefido, totidem staminibus clauso. Styli duo. Capsula vnilocularis.

H. *Gomphrena* caule erecto, foliis ovato-lanceolatis, capitulis solitariis, pedunculis diphyllis. *globosa*. LINN. *Sp.* 1. HALL. *Gott.* 348. α . capitulis rubris, β . capitulis argenteis. *Amaranthus* globosus vulgo. In India. A.

BLITVM. LINN. 14. *Chenopodium* morus T.

Habitus fere *Chenopodii*. Calyx trifolius. Stamina vnicam. Calyces conglomerati succulenti maturi baccam referunt.

H. *Blitum* capitellis spicatis lateralibus. *capitatum*. LINN. Sp. 1. *Chenopodium* morus foliis rari-ter ferratis. HALL. Gott. p. 25. Hac quoque pertinet *Blitum Helv.* 176. *Erdbeeren-Spinat*. In Rhætia & Tyrolensi Comitatu. Anglia. A.

H. *Blitum* capitellis sparsis lateralibus. *virgatum*. LINN. Sp. 2. *Chenopodium* morus foliis argute dentatis. HALL. Gott. l. c. In Tataria. A.

AXYRIS. HALL. Comment. Soc. Sc. Gott. T. II. p. 345. LINN. Gen. 429.

Monophyta. Mares dense spicati terminales, obsecuri singuli, stamine vnico. Fem. alares. Calyx exterior bi-aut trifolius, interior monophyllus, quadrifidus, lacer, plumosus, tubae duae, semen compressum.

H. *Axyris* herbacea amentis masculis corymbosis pedunculatis. HALL. Gott. p. 26. & Comm. Gott. 1752. T. II. p. 345. Tab. XV. f. 2. An *Axyris* caule erecto spicis simplicibus. *Amaranthoides* LINN. Spec. 2? Folia ovato-lanceolata punctata. A.

H. *Axyris* foliis ovatis, caule erecto, spicis conglomeratis. *hybrida*. LINN. Spec. 3? Caulis ramosus, lanatus in primis in summitate ramulorum. Folia ovata, inferne incana, villosa, superne punctata; spica marium pedunculata, conglomerata, glomerulis densissimis. A.

CAMPHORATA. HALL. 183. Camphorosma.

LINN. nov. gen. 1042.

H. Camphorata foliis lanceolatis, hirsutis, ciliatis, glomerulis florum laxè spicatis. *Camphorosma. Monspeliaca*. LINN. *Spec.* 1. *Mat. med.* 43. An Camphorata hirsuta. C. BAVH. *Pin.* 485. Glomeruli florum ad alas foliorum; cuilibet flosculo præter foliolum a latere positum maius, calyx quadrifolius, foliis duobus maioribus, glumarum instar carinatis, duobus minoribus, planis, stamina quatuor calyce minora, styli duo calice multo longiora, capsula monosperma, semen lenticulare. Vide quam diversa sit fabrica a caractere Linnaeano. In Gallia Narbonensi. Tataria. P.

CORISPERMUM. LINN. 12.

Flores solitarii alares. Calyx bifolius. Stamen unum, duo, tria. Tubae duae. Semen compressum, marginatum.

H. *Corispermum* floribus lateralibus. *hyssopifolium*. LINN. *Spec.* 1. *Corispermum* floribus alternis. *Cliff.* 3. ROY. 205. *Corispermum* hyssopifolium. JUSSIEV. *Art. Par.* 1712. p. 244. HALL. *Gott.* 26. In Tataria. Monspeli. A.

CERATOCARPUS. LINN. 921.

Monoph. Flores alares. *Mas.* Calyx bipartitus, stamen unicum. *Fem.* Calyx bifolius. Semen compressum bicornè.

H. *Ceratocarpus arenarius*. LINN. *Spec.* 1. p. 969. BUXBAVM. *Art. Petropol.* 1. p. 241. *Ceratocarpum GERBERI*. In arenosis Tatariae. A.

SAL-

SALSOLA. LINN. 275. Kali.

TOVRN. 128.

Calyx quinquefolius. Stylus bipartitus. Capsula globosa. Semen cochleatum. Flores in alas congesti.

H. *Salsola* (herbacea) foliis pungentibus. LINN. *Cliff.* 86. HALL. *Gott.* 22. *Salsola Kali.* LINN. *Sp.* 1. *Kali spinosum cochleatum.* BAVH. *Pin.* 289. In littoribus marinis. A.

CLASSIS VI.

VAGINALES. LINN.

Caulis geniculatus, vaginis per intervalla cinctus. Calyx quinque aut sexpartitus, plerumque coloratus. Stamina quinque ad octo. Tubae duae vel tres. Fructus calyce comprehensus, monospermus, semen triquetrum aut ovato-compressum. Flores in plerisque spicati, aut racemosi terminales, in paucioribus ad alas foliorum congesti.

RHEVM. LINN. 454. Rhabarbarum.

TOVRN. 18.

Calyx sexpartitus. Stamina novem. Tubae tres plumosae. Semen triquetrum, margine membranaceum. Flores paniculati.

H. *Rheum* foliis glabris cordiformibus. HALL. *Gott.* 15. *Rheum Rhaponticum.* LINN. *Spec.* 1. *Mat. med.* 198. *Rhaponticum.* ALPIN. *Exot. p.* 137. BLACKW. 262. & *Offic.* In Thracia. Tartaria. P.

H. Rheum foliis acutis crispis subhirsutis. HALL. *Gott.* 15. Rheum ^{visulatum} *Rhabarbarum*. LINN. *Spec.* 2. *Mat. med.* 197. Rhabarbarum verum *Offic.* In China prope murum & in Sibiria. P.

LAPATHVM. HALL. 168. Rumex.
LINN. 407.

Calyx sexfolius. Stamina sex. Tubae tres plumosae. Semen triquetrum. Sexus variat.

Lapathum acetosum sexu distinctum, foliorum sagittatorum hamis longis & divergentibus. HALL. *Helv.* 168. Rumex floribus dioicis, foliis lanceolato-hastatis. *Acetosella*. LINN. *Sp.* 22. Acetosa arvensis lanceolata. BAVH. *Pin.* 114. In agris lapidosis, rupibus, muris. P.

Lapathum acetosum sexu distinctum, foliorum sagittatorum hamis brevibus. HALL. *Helv.* 168. Rumex floribus dioicis, foliis oblongis sagittatis. *Acetosa*. LINN. *Sp.* 21. Acetosa pratensis. BAVH. *Pin.* 14. BLACKW. 230. α . Acetosa montana maxima. BAVH. *Pin.* 114. β . Acetosa montana arilato rotundo folio. BOCCON. *Sauerampfer*. In prat. α . β . in montibus altioribus. P. ^{in montibus altioribus} *apud molen garr* *alta de Sm. 42159*

H. Lapathum floribus superioribus masculis, inferioribus femininis, foliis cordatis. Lapathum obtusifolium, spica confertissima. HALL. *Helv.* 6. 170. Rumex alpinus. LINN. *Sp.* 3. Lapathum rotundifolium. CLVS. *hist.* 2. 69. In alpibus P.

Lapathum (*Rumex*) floribus monoicis, calycibus fructus vncinatis. LINN. *Cliff.* 139. ROY. 230. Rumex.

Rumex spinosus. LINN. *Spec.* 20. Beta cretica, semine aculeato. BAVH. *Pin.* 118. *Prodr.* 57. Mares superiores longe spicati, feminini flores alares, congesti. In Creta. P.

H. *Lapathum* (*Rumex*) valvulis laevibus, caule arboreo, foliis subcordatis. *Eunaria*. LINN. *Spec.* 15. *Lapathum acetosum* calycibus triphyllis. HALL. *Gott.* 16. *Acetosa* arborescens, subrotundo folio. BLACKW. *T.* 252. f. 3. In Canariis. Fr.

H. *Lapathum acetosum*, valvis exterioribus amplissimis strepentibus. HALL. *l. c.* *Rumex vesicarius* LINN. *Sp.* 16. In Africa. A.

H. *Lapathum acetosum* pedunculis incrassatis bicornibus. HALL. *l. c.* *Rumex* fructibus dentatis calyce reflexis, caput bovinum referentibus. LINN. *Cliff.* 139. *bucephalophoros*. *Sp.* 13. *Acetosa* ocyimi folio, bucephalophoros. COL. *Eschr.* *T.* 150. In Italia & Germania meridionali. A.

Lapathum foliis oblongis, paniculis aphyllis. HALL. *Gott.* 17. α . *Lapathum* folio acuto, crispo. BAVH. *Pin.* 115. β . *Lapathum aquaticum* folio cubitali. *Rumex* LINN. *Sp.* 14. C. B. *Pin.* 116. γ . *Lapathum acutum* f. *Oxylapathum* f. BAVH. *Hist.* 2. p. 983. *Rumex*. LINN. *Sp.* 10. HALLERVS omnes hasce plantas tanquam varietates ad vnam speciem refert. Habitat α . β . in paludosis. γ . in siccoribus. P.

H. *Lapathum*, (*Rumex*) valvulis integerrimis unica granifera, foliis cordato-lanceolatis. *sanguineum*. LINN. *Sp.* 2. *Lapathum* folio acuto rubente. C. BAVH. *Pin.* 115. In Virginia. P.

- + *Lapathum foliis acutis verticillatum.* HALL. *Gott.* 17. *Rumex* ad alas foliorum conglomeratis, foliis lanceolatis integerrimis. BOEH. *Fl. Lips.* 576. *Hydrolapathum minus.* LOBEL. *icon.* p. 286. In fossis. P. *Hab. a. p. n. 1590.*

Lapathum foliis obtusis, ora crispa, calycis valvis interioribus ferratis. HALL. *Gott.* 18. *Rumex obtusifolius.* LINN. *Spec.* 11. *Lapathum vulgare folio obruso.* I. BAVH. Ad vias. P.

H. *Lapathum arvense subhirsutum folio retuso, caule longius brachiato, capsula seminis crenata.* HALL. *Gott.* 18. In TILLII *hort. Pis.* p. 93, T. 37. f. 2. folium pingitur. Caulis ramosissimus ad angulos rectos brachiatus. A.

- + *Lapathum perenne foliis longissimis, calycibus ciliatis.* HALL. *l. c.* *Lapathum anthoxanthum.* I. B. II. 987. In vdis circa Göttingam. In Saxonia superiori. Circa Berolinum, in *Lam. montan.* l. p. 21. 1778.

H. *Lapathum annum foliis lingulatis, verticillis densissimis, ciliis calycis longissimis.* HALL. *l. c.* *Rumex aegyptiacus.* LINN. *Sp.* 6. In Aegypto. A.

BISTORTA. HALL. 178. *Polygonum* *lian.*

Calyx quinquepartitus. Stamina octo. Tubae duae. Semen triquetrum. Radix contorta. Flores dense spicati.

Bistorta foliis ovato-oblongis acuminatis. LINN. *Cliff.* 150. ROY. 217. *Bistorta foliis imis appendiculatis, semine triquetro.* HALL. *Helv.* 178. *Polygonum Bistorta.* LINN. *Sp.* 2. *Bistorta radice* α . magis, β . minus

minus intorta. C. B. *Pin.* 192. Bistorta BLACKW. *T.* 254. & offic. *Krebswurzel, Schlangenwurzel.* In pratis vdis. P.

Bistorta foliis lanceolatis. LINN. *Cliff.* 150. Bistorta foliis ad oram nervosis, imis ovalibus, superioribus linearibus. HALL. *Helv.* 1. 179. Polygonum viviparum. LINN. *Sp.* 3. Bistorta alpina α . media β . minor. In Alpibus. P.

PERSICARIA. HALL. 179. LVDW. 778.

Flores densius spicati. Stamina quinque ad octo. Tuba aut bifida aut duplex. Semen ovato compressum.

Persicaria foliis serratis, spica densa, cylindrica. HALL. *Helv.* 1. 180. Persicaria florum staminibus quinis corollam superantibus, stylo bifido. LINN. *Cliff.* 41. Polygonum amphibium. LINN. *Spec.* 7. Potamogeton falicis folio. BAVH. *Pin.* 193. In fossis folia laevia, & caulis procumbens, in siccioribus locis caulis erectus & folia hirsuta Glandulae ad basin segmentorum.

Persicaria vaginis florum ciliaribus, spicis interruptis densioribus. HALL. *Helv.* 180. Gott. 28. Persicaria floribus hexandris digynis. LINN. *Fl. Succ.* 319. Polygonum Persicaria. LINN. *Sp.* 11. Persicaria mitis. BLACKW. *T.* 118. maculosa, & non maculosa. BAVH. *Pin.* 101. α . Persicaria foliis subtus tomentosis. *Helv.* 181. β . Persicaria angustifolia. C. BAVH. *Prodr.* 43. HALL. *Helv.* n. 11. Flöhkraut. Pfersingkraut. In hortis & ad vias. A.

Perficaria acris, spicis longis strigosis, vaginis non ciliatis. HALL. *Helv.* 181. *Gott.* 29. *Perficaria* florum staminibus senis, stylo bifido. LINN. *Cliff.* 46. *Polygonum Hydripiper*. LINN. *Spec.* 10. *Perficaria vrens*, f. *Hydropiper*. BAVH. *Pin.* 101. BLACKW. *T.* 119. *Wasserpfeffer*. In foliis. A.

H. *Perficaria* florum staminibus sex pluribusque, stylo duplici. LINN. *Cliff.* 42. *Perficaria Orientalis* Nicotianae folio, calyce florum purpureo. TOURN. *Cor.* 38. COMMEL. *rar.* 43. *T.* 43. HALL. *Gott.* 29. *Polygonum orientale*. LINN. *Spec.* 13. Glandulae octo ad basin segmentorum. In Oriente. A.

FAGOPYRVM. HALL. 172. LVDW. 794.

HELXINE LINN.

Stamina octo, tubae tres, glandulae octo ad basin segmentorum. Flores laxè spicati.

H. *Fagopyrum (Helxine)* caule diffuso, foliis lanceolatis. LINN. *Hort. Vps.* 96. *Polygonum divaricatum*. LINN. *Spec.* 18. *Perficaria montana*, angustifolia, racemosa. AMMAN. *Ruth.* 240. p. 187. HALL. *Gott.* 29. Glandulae obscurae. In Sibiria. P.

Fagopyrum caule erecto, fructu ex calyce eminente laevi. HALL. *Gott.* 29. *Helv.* 172. *Polygonum Fagopyrum*. LINN. *Sp.* 23. *Mat. med.* 188. p. 63. *Frumentum Sarracenicum*. MATH. 394. *Heydekorn. Buchweize*. Sponte in Asia. Nobis colitur. A.

H. *Fagopyrum* caule erecto, fructu ex calyce eminente serrato. HALL. *l. c.* *Polygonum Tataricum*. LINN. *Spec.* 22. In Sibiria. Finnia. A.

Fago-

Fagopyrum scandens, femine calyce incluso. HALL. *Helv.* 173. Helxine caule volubili. LINN. *Lapp.* 154. ROYEN. 217. *Polygonum Convolvul.* LINN. *Sp.* 24. *Fagopyrum vulgare scandens*. L. R. N. 511. *Vogelzunge, wild Heydekorn*. Vbi-
que in agris & hortis, A.

Polygonum dumetorum. Hall. L. N. 512. *P. hibernicum*.

POLYGONVM. HALL. 182.

Stamina octo. Tubae tres. Glandulae nullae.
Flores solitarii alares.

Polygonum caule procumbente herbaceo, foliis lanceolatis integerrimis. LINN. *Mat. med.* 186. *Polygonum aviculare*. LINN. *Spec.* 15. *Polygonum vulgare* α . angustifolium β . latifolium. BAVH. *Pin.* 281. *Polygonum* HALL. *Helv.* 182. BLACKW. *T.* 315. *Wegtritt, Weggras*. In cultis & rudera-
tis, A.

h. *Polygonum* foliis lanceolatis, caule stipulis obrecto fruticoso, *maritimum*. LINN. *Sp.* 8. *Polygonum* perenne, vaginis amplissimis. HALL. *Gott.* 30. *Polygonum* *maritimum* latifolium. BAVH. *Pin.* 280. In Europa australi, LINNAEVS inter species pentandras refert. P.

Ad. *Polygonum hibernicum*
in *Pin.* 281. *Polygonum* *maritimum* latifolium. BAVH. *Pin.* 280. In Europa australi, LINNAEVS inter species pentandras refert. P.

HERNIARIA. LINN. 272.

Calyx patens, interne coloratus. Stamina quin-
que sterilia, quinque abortiva. Tuba bifida. Glo-
meruli florum alares.

Herniaria calycibus bractea nudis. LINN. *Mat. med.* 104. HALL. *Helv.* 182. *Herniaria glabra*. LINN. *Spec.* 1. BLACKW. *T.* 320. *Polygonum mi-*

nus.

nus. f. Millegrana maior BAVH. *Pin.* 281. *Bruckkraut*. In arenosis, siccis, v. g. circa *Northheim*, *Herzberg*. A.

H. *Herniaria squamis nitidis flores superantibus*. LINN. *Cliff.* 41. ROY. 215. HALL. *Gott.* 31. *Illecebrum Paronychia*. LINN. *Sp.* 4. *Paronychia hispanica*. CLVS. *Hist.* 2. 183. In Gallia *Narbonensi* & *Hispania*. A.

ILLECEBRVM. LINN. 256.

Calyx coloratus, cartilagineus. Stamina quinque. Tuba simplex. Flores dense verticillati.

Illecebrum floribus verticillatis nudis, caulibus procumbentibus. verticillatum. LINN. *Spec.* 1. *Illecebrum*. HALL. *Gott.* 31. *Illecebra spuria*. RVPP. 99. In arenosis *Collensibus* & circa *Berolinum*. A.

Quae sequuntur tria genera ob habitum hic relinquenda, etsi petalis, numero staminum, aut capsula polysperma differant.

MOLLVGO. LINN. 99.

Calyx interne coloratus. Stamina tria. Tubae tres. Capsula polysperma.

H. *Mollugo foliis quaternis obovatis, panicula dichotoma*. LINN. *Spec.* 2. *Triclis tristemon foliis coniugatis*. HALL. *Gott.* 25. Vid. descriptionem & iconem in *Comment. Soc. Sc. Gott. T. V.* Petala quinque calyce minora. Folia opposita, quae ad ramorum ortum quaterna videntur. In Gallia *Narbonensi*. A.

h. *Mollugo* foliis verticillatis cuneiformibus acutis, caule subdiviso procumbente, pedunculis unifloris. *verticillata*, LINN. *Sp.* 4. *H. Vpf.* 24. Vid. icon floris in *Comm. Gott. T. V.* In *Africa. Virginia. A.*

PHARNACEVM. LINN. 341.

Omnia fere prioris, Differt tantum staminibus quinque & floribus umbellatis.

h. *Pharnaceum* glabrum, pedicellis folia aequantibus. *Cerviana*, LINN. *Sp.* 1. *Phil. Bot.* 258. Tridis pentastemon, floribus umbellatis. HALL. *Comm. Sc. Sc. Gott. T. II. p.* 346. *T. XV. f.* 2. *Rostochii.* In *Sibiria. A.*

CORRIGIOLA. LINN. 340.

Calycis folia & petala & stamina quinque. Tuba trifida. Capsula monosperma. Folia alterna.

Corrigiola caule fasciculis florum terminato. LINN. *Cliff.* 112. *Spet.* p. 271. HALL. *H.* 391. *Polygonifolia.* DILL. *Nov. gen.* 95. *Tab. III.* In arenosis. v.g. *Cellensibus.*

CLASSIS. VII.

TRICOCCAE.

Stamina in multis plura, quam segmenta calycis, in aliis segmentorum numero aequalia. Petala in aliis adsunt, in aliis desunt. Fructus tricoccus, trilocularis, tot styli quot loculi fructus. Sexus variat. Omnes ires, fuspectae.

EUPHOR

EUPHORBIA. LINN. 536. Tithymalus.

HALL. 189. TOVRN. 18.

Ex margine calycis monophylli petala quatuor aut quinque, crassa, plana. Fructus ex flore pendulus, petiolatus. Styli tres bifidi. Vbique lacte turget.

h. Euphorbia aculeata nuda subquingularis, aculeis geminatis. *Canariensis*. LINN. Sp. 2. Euphorbium tetragonum & pentagonum spinosum Canarium. BOERH. p. 258. HALL. Gott. 32. Euphorbium. BLACKW. T. 340. f. 1. In Africa calidiori. Fr.

h. Euphorbia aculeata nuda multangularis, spinis solitariis subulatis. *cereiiformis*. LINN. Sp. 5. Euphorbium cerei effigie, caulibus gracitioribus. BOERH. Lugdb. 1. p. 258. In Aethiopia. Fr.

h. Euphorbia aculeata nuda multangularis, aculeis geminatis. *officinarum*. LINN. Sp. 6. Euphorbium polygonum spinosum, cerei effigie ISNARD. Act. Par. 1720. T. 10. BLACKW. T. 340. f. 2. In Africa calidiorē. Fr.

h. Euphorbia inermis imbricata tuberculis, foliolo lineari instructis. *caput Medusae*. LINN. Sp. 8.

α. Euphorbium erectum, aphyllum, ramis rotundis, tuberculis tetragonis BURM. T. 7. f. 2. H. Gott. 32.
β. Tithymalus aizoides africanus, simplici squamoso caule. COMMEL. Prael. 58. T. 8. In Aethiopia. Fr.

h. Euphorbia inermis nuda fruticosa, filiformis volubilis: cicatricibus oppositis. *viminalis*. LINN. Sp. 9. Felfel Tavil, ALP. Aegypt. 190. T. 190. In Africae maritimis. Fr.

H.

h. *Euphorbia inermis* seminuda fruticosa, filiformis, flaccida, foliis alternis. *Mauritanica*. LINN. Sp. 10. *Tithymalus aphyllus Mauritaniae*. DILL. Elth. T. 289. In *Africae* maritimis. Fr.

h. *Euphorbia inermis* subnuda fruticosa filiformis erecta, ramis patulis determinate confertis. *Tirucalli*. LINN. Sp. 11. *Tithymalus indicus frutescens*. COMMEL. H. Amst. l. p. 27. T. 14. In *India*. Fr.

h. *Euphorbia umbella* quadrifida, dichotoma, foliis oppositis integerrimis. *Lathyris*. LINN. Sp. 31. *Tithymalus*. HALL. Helv. n. 1. 189. *Lathyris* f. *Cataputia minor*. 1. B. III. p. 880. In *Helvetia*, *Gallia*, *Italia*. P.

Euphorbia (*Tithymalus*) foliis subhirsutis, ad caulem ellipticis, cymbis florum perfoliatis, petalis lunatis acutis. HALL. Helv. 190. *Euphorbia sylvatica*. LINN. Sp. 55. *Tithymalus sylvaticus lunato flore*. BAVH. Pin. 290. In sylvis. P.

h. *Euphorbia involucellis* perfoliatis emarginatis, petalis obtusis foliis integerrimis, hirsutis, caule frutescente. *Euphorbia Characias*. LINN. Sp. 56. *Tithymalus characias* L. GLVS. Hist. 2. 88. In *Gallia*, *Hispania*. Fr. *inter L. uliginosam et hirsut. inveniuntur. a. 3. Charac.*
(Haupt. 1803. 2. p. 167.)

Euphorbia (*Tithymalus*) altissimus, levis, foliis sub ^{ternis} foliis ternis, cordatis, fructu verrucoso. HALL. Helv. 191. n. 6, Gott. 34. *Euphorbia palustris*. LINN. Sp. 51. *Tithymalus palustris fruticosus*. BAVH. Pin. 292. P. In pratis paludosis *Thuringiae Helvetiae*

Euphor-

Euphorbia (*Tithymalus*) foliis glabris, ferracuminatis, involucris bifoliis, cordatis, fructu to verrucoso. *Helv.* 191. *E. platyphyllus*. LINN. 47. *Tithymalus arvensis latifolius Germanicus* Pin. 191. Vbique in agris. A.

H. Euphorbia (*Tithymalus*) foliis ovato-lanceolatis, subhirsutis, umbellae involucri quinquefo HALL. *Gott.* 34. Num Varietas?

H. Euphorbia (*Tithymalus*) foliis ovatis, fterminatis, subhirsutis, petalis longe bicorni HALL. *Gott.* l. cit. A.

H. Euphorbia (*Tithymalus*) foliis ellipticis, nula terminatis, petalis lunatis, ferratis. HALL. 35. *E. Pithyusa*. LINN. Sp. 35. *Tithymalus f brevis aculeatis*. BAVH. Pin. 292. In arce Belgii, Hispaniae. P.

H. Euphorbia (*Tithymalus*) foliis linearibus nula terminatis, radiis umbellae plurimis r f petalis obtusis HALL. *Gott.* l. c. *Helv.* 192. n. *E. paralias*. LINN. 37. *Tithymalus marit* BAVH. Pin. 291. In maritimis. P.

Euphorbia (*Tithymalus*) foliis linearibus, o fis, petalis emarginatis. *Helv.* l. c. *Tithymalus an daloides angustifolius*. I. R. H. TABERN. p. 5 RVPP. p. 273.

Euphorbia (*Tithymalus*) foliis linearibus, obt confertis, petalis lunatis. HALL. *Helv.* 193. n. *E. cyparissias*. LINN. Sp. 49. Rarius circa *Gott gam* v. g. circa *Grossen Schneen*.

Euphorbia (Tithymalus) foliis linearibus obtusis, petalis argute bicornibus. *Helv.* 193. n. 12. *Esula exigua*. LINN. *Sp.* 29. c. BAVH. *Pin.* 291. Vbi-
que in hortis. A.

Euphorbia (Tithymalus) foliis subrotundis serratis, radiis umbellae quinque latissimis. *Helv.* 194. n. 14. E. *Helioscopia*. LINN. *Sp.* 40. *Tithymalus helioscopius*. BAVH. *Pin.* 291. In cultis & hortis. A.

Euphorbia (Tithymalus) foliis ovatis, involucri bifoliis cordatis, obtusissimis. *Helv.* 193. n. 13. E. *Peplus*. LINN. *Spec.* 27. *Peplus* f. *Esula rotunda*. BAVH. *Pin.* 292. A.

H. *Euphorbia* foliis subrotundis obtusis, floribus solitariis in foliorum alis. GVETT. *Stamp.* 2. p. 240. *Euph. Peplis*. LINN. *Sp.* 23. *Peplis maritima folio obtuso*. BAVH. *Pin.* 293. *Peplis* CLVS. *Hist.* 2. p. 187. Petala argute bicornia, fructus in dorso varum verrucosus. In *Narbonae* maritimis. A.

H. *Euphorbia* dichotoma, foliis serratis oblongis, floribus axillaribus solitariis, ramis patulis. *maculata*. LINN. 21. Synonymum PLVCK. 372. T. 65. f. 8. ad sequentem potius retulerim. Folia inferiora maculata, superiora pura, petala obtusa, fructus glaber, laevis. In *America* septentrionali. A.

H. *Euphorbia* dichotoma, foliis crenulatis oblongis, floribus solitariis axillaribus, caulis procumbentibus. *Chamaesyce*. LINN. *Sp.* 22. *Chamaesyce*. BAVH. *Pin.* 293. CLVS. *Hist.* 2. p. 187. Habitus prioris, sed tota planta omni ex parte minor, & hirsuta, & procumbens. Folia semper maculata
D prioris

prioris spec. similia, petala integra, fructus hirsutus.
In *Sibiria* & *Europa* australi. A.

CLVTIA. LINN. 1009.

Pl. Diphyta. In utroque sexu calyx quinquefolius, quinque petala, & nectaria intra petala posita. *Mar.* Stamina quinque ex columella media orta, patentia. *Fem.* Styli tres, bifidi. Capsula sexsulca.

H. *Clusia* foliis ovatis integerrimis, floribus lateralibus. *pulchella*. LINN. *Sp.* 2. *Clusia foliis petiolatis*. Cliss. 500. HALL. *Gott.* 36. In *Aethiopia*. Fr.

CNEORVM. LINN. 47.

Calyx tridentatus. Petala tria, totidemque stamina. Stylus trifidus.

H. *Cneorum tricocon*. LINN. *Sp.* 1. ROY. 119. *Chamalaea tricoceps*. BAVH. *Pin.* 462. HALL. *Gott.* 36. In *Hispania*, *Gallia Narbonensi*. Fr.

PHYLICA. LINN. 236.

Plurimi flores in discum collecti. Calyx proprius trifolius. Petalum limbo quinquefido; squamae stamina munientes. Stylus simplex.

H. *Phyllica* foliis ovato-linearibus. ROY. 199. HALL. *Gott.* 36. Ph. *ericoides*. LINN. *Sp.* 1. *Alternoides Africanae ericae foliis, floribus albicantibus & muscosis*. COMMEL. *Hort. Amst.* II. p. 1. T. 1. In *Aethiopia*. Fr.

CROTON.

CROTON. LINN. 960.

H. Croton foliis ovatis tomentosis integris serratis. *argenteum*. LINN. *Sp.* 4. *Cliff.* 441. ? Flores in corymbo, superius masculo, inferius feminino. Calyx in *femina* multo maior ab exteriori latere tripartitus in folia oblonga tomentosa, quibus singulis ad basin squama glandulifera adhaeret, ab interiori fissus in duas lacinias minimas. Petala nulla vidi. *Fl. mas* multo minor, calyx similis, filamenta staminum hirsuta ex receptaculo hirsuto, vix coalita. Conf. *epist.* ad III. WERLHOF, ubi fusus hanc descripsi.

JATROPHA. LINN. 961.

Monoph. *Mas.* Petalum hypocrateriforme, limbo quinquefido. Stamina decem. *Fem.* Corolla pentapetala. Styli dichotomi.

H. Jatropha foliis cordatis angulatis. ^{*Currat*} LINN. *Sp.* 3. *Mat. med.* 439. HALL. *Gott.* 36. In *America* calidiori. Fr.

RICINUS. LINN. 962.

Monoph. *Mares* inferius positi. Calyx quinquepartitus. Filamenta staminum inferne ramosae coalita. *Fem.* Calyx tripartitus. Styli bipartiti,

H. Ricinus foliis peltatis serratis: petiolis glanduliferis. LINN. *Cliff.* 4501 *Mat. med.* 438. HALL. *Gott.* 37. Ric. *communis*. LINN. *Sp.* 1. *Ricinus vulgaris*. BAVH. *Pin.* 439.

Var. *α.* *Ricinus amer. fructu glabro.*

β. *Ricinus caule rutilante.* Sponte in *Africa*, *Asia*, *America* & *Europa* calidiori. P. nobis annua.

prioris spec. similia, petala integra, fructus hirsutus.
In *Sibiria* & *Europa* australi. *A.*

CLVTIA. LINN. 1209.

Pl. Diphyta. In utroque sexu calyx quinquefolius, quinque petala, & nectaria intra petala posita. *Mas.* Stamina quinque ex columella media orta, parentia. *Fem.* Styli tres, bifidi. Capsula sexsulca.

H. *Clutia* foliis ovatis integerrimis, floribus lateralibus. *pulchella*. LINN. *Sp.* 2. *Clutia foliis petiolatis*. *Cliff.* 500. HALL. *Gott.* 36. In *Aethiopia*. Fr.

GNEORVM. LINN. 47.

Calyx tridentatus. Petala tria, totidemque stamina. Stylus trifidus.

H. *Cneorum tricocton*, LINN. *Sp.* 1. ROY. 119. *Chamalaea tricoctos*. BAVH. *Pin.* 462. HALL. *Gott.* 36. In *Hispania*, *Gallia Narbonensi*. Fr.

PHYLICA. LINN. 236.

Plurimi flores in discum collecti. Calyx proprius trifolius. Peralum limbo quinquefido; squamae stamina munientes. Stylus simplex.

H. *Phylica* foliis ovato-linearibus. ROY. 199. HALL. *Gott.* 36. Ph. *ericoides*. LINN. *Sp.* 1. *Alaternoides Africana ericae foliis, floribus albicantibus & muscosis*. COMMEL. *Hort. Amst.* II. p. 1. T. 1. In *Aethiopia*. Fr.

CROTON.

CROTON. LINN. 960.

H. *Croton* foliis ovatis tomentosis integris serratis. *argenteum*. LINN. *Sp.* 4. *Cliff.* 444. Flores in corymbo, superius masculo, inferius feminino. Calyx in *femina* multo maior ab exteriori latere tripartitus in folia oblonga tomentosa, quibus singulis ad basin squama glandulifera adhaeret, ab interiori fissus in duas lacinias minimas. Petala nulla vidi. *Fl. mas* multo minor, calyx similis, filamenta staminum hirsuta ex receptaculo hirsuto, vix coalita. Conf. *epist.* ad III. WERLHOF, ubi fustus hanc descripsi.

JATROPHA. LINN. 961.

Monoph. *Mas*. Petalum hypocrateriforme, limbo quinquefido. Stamina decem. *Fem.* Corolla pentapetala. Styli dichotomi.

H. *Jatropha* foliis cordatis angulatis. ^{*Currae*} LINN. *Sp.* 3. *Mat. med.* 439. HALL. *Gott.* 36. In *America* calidiori. Fr.

RICINUS. LINN. 962.

Monoph. *Mas* inferius positi. Calyx quinquepartitus. Filamenta staminum inferne ramosse coalita. *Fem.* Calyx tripartitus. Styli bipartiti,

H. *Ricinus* foliis peltatis serratis: petiolis glanduliferis. LINN. *Cliff.* 450. *Mat. med.* 438. HALL. *Gott.* 37. Ric. *communis*. LINN. *Sp.* 1. *Ricinus vulgaris*. BAVH. *Pin.* 439.

Var. *a.* *Ricinus amer. fructu glabro.*

β. *Ricinus caule rutilante.* Sponte in *Africa, Asia, America & Europa* calidiori. P. nobis annua.

ACALYPHA. LINN. 959.

Monoph. *Mares* plurimi in iulum fere conferti, calyce minimo quadrifolio, stamina vt in Ricino. *Feminae* plures involucro foliaceo magno crenato exceptae. Styli ramosi.

H. *Acalypha* involucris femineis cordatis incisis, foliis ovato lanceolatis petiolo longioribus. *Virginica*. LINN. Sp. 1. *Mercurialis tricocos hermaphroditica, ad foliorum iuncturas e foliolis cristatis tulifera simul & fructum ferens*. BURM. *Ceylan*. 248. T. 99. fig. 4. Sponte in *Ceylona*. *Virginia*. A.

CLIFFORTIA. LINN. 1004.

Diph. *Mas*. Calyx trifolius. Stamina plurima. *Fem*. Calyx similis germiui insistens. Fructus bilocularis.

H. *Cliffortia* foliis subcordatis dentatis. *ilicifolia*. LINN. Sp. 1. *Hort. Cliff*. 463. T. 30. *Arbustula Afra folio acuto ilicis caulem amplexo rigido*. DILL. *Elth*. 36. T. 31. f. 38. In *Aethiopia*. Fr.

MERCVRIALIS. LINN. 998.

HALL. 194.

Diph. *Mares* in iulos congesti, calyce trifolio, staminibus duodecim circiter. *Feminae* solitariae. Styli duo reflexi. Capsula bilocularis.

Mercurialis caule simplicissimo, foliis scabris, *perennis*. LINN. Sp. 1. HALL. *Helv*. 194. 2. Gott. 37. *Mercurialis montana* α . *testiculata* β . *spicata*. BAVH. *Pin*. 122. In sylvis primo vere. P.

TRICOCCAE.

53

Mercurialis caule brachiato, foliis glabris. annua.
LINN. *Spec.* 2. *Mat. med.* 464. HALL. *Helv.* 194.
1. Mercurialis BLACKW. *T.* 168. & *offic. Bingel-*
kraut. In ruderatis, vmbrosis. A.

Quae sequuntur tria genera, partibus quibusdam
anomala, ob habitum tamen hic relinquenda.

THELYGONVM. LINN. 947.

Pl. Monoph. *Mas.* Calyx bifolius stamina duo-
decim. *Fem.* Fructus globosus unilocularis, stylus
simplex.

H. Thelygonum Cynocrambe. LINN *Sp.* I. HALL.
Ger. 38. Cynocrambe Dioscoridis. BAVH. *Pin.* 122.
Pedr. 59. In *Italia, Monspeli.* P.

ASARVM. LINN. 522. HALL. 195.

Calyx monophyllus, semitrifidus. Stamina duo-
decim, stylus stigmate sexfido. Fructus sexlocularis
sub calyce.

Asarum foliis reniformibus obtusis. Europaeum.
LINN. *Sp.* I. *Mat. med.* 219. Asarum. HALL. *Helv.*
195. BLACKW. *T.* 383. & *offic. Haselwurz.* In sylvis. P.

ARISTOLOCHIA. LINN. 911. HALL. 196.

Petalum ex tubo ventricoso lingulatum, sub flo-
re capsula sexlocularis; stylo brevi, sub stigmate sex-
partito stellato, adnascuntur sessiles antherae sex
quadriloculares.

H. Aristolochia foliis cordatis, caule erecto, flori-
bus axillaribus confertis. LINN. *Sp.* 13. Aristolochia
Clematitis. BLACKW. 255. vulgaris CLVS. *Hist.* 2.
LXXI. Sponte in *Asia, Helvetia, Gallia.* P.

H. Aristolochia caule infirmo ramoso, foliis cordatis crenulatis petiolatis, floribus solitatis ^{Pistolochia} LINN. Spec. 10. Aristolochia *Pistolochia dicta*. BAVH. Pin. 307. In Hispania. Gallia Narbonensi. P.

CLASSIS VIII.

INVNDATAE LINN.

Caulis herbaceus, geniculatus. Folia verticillata, aut opposita. Flores aut in alis foliorum, aut in spicam digesti. Stamina vnicum ad octo. Semina saepe nuda, toridemque semper tubae. Omnes in paludosis aut stagnis; perfecta florescentia sub aqua latent.

SALICORNIA. LINN. 10.

Habitus aphyllus. Flores in conum congesti. Calyx truncatus. Stamen vnicum. Stylus apice bifido. Semen vnicum.

H. *Salicornia* articulis apice crassioribus obtusis *Europaea*. LINN. Sp. 1. Mat. med. 8. a. annua. *S. sempervirens*. *Salicornia* HALL. Gott. 5. *Kali geniculatum* BAVH. Pin. 289. Soda offic. Ad aquas salas & in littoribus marinis.

CHARA. LINN. 1066.

Folia verticillata in quorum alis anthera sessilis. Semen vnicum.

Chara caulibus laevibus. *vulgaris*. LINN. Spec. 2. p. 1156. ROY. 533. *Chara vulgaris foetida*. VAILL. HALL. Gott. 38. *Helv.* 196. n. 1. In aquis pigris.

LIMNO-

LIMNOPEUCE. HALL. 197. Hippuris

LINN. II.

Folia verticillata, in quorum alis semen vnicum & stamen vnicum.

Limnopeuce. HALL. *Gott.* 38. *Helv.* 197. Hippuris vulgaris. LINN. *Spec.* I. ROY. 205, In rivis & stagnis. A.

ZANNICHELLIA. LINN. 920.

HALL. 198.

Folia coniugata, in eadem ala folii, flos *mas* stamine vnico, & *femina* seminibus quatuor nudis, stigmate lato patente, calyce communi exceptis.

Zannichellia tubae ligula integra. HALL. *Helv.* 198. *Gott.* I. c. Zannichellia palustris. LINN. *Sp.* I. Vid. iconem MICHEL. *gen.* 71. T. 34. f. 1. 2. In rivo ad *Rasemühle*. A.

STELLARIA. HALL. 198: Callitriche,

LINN. 13.

Folia coniugata. Flores alares, *mares* stamine vnico, *feminae* fructu biloculari. Vtrique sexui petala duo lunata.

Stellaria foliis omnibus angustis, apice resecto. HALL. *Helv.* 199. n. 1.

β. Stellaria foliis ad caulem angustis, in summitate subrotundis. HALL. I. c. n. 2.

γ. Stellaria foliis omnibus subrotundis. HALL. n. 3. Callitriche palustris LINN. *Sp.* I. omnes, ut varietates, iungit. In paludibus. A.

POTAMOGETON, LINN. 160.

-HALL. 199.

Flores spicati ex alis foliorum. Petala quatuor, totidemque stamina antheris gemellis, & semina nuda quatuor.

Potamogeton foliis oblongo-ovatis petiolatis, *natans*, LINN. *Sp.* 1. HALL. *Helv.* 1. c. n. 1. *Potamogeton rotundifolium*, BAVH. *Pin.* 193. In fluviis & stagnis.

Potamogeton foliis ovatis longissime petiolatis. *Helv.* 199. *Potamogeton foliis lanceolatis planis in petiolos desinentibus. lucens*, LINN. *Sp.* 4. *Fontinalis lucens maior*, BAVH. *Hist.* III. p. 769. In stagnis. v. g. *Lichtenmeer auf dem Heimberg*.

Potamogeton foliis cordatis amplexicaulibus, *perfoliatum*, LINN. *Sp.* 2. HALL. *Helv.* 199. n. 3. *Potamogeton rotundifolium alterum*, LOESEL. *Pruss.* ic. 65. In lacubus, v. g. *lacu Seeburgensi ad Baerenshausen*.

Potamogeton foliorum oris plicatis, molliter spinosis, *Helv.* 202. n. 5. *Potamogeton crispum*, LINN. *Sp.* 5. *Fontinalis crispa*, 1. B. *Hist.* III. p. 778. *Potamogeton foliis crispis f. Lactuca ranarum*, BAVH. *Pin.* 193. In rivis v. g. *am Schützenhof*.

Potamogeton foliis lanceolatis oppositis acuminatis, *setaceum*, LINN. *Sp.* 2. HALL. *Helv.* 200. n. 6. *Fontinalis media lucens*, 1. B. III. p. 777. In aquis lente fluentibus.

Potamogeton foliis planis longissimis, spica densa & crassa, HALL. 1. c. n. 8. *Potamogeton racemosum angu-*

angustifolium. C. B. *Prodr.* p. 101. n. 3. In rivo
am Schützenbause & alibi.

Potamogeton foliis lineari-lanceolatis alternis
sessilibus, stipula latioribus. *gramineum*. LINN. *Sp.*
10. Potamogeton HALL. *Helv.* n. 7. Potamogeton
gramineum latifolium. LOESEL. *icon*, 66. p. 206.
In fluviis lente fluentibus vt in *Leina*.

Potamogeton foliis gramineis vndulatis spicis exi-
guis. HALL. *l. c.* n. 9. P. *compressum*. LINN. *Sp.* 7.
Potamogeton *caule compresso, folio graminis candi-*
ni. RAI. *Hist.* 189. In fossis paludosis.

Potamogeton pedunculis florigeris longissimis,
glomerulis florum remotis. HALL. *l. c.* n. 11. Po-
tamogeton *marinum*. LINN. *Sp.* 11. Potamogeton
pufillum. BOCCON. *Sicil.* 42. T. 20. f. 5. Circa
Gottingam.

MYRIOPHYLLVM. LINN. 945. Pentapteris,

HALL. 201.

Monophyta. *Mas.* Calyx quadrifolius, stamina
octo. *Fem.* Capsula tetrasperma, tubis quatuor.
Interdum & androgyni adfunt. Folia verticillata.

Myriophyllum (*Pentapteris*) scapo florifero nu-
do. HALL. *Gott.* 40. *Myriophyllum spicatum*.
LINN. *Sp.* 1. Myrioph. *aquaticum pennatum spi-*
catum. C. BAVH. *Prodr.* 73. T. 73. Flores mascu-
li verticillati quaterni, staminibus, calyce explicato,
longe pendulis, feminini calyce trifolio mihi ob-
servati sunt. In lacu *Seeburgenfi*.

o Myriophyllum (*Pentapteris*) floribus in alis sessilibus. *Helv.* 202. n. 2. *M. verticillatum*. LINN *Sp.* 2. Millefolium aquaticum flosculis ad foliorum nodos. BAVH. *Pin.* 141. In fossis v. g. am Schützenbaufs.

CERATOPHYLLVM. LINN. 244.

HALL. 202.

Pl. monophyta. *Mas.* Calyx multipartitus, stamina plurima. *Fem.* Calyx similis, capsula monosperma, acuminata. Folia verticillata, flores alares.

Ceratophyllum foliis asperis, bis bifurcatis. HALL l. c. *C. demersum*. LINN. *Sp.* 1. Dichotophyllum DILL. *nov. gen.* p. 100. T. 3. In fossis.

CLASSIS. IX.

GRAMINEAE.

In omnibus amplissimae huius classis plantis caulis, qui *culmus* dicitur, est simplex, articulatus foliis vaginantibus cinctus; folia longa sunt, angusta, integerrima, sine acuto: flores in summitate caulis aut in spicam aut in paniculam dispositi, simplices, aut in capitulum, quod *torus* dicitur, collecti: calyx & folliculus glumofus; ut plurimum bivalvis; stamina in plurimis tria, filamentis capillaribus, antheris sagittatis; ex apice germinis stylus duplex aut simplex: stigmata plumosa. Semen unicum folliculo comprehensum.

ORDO. I.

GRAMINA.

Calyx biglumis, rarius vniglumis, & triglumis.
 Folliculus bivalvis, (rarius vnivalvis,) valvula interiori planiori, exteriori concaua, saepe aristata.
 Stamina tria, (excepto *Anthoxantho*, cui duo,) styli duo, semen vnicum.

I. Locusta vniflora.

ANTHOXANTHVM. LINN. 40.

Calyx & folliculus bivalvis, hic aristatus, nectarium diphyllum. Stamina duo.

Anthoxanthum spica ovato-oblonga, flosculis subpedunculatis arista longioribus, *odoratum*, LINN.
Spec. I. Gr. pratense spica flavescente, BAVH. *Pin.*
 3. In ficcis.

NARDVS. LINN. 65.

Calyx nullus, folliculus bivalvis. Stylus simplex.
 ex LINN.

Nardus spica setacea recta, *stricta*, LINN. *Sp. I.*
Gramen sparteum iuncifolium, BAVH. *Pin. 5.* In
Sylva bercynica. v. g. um die Oderbrücke.

PHALARIS. LINN. 69.

Calyx biglumis; folliculus bivalvis; panicula spiciformis.

Phalaris panicula subovata, glumis carinatis.
Canariensis, LINN. *Sp. I. α.* *Phalaris maior sermone albo aut nigro*. B. *Pin. 28. β.* *Gramen spicatum perenne*,

perenne, semine miliaceo, fibrosa radice. HALL. Gott. 51. In Europa australi. Canariis.

Phalaris panicula oblonga ventricosa. arundinacea. LINN. Sp. 3. Gramen *arundinaceum spicatum.* B. Pin. 6. Th. 94. Gr. *aquat. paniculatum phalaridis semine.* ^{non. fol. 2. 1524.} SCHEVCHZ. HALL. Gott. p. 46. In stagnis & fossis.

Phalaris panicula cylindrica. phleoides. LINN. Sp. 11. Gramen *typhoides asperum primum.* C. B. Pin. 4. HALL. Gott. 50. In versuris agrorum. ^{Nov. Fl. Soc. 14.}

PHLEVM. LINN. 71. Typhoidis spec.

HALL.

Calyx biglumis, apice bicuspidate, glumis superne latis aristatis. Folliculus bivalvis. Spicatum.

Phleum spica cylindrica longissima. pratense. LINN. Sp. 1. Gramen *typhoides maximum, spica longissima.* C. B. Pin. 4. In pratis & agris.

^{*Phleum nodosum*}
β. Gramen *typhoides asperum alterum.* C. B. Th. p. 32. Caulis fere bulbosus. Num varietas?

^{*Phleum spica cylindrica basi scabro, culmo adscensu, inde bulbosum. det. p. 1. 177. p. 59.*}

ALOPECVRVS. LINN. 72. HALL.

Gott. 42.

Calyx biglumis. Folliculus univalvis, ex basi arista. Spicatus.

^{*glumis villosis*}
Alopecurus culmo spicato erecto. pratense. LINN. Sp. 1. Gramen *spicatum, spica cylindracea longiore tenuissima.* SCHEVCHZ. 69. In pratis v. g. *auf dem Rosdorffer Weg.*

^{*glumis villosis*}
Alopecurus culmo spicato erecto, glumis mediis aristatis. Spec. Edis. 2 p. 69. culm. spicato in horto
Simulacrum pingu. Chy. haufen. Junio 1766. In via
in Rosdorff 1774. In 1775 vixit. HALL. 1785.
785.

Alopecurus culmo spicato infracto, geniculatus, LINN. *Sp.* 2. *Alopecurus culmo infracto aristis gluma longioribus*, ROY. 54. *Gramen aquaticum geniculatum spicatum*, C. B. *Pin.* 3. In fossis natatile.
Alopecurus panicum V. d. in f. a.

PANICVM, LINN. 70. HALLERI *Panici*

& *Milii species*, *Millot. & xylot.*

Calyx triglumis, gluma tertia minima, post alteram posita. Folliculus bivalvis interdum aristatus. Flores spicati aut paniculati.

H. *Panicum spica composita, aristis flosculo brevioribus*, LINN. *Hort. Vrs.* 19. *Panicum italicum*, LINN. *Spec.* 4. *Panicum italicum f. panicula major*, B. *Pin.* 27. In Indiis. A.

Panicum spica tereti, involucellis bifloris fasciculato-pilosis, glaucum, LINN. *Sp.* 2. Var. *β. Gram. panicum f. panicum sylvestre, spica simplici*, SCHEVCHZ. 46. In agris, exeunte aestate.

Panicum spicis alternis remotis laxis compositis, LINN. *Cliff.* 27. *Pan. Crus Galli*, LINN. *Sp.* 5. Gr. *panicum, spica divisa*, C. B. *Pin.* 8. Ad fossas.

Panicum spica composita, aristis spica longioribus, ROYEN. 55. *Cynosurus panicum*, LINN. *Sp.* 8. *Gram. panicum spica aspera*, B. *Pin.* 8. In agris cultis.
Alopecurus panicum Dinter in *Spec. Ed. 2. p. 10. f. 116.*
panicum *Strob. V. d. in f. a.*

Panicum spicis aggregatis, basi interiori nodosis, flosculis geminis muticis, vaginis foliorum punctatis, sanguinale, LINN. *Sp.* 8. *Gram. dactylon folio latiore*, C. B. *Th.* 119. HALL. *Gott.* 146. In segetibus auf dem Hagen.

H. Panicum panicula laxa flaccida, foliorum vaginis pubescentibus. *Miliaceum*. LINN. Sp. 16. *Mat. med.* 34. Milium semine luteo & albo. B. Pin. 26. In India. A nostris colitur.

MILIVM. LINN. 73.

Calyx & folliculus bivalvis, muticus, valvulis ovatis. Stigmata penicilliformia. Paniculatum.

Milium floribus dispersis. *effusum*. LINN. Sp. 1. Gramen sylvaticum; panicula miliacea sparsa. C. B. Pin. 8. In Europae umbrosis, nemoribus.

AGROSTIS. LINN. 74.

Locusta acuminata. Calyx & folliculus bivalvis, aristatus. Paniculata.

Agrostis petalo exteriori exferente aristam rectam strictam longissimam. *Spica vent.* LINN. Spec. 1. Gram. segetum panicula sparsa. C. B. (*Bromus*.) HALL. Helv. 229. Inter segetes.

Agrostis panicula inferiori verticillatim laxa, superiori contracta. LINN. Lapp. 46. Agr. rubra. LINN. Sp. 4. Gram. serotinum arv. panicula contracta pyramidalis. (*Arundo*.) HALL. Gott. 46. Bey Stollberg.

Agrostis panicula tenuissima. LINN. Lapp. 45. Agr. capillaris. LINN. Sp. 8. Gram. mont. panicula spadicæa delivatiore. B. Prodr. 12. (*Poa*.) HALL. Gott. 46. In pratis & locis siccis.

LAGURVS. LINN. 86.

Calyx biglumis, glumis arista villosa. Folliculi valvula maior arista triplici, media e dorso longissima. Spicatus.

H. Lagurus spica ovata. ^{acutiuscula} LINN. Sp. 1. Gram. *alopecuroides spica rotundiore*. BAVH. Pin. 4. (Typhoides) HALL. Gott. 36. In Europa Australi. A.

II. Locusta multiflora.

AIRA. LINN. 75. *Helv.*

Calycis biflori glumae ovato-acutae. Folliculus calyci similis. Fl. paniculati.

Aira panicula spicata, floribus muticis pedunculo longioribus, glumis acuminatis inaequalibus. *tristata*. LINN. Sp. 3. Gram. *spica tristata subirsutum*. C. B. Pin. 2. Prodr. 8. (Poa) HALL. Gott. 46. In siccioribus v. g. Auf dem Heymberg.

Aira foliis planis, panicula patente. floribus muticis laevibus calyce longioribus. *aquatica*. LINN. Sp. 6. Gramen *aquaticum miliaceum*. SCHEVCHZ. 218. HALL. *Helv.* 218. In aquosis. v. g. ad viam versus Geismar.

Aira foliis planis, panicula patente, petalis basi villosis aristatisque; arista recta brevi. *cespitosa*. LINN. Sp. 8. Gramen *segetum panicula arundinacea*. C. B. Pin. 3. In agris. B.

Aira foliis setaceis, panicula angustata, flosculis basi pilosis aristatis; arista tortili longiore. *montana*. LINN. Sp. 14. Gram. *avenaceum capillaceum*, minoribus

noribus glumis. C. B. Pin. 10. (*Avena*.) H
Helv. 228. In sylvis super Hardeggen. B.

(Aira flore masculino aristato, feminino mut)
LINN. Cliff. 27. *Holcus lanatus*. LINN. Sp.
Gram. pratense paniculatum molle. C. B. Prodr.
In vdis.) *Scribitur etiam p. 63.*

H. *Holcus lanatus* 1752.
MELICA. LINN. 76.

Calyx & folliculus bivalvis, muticus. Corpusculum
inter flosculos. Paniculata.

den unistrom
in sp. f. p. 10.
in sp. f. p. 10.
in sp. f. p. 10.
Melica petalis imberbibus, panicula nutante
plicis. *nutans*. LINN. Sp. 2. Gram. montanum
naceum, locustis rubris. C. B. Prodr. 20. In ri
bus. Vm Pless, & in M. Hengst.

H. *Melica altissima* In Sibira Canada
altissima 1752. 113.
POA. LINN. 77. *Dactylis*.

Locusta ovato-oblonga mutica, disticha. C.
& folliculus ut in Aira. Fl. Paniculati.

Poa flosculis confertis vno versu dispositis
Cliff. 28. *Dactylis glomeratus*. LINN. Sp. 2.
spicatum, folio aspero. C. B. Prodr. 9. HALL. H
231. Ad vias.

Poa panicula diffusa, spiculis sexfloris linearibus
aquatica. LINN. Sp. 1. Gram. palustre panicula
altissimum. C. B. Pin. 2. Im Denckerhäuser-Te

Poa panicula diffusa, spiculis quinquefloris
pubescentibus, culmo erecto tereti. *trivialis*. L.
Sp. 3. Gram. pratense paniculatum medium. C.
Pin. Ubique in pascuis. *napula infra panicea aquatica*

Poa panicula diffusa, spiculis quadrifloris pu
bescentibus, culmo erecto tereti. *angustifolia*. L.

Sp. 4. Gr. pratense paniculatum maius angustiori folio. C. B. Pin. 2. Prodr. 6. Inversuris agrorum.

*Poa panicula diffusa, spiculis trifloris glabris, culmo erecto setati. pratensis. LINN. Spec. 5. Gram. pratense paniculatum maius C. B. Pin. 2. In pratis. *Spiculis in tria rursus subdivis.**

Poa panicula diffusa angulis rectis, spiculis obtusis, culmo obliquo compresso. annua. LINN. Sp. 6. Gram. pratense paniculatum minus. C. B. Pin. 2. Ad vias.

Poa panicula coarctata secunda, culmo obliquo compresso. compressa. LINN. Sp. 15. Gram. minusculum radice repente. C. B. Prodr. 2.

Poa panicula attenuata, spiculis subbifloris mucronatis scabris, culmo incurvo. nemoralis. LINN. Sp. 16. α. Gr. panicul. angustifolium alp. locustis varioribus & angustioribus non aristatis. SCH. 164. β. Gr. cirrhosa & villis spongiosa ad singula genitalia donatum. ROCCON. Mus. 2. p. 70. T. 59. Ad radices umbras montium.

Poa infusa L. 63.

BRIZA. LINN. 78.

Locusta cordata disticha. Calycis valvulae cordatae. Paniculata.

Briza spiculis ovatis, flosculis septem. media. LINN. Sp. 2. Gram. tremulum maius. C. B. Pin. 2. In siccioribus & pratis.

*H. Briza spiculis cordatis, flosculis septemdecim. maxima. LINN. Sp. 3. Gram. tremulum maximum. C. B. Pin. 2. In Italia. Monspelii. & sefor. *in aqua. 0. 1. 2. 3. 4. 5. 6. 7. 8. 9. 10. Tab. 60.**

FESTV.

FESTUCA. LINN. 82.

Locusta oblonga, apice acuminato. Folliculus bivalvis aristatus. Culmus alternis incisuris exsculptus, quem media plana facies locustae respicit.

Festuca panicula secunda coarctata aristata, culmo tetragono nudiusculo, foliis sataceis. ovina, LINN. *Sp.* 1. Gr. *foliis iunceis brevibus, maius, radice nigra*. C. B. *Prodr.* 34. In locis ficcis.

Festuca panicula nutante inferne ramosa, spicis adscendentibus hispidis, foliis setaceis. duriuscula. LINN. *Spec.* 2. Gr. *pratense, panicula duriore laxa, unam partem spectante*. RAII *hist.* 1286. In ficcis.

*Itica elatior.
sub gran. T. 2.
na & Secum Vang.*

Festuca fluitans . oblonga fere . . . *foliis sataceis. Tab. 3.*
TRITICVM. LINN. 94.

Locusta saepius triflora, obtusa. Folliculi valvula exterior obtusa cum acumine. Culmus vt in prioribus.

Triticum radice annua spica glabra aristata. aestivum, LINN. *Spec.* 1. Triticum aestivum. C. B. *Pin.* 21. *Sommer-Weitze*. Rarius seritur.

Triticum radice annua, spica mutica, LINN. *Cliff.* 24. Triticum hybernum, LINN. *Sp.* 2. *Winter-Weitze*. A nostris seritur.

Triticum flosculis aristatis hermaphroditis, intermedio neutro. Spelta, LINN. *Sp.* 4. *Zea dicoccos* vel *Spelta maior*. B. *Pin.* 22. *Spelt*.

Triticum radice repente, foliis viridibus, LINN. *mat. med.* 37. Triticum repens, LINN. *Sp.* 6. *Gramen caninum arvense* f. *gramen Dioscoridis*. B. *Pin.* 1. HALL. *Helv.* 2. *Quecken*. In hortis & arvis.

Var. p. Hall. Hist. n. 1426. aristata. In 1720.
Triti-

Triticum. Gramen latifolium spica triticea compacta. C. B. Prodr. 53. HALL. H. 207. n. 1.
Gr. de laire variet. helv. Ital. Helv. 17. n. 1487.
 β. Gramen latifolium, spica triticea divulla. C. B. Prodr. 54. HALL. l. c. β. ad rivus.

Triticum. caninum. LINN. Sp. 7. Gramen foliaceum, radice repente, f. *Gram. offic. longioribus aristis donatum*. HALL. Helv. 208. Gramen spica triticea compacta birsutum aristatum. HALL. n. 6. Num quæ hæc plantæ modo dictæ varietates sunt *graminis canini vulgaris officinarum*?

Triticum culmo indiviso, spiculis alternis subsessilibus. *Bromus pinnatus*. LINN. Sp. 10. Gramen foliaceum corniculatum. TOURN. Gram. spica Brizæ maius. C. B. Prodr. 53. In sylvis.

β. Gramen foliaceum corniculatum spicis glabris. I. B. H. Festuca dumetorum. C. B. Prodr. 69. In pratis.

LOLIVM. LINN. 90.

Locustæ oblongæ acuminatæ distichæ, dorso culmi alternis incisuris obversæ & appressæ. Calyx uniglumis, folliculus bivalvis. Fl. spicati.

Lolium spica mutica, *perenne*. LINN. Sp. 1. Gramen foliaceum angustiori folio & spica. C. B. Pin. 9. Ad vias & versuras agrorum.

Lolium spica aristata. *remulentum*. LINN. Spec. 2. Gram. foliaceum spica longiore f. *Lolium Diascoridis*. C. B. Pin. 9. In segetibus vdis.

Lolium spicis mediis latioribus. ROY. p. 69. Gr.
loliaceum spicis brevioribus & compressis. MORISON.
 III. S. VIII. T. 2. f. 2. Ad vias.

Elymus canadensis.

BROMVS. LINN. 83.

Locusta ovata. Calyx biglumis, muticus. Folliculus aristam infra apicem emittit. Spiculae paniculatae.

Bromus panicula patente, spicis ovatis, aristis rectis. *secalinus*. LINN. Spec. 1. Gr. *avenaceum* locustis villosis crassioribus. I. R. H. Festuca graminea, glumis birsutis. C. B. Pin. 9. Inter segetes.

Bromus panicula patula, spiculis oblongis distichis, glumis subulato-aristatis. *sterilis*. LINN. Sp. 5. *Bromus* herba. DODON. Cereal. p. 141. Ad vias.

Bromus panicula nutante, spiculis ovato-oblongis. *arvensis*. LINN. Spec. 6. Festuca graminea tuba effusa. C. B. Pin. 192.

Bromus panicula nutante, spicis linearibus. *rectorum*. LINN. Spec. 7. Gr. *avenaceum*, locustis villosis & candicantibus. I. R. H. 526. In agris.

Bromus mollis.

Bromus. HALL. Helv. n. 7. p. 229. Gram. Typhoides molle. C. B. Tk. 50. Gr. spicat. spica cylindracea molli & densa. I. R. H. Folia lanuginosa. Ad ripas Leinae.

Bromus mollis.

Bromus mollis.

Bromus mollis.

Bromus mollis.

Bromus mollis.

Bromus mollis.

Bromus mollis.

Bromus mollis.

Bromus mollis.

Bromus mollis.

Bromus mollis.

Bromus mollis.

Bromus mollis.

Bromus mollis.

Bromus mollis.

Bromus mollis.

Bromus mollis.

Bromus mollis.

Bromus mollis.

Bromus mollis.

AVENA. LINN. 85.

Locusta laxa. Calyx muticus. Ex dorso folliculi arista geniculo reflexa, spiralis. Fl. paniculati.

genuis Loph. Hal. 26. 1. 116. *aristata* Br. 1. 57.

aristata Loph. Hal. 26. 1. 116. *aristata* Br. 1. 57.

aristata Loph. Hal. 26. 1. 116. *aristata* Br. 1. 57.

aristata Loph. Hal. 26. 1. 116. *aristata* Br. 1. 57.

aristata Loph. Hal. 26. 1. 116. *aristata* Br. 1. 57.

aristata Loph. Hal. 26. 1. 116. *aristata* Br. 1. 57.

aristata Loph. Hal. 26. 1. 116. *aristata* Br. 1. 57.

ARUNDO: LINN. 87.

Locusta vni-vel multiflora, mutica. Folliculus intus lanuginosus, cum calyce bivalvis. Fl. paniculati.

H. Arundo calycibus trifloris, panicula diffusa. D. max. LINN. Sp. 2. Arundo sativa. BAVH. Pin. 1. Arundo Indica Laconica versicolor. Cliss. 26. Arundo Americana striata. MONT. F. 108. Variet. I Hispania. Gallo-Provincia.

Arundo calycibus quinquefloris, panicula laxa phragmites. LINN. Sp. 3. Arundo vulgaris f. phragmites. Discoloridis. C. B. Pin. 17. Schilf-Rohr. I lacubus & fossis.

STIPA. LINN. 84.

Locusta uniflora, laxa, biglumis. Folliculus atq. longa ex apice prodeunte, basi articulata. I paniculati.

H. Stipa aristis lanatis. pennata. LINN. Sp. 1. C. spartum pennatum. C. B. Pin. 5. In Marchia. Helvetia. Gallia.

H. Stipa aristis nudis. juncæa. LINN. Sp. 2. F. Juncæa juncæa folio. C. BAVH. Pin. 9.

CYNOSVRVS. LINN. 81.

Involucrum laterale trifolium, saepe pinnatum. Calyx biglumis. Folliculus aristatus.

Cynosurus bracteis pinnatifidis. cristatus. LINN. Spec. 1. Gramen pratense cristatum. B. Prodr. In pratis.

Cynosurus bracteis integris, coeruleus. LINN. *Sp.* 5. *Gr. glumis variis*. C. B. *Prodr.* 21. Ad rupes praeruptas M. *Pless.*

H. *Cynosurus paniculae* spiculis sterilibus pendulis ternatis, floribus aristatis. *aureus*. LINN. *Sp.* 9. *Gramen panicula, pendula, aurea*. C. B. *Pin.* 3. HALL. *Gott.* 49. vbi vide characterem a reliquis speciebus diversissimum.

HORDEVM. LINN. 93.

A. Praeter involucrium triflorum, ex sex aristis compositum; calyx nullus. Folliculus aristatus.

H. *Hordeum flosculis omnibus hermaphroditis aristatis, ordinibus duobus erectioribus, vulgare*. LINN. *Sp.* 1. *Hordeum polystychon vernum*. C. B. *Pin.* 22. *Gerste*. Ad cerevisiam coquendam inprimis feritur.

Hordeum flosculis lateralibus masculis aristatis, marinum. LINN. *Sp.* 5. *Gram. bordeac. minus & vulgare*. C. B. *Pin.* 8. β . *Gr. spicatum secalinum minus*. SCH. 17. Vbiq. ad sepes.

Hordeum flosculis lateralibus imperfectis, mere aristatis, Gramen bordeacum montanum spica striatoflore, brevius aristata. HALL. *Helv.* 204. *Gott.* 12. *Gr. secalinum maius*. In sylvis über Herperhausen.

SECALE. LINN. 92.

Involucrum biflorum aristae duae. Folliculus bivalvis, aristatus.

H. *Secale glumarum ciliis scabris, cereale, L. f. n.*
Sp. 1. α. hybernium vel maius. β. vernum vel mi-
nus, B. Pin. 23, Korn, Rothen, Vbique colitur,

GRAMINIBVS ADFINES.

Culmo, foliis & toto reliquo habitu, pleraeq;
 etiam staminibus, cum prioribus conveniunt, sed
 in flore diuersitas est. His enim stylus vt plurimum
 vnicus & semen est vel vnicum, in bivalvi quidem
 folliculo, sed flores sexu sunt distincti; aut semen
 vnicum, ad vnicam glumam; aut semen vnum,
 plurave capsulis recta, cum flore apetalo sexfolio &
 staminibus tribus vel sex.

I.

Semen stylo bicorni in folliculo biglumi, flores
 sexu distincti.

AEGILOPS. HALL, Gott. 52.

In eadem locusta tri- aut quadriflora, flores &
hermaphroditi laterales, & *masculi* intermedii,
 Calyx biglumis, aristatus. Folliculi bivalvis gluma
 exterior triplici arista terminata, interior mutica,
 quae semen hirsutum obvolvīt. Petala duo plumo-
 sa. Styli duo. Vide quomodo differat character Lin-
 naeanus *Gen. 1018.*

H. *Aegilops spica ovata, aristis brevior, ovata,*
LINN. Sp. 1. Aegilops DOD. Pempt. 73. Cereal.
p. 143. Festuca altera capitulis duris, BAVH. Pin.
10. In Europa australi, B.

CENCHRVS. LINN. 1017. *Panicastrella*

MICH. 31.

Capitula ex locustis bifloris, ternis calyce communi echinato exceptis, in spicam disposita. Flos alter *androgynus*, alter *masculus*. Calyx proprius & folliculus bivalvis, acuminatus.

h. *Cenchrus* spica oblonga conglomerata. *echinata*. LINN. *Sp.* 3. *Gram. locustis tumidiaribus echinatis* SCHEVGHZ. 77. In *Jamaica*. A.

ZEa, LINN. 926. Mays. TOVRN,

Monoph. Vtrique sexui calyx & folliculus bivalvis muticus. *Mares* in locustis bifloris laxè spicatis, *Feminae* foliis rectae; singulis stylus filiformis longissimus. Semina in receptaculo cylindrico ad dimidium demersa.

h. *Zea Mays*. LINN. *Sp.* 1. ROY. 73. Frumentum indicum Mays dictum. BAVH. *Pin.* 25. a. *gracilis aureis*, β. *granis rubentibus*. In *America*. A.

COIX. LINN. 927.

Monoph. *Mares*. Locustae biflorae laxè spicatae, singulae calyce & folliculo mutico. *Feminae* debasin priorum spicatae; fabrica *masculis* simili.

h. *Coix* seminibus ovatis. LINN. *Sp.* 1. ROYEN. 72. *acryma Jobi*. CLVS. *Hist.* 2. p. 216. ZANONI p. 36. T. 101. In *Indiis*. A.

h. *Coix* feminibus angulatis. *dactyloides*. LINN. p. 2. *Gram. dactylon maximum Americ.* PLUCK.

T. 190. f. 2. *Sesamo Indico perpetuo*. ZANON. T. 66. In *America*. P.

CYPEROIDEAE.

II. Stamina tria, & semen vnicum, stylo tricorni ad glumam vnicam.

CAREX. LINN. 928. *Cyperoides*. *Carex*
HALLERi.

Monophyta vt plurimum. Vtriusque sexus flores spicati. Feminis semen triquetrum in nectario apice bidentato.

1. Flores masculi & feminini in eadem spica.

Carex spica paniculata. HALL. *Helv.* n. 12. p. 246. *Carex ra. emq. composito*, LINN. *Nov. Sp. Cent.* I. n. 93. Gram. *cyperoides palustre*, *spica longiori laxa*. RAIL. *Cyperus longus inodorus sylvaticus*. C. BAVH. *Prodr.* n. 24. Ad oras stagnorum v. g. bey der Rasemühle. Seeburg. Landolfshausen.

Carex spicis alternis, latis, laxius in spicam dispositis. *Helv.* 244. Circa Oderbrücke in sylva *Hercynica*. *Adelungia* L. *Hall. Hist.* 1359.

Carex spicis teretibus acutis, alternis, se contingentibus. HALL. *Helv.* 244. n. 5. *Carex brizoides*, LINN. *Nov. Sp. Cent.* I. n. 91. Gram. *cyperoides elegans, spica composita asperiore*. Gott. p. 54. Circa Gottingam & in sylva *Hercynica*.

Carex spicis obefis subrotundis, in summo catule se alterne contingentibus. HALL. n. 7. p. cit. *Carex leporina*, LINN. *Sp.* 6. Gram. *cyperoides palustre*

GRAMINA.

Iustre maius spica divisa. B. Pin. 6. Prope Gettn.
gam versus Lenglern, in locis vdis.

Carex spica composita ex plurimis spicis teretibus, acuris, squamae latae infidentibus. HALL. *Helv. p.* 245. n. 8. *Gottingae* in pratis vdis. v. g.

auf dem Misch. Sympneus zu sein in einem oder in beiden Th. b.
A. 1862. Sympt. aus der Natur

Carex spicis brevibus sessilibus, echinatis in summo caule cohaerentibus. HALL. l. c. n. 9. *Carex vulpina*, LINN. Sp. 7. Gr. *cyperoid. palustre maius, spica compacta.* BAVH. Pin. 6. Ad sepes vdas,

Carex spicis brevibus, echinatis, sessilibus, distinctis. HALL. 245. n. 10. *Carex muricata.* LINN. Sp. 8. Gramen nemorosum, spicis parvis asperis. B. Pin. 7. In nemoribus humentibus. v. g. im Eulentoch.

Carex spicis parvis sessilibus in foliorum alis.
HALL. 243. n. 1. *C. remota*. LINN. *Sp. nov. Cent.*
I. n. 92. MORIS. III. p. 243. T. 12. f. 17. In
sylva Hercyniae.

2. Flores masculini & feminini in spicis distinctis. Cyperoides HALLERI.

Carex (Cyperoides,) spicis ex luteo & spadiceo variis, feminali suprema super spicam matrem elata. MALL. *Helv.* 236. n. 10. *Carex digitata*, LINN. *Sp.* 14. Gram. *caryophyllum*, mont. *spica varia*. C. B. *Prodr.* 9. In nemoribus.

β. Gr. caryophyllatum, nemorosum, spica mul-
tiplici. C. B. Prodr. n. 22. Num varietas? ^{Caryophyllatum Linn.} Herb. hisp. n. 1375.

Carex (Cyper.) capsulis subrotundis spicae vndique rigentibus. HALL. *Helv.* n. 12, Car. *flava*. LINN.

LINN. *Sp.* 12. Gr. *Cyper. aculeatum Germanicum*
f. *minus*. BAVH. *Pin.* 7. In paludibus.

Carex (Cyper.) spicis feminalibus breviter pedunculatis, & inter se & a mare remotissimis. HALL. *Helv.* 237. n. 13. *Cyperoides spicis parvis longe distantibus* 1. R. H. ^{Carex spicis parvis longe distantibus} In vdis.

Carex (Cyper.) spicis feminalibus (longius) pedunculatis, nutantibus, suprema marem erectam & gracilem contingente. HALL. *l. c.* n. 15. *Car. pallens.* LINN. *Sp.* 23. In paludibus.

Carex (Cyper.) spicis feminalibus breviter pedunculatis, rufis, marem obesam contingentibus. HALL. *l. c.* n. 14. *Cyperoides minus spicis densioribus.* 1. R. H. *Bey Robringen & ad Lützelengen.* ^{et al. Hist. 1381. v. n. 1}

Carex spicis masculis pluribus, femineis pedunculatis, capsulis inflatis acuminatis. *vesicaria*, LINN. *Sp.* 29. α. *Cyperoides vesicarium, spicis viridantibus aut subfuscis.* SCHEVCHZ. 470. β. *Cyperoides maius praecox, spicis turgidis, teretibus, flavescens.* MORIS. *Hist.* III. p. 242. γ. *Gram. Cyperoides sylvarum tenuius spicatum.* SCHEVCHZ. 418. In vdis sylvaticis.

Carex (Cyper.) spicis feminalibus longis & asperis, ex longis pedunculis pendulis. HALL. *Helv.* 239. n. 25. *Carex. Pseudo-Cyperus.* LINN. *Spec.* 26. Gr. *cyperoides spica pendula brevior.* C. B. *Pin.* 6. In vdis.

Carex (Cyperoid.) latifolium, spicis feminalibus semimasculis, ex longis pedunculis pendulis. HALL. *Helv.*

Helv. 239. n. 26. Gr. *Cyperoides spica pendula*
longiore & angustiore. c. B. in vdis. ~~an~~ *pendula*, in

Carex (Cyperoid.) spicis maribus obesis, femi-
ninis imis longe petiolatis, gracilibus. HALL. Helv.
31. Gramen *nigro luteum vernum*. I. B. II. p. 494.
Carex. LINN. *Spec.* 28. *a.* In locis gramineis.

Carex (Cyper.) foliis latis & asperis, spicis plu-
rimis, fere sessilibus, duabus androgynis. HALL.
Helv. 29. Gr. *Cyperoid. spica rufa*, f. *caule trian-*
gulo. BAVH. Pin. 6. Carex. LINN. *Spec.* 28. *β.*
In vdis.

Carex. (Cyperoides) HALL. Helv. n. 32. Gram.
Cyperoid. caryophyllaeum, elatius, granis tumi-
dioribus, spicis pedunculis tenuissimis appensis.
RORIS. III. p. 243. n. 13. Im Wald beyrn Hessen-
weisch. Junge. Hall. *an* *Cyperoides*.

Carex: (Cyperoides) HALL. n. 33. Cyperoides *an*
alutro, spicis purpureo spadiceis, tenuibus pe-
dunculis infidentibus. SCH. 467. In vdis. *an* *Cyperoides*.

Carex. (Cyper.) HALL. n. 34. Cyperoides spicis
sessilibus erectis e feminibus confertis compositis.
RAII. III. p. 418. In paludibus versus sylvam supra
Teimar. Hall. 1798. v. infra

Carex spicis remotis, femineis subsessilibus, capsu-
is hirtis. *hirta*. LINN. *Spec.* 13. Cyperoides *poly-*
tachyon lanuginosum. RAI. I. R. H. In dumetis.

3. Flores masc. & fem. in plantis distinctis. *an* *Cyperoides*.

Carex spica simplici dioica. LINN. Sp. I. Vlva.
HALL. Helv. 242. In vdis circa M. Bructerum.

an *Cyperoides*

Carex canaliculata Hall. *an* *Cyperoides* *CYPE*
— *an* *Cyperoides* *an* *Cyperoides* *an* *Cyperoides*
— *an* *Cyperoides* *an* *Cyperoides* *an* *Cyperoides*
— *an* *Cyperoides* *an* *Cyperoides* *an* *Cyperoides*

12.

•

•

•

[illegible]

LNAGROSTIS. HALL. p. 250. Eriophorum.

LINN. 63.

Priori simillima, a quo tantum differt villo seminis spica ipsa longiore.

Linagrostis foliis teretibus, spica subrotunda, vnic. HALL. *Helv.* 250. n. 2. Eriophorum vaginatum. LINN. *Spec.* 1. Gram. *tomentosum alpinum* *minus.* B. *Pin.* 5. In sylva *Hercyn.* & in vdis *Cellensibus.*

Linagrostis foliis planis, spicis multiplicibus. HALL. *l. r.* n. 1. Eriophorum pectinatum. LINN. *Spec.* 2. Gr. *prat. tomentosum, panicula diffusum.* B. *Pin.* 4. In pratis vliginosis.

III. Stamina tria (duo), semen vnicum, stylo (bicorni) tricorni ad glumas sex.

SCHOENVS. LINN. 60.

Calyx locustae multiliorae biglumis. Panicula foliolo excepta. Semen aritis cinctum.

Schoenus flosculis fasciculatis. LINN. *Cliff.* 22. HALL. *Helv.* 251. *albus.* LINN. *Spec.* 9. *Schoenus locustis albicantibus erectis.* Gott. 57. In *Cellensi Ducatu.*

Schoenus spicis gemellis, obesis. HALL. *Gott. l. c.* n. 2. vbi vide characterem a priori satis diversum.)

(**MARISCVS. HALL. Gott. 57.**)

Habitus prioris. Locustae biflorae. Stamina duo, stylus bicornis. Capsula nucleo foeta, aritis non cincta.

Maris-

Mariscus. HALL. *Helv.* p. 251. *Schoenus* cu-
mo tereti, foliis margine dorsoque aculeatis. LINN.
Spec. 1. *Cyperus longus inodorus Germanicus*.
Pin. 14. Im Denckerhäuser Teich.

IV. Stamina sex, capsula tripartita, stylo tri-
corni ad glumas sex.

JVNCVS. LINN. 396.

Calyx communis biglumis. Capsula triquetra
vnilocularis.

Juncus culmo aphylo, mollior, panicula com-
pactiori. HALL. *Helv.* 252. n. 2. *Juncus conglom-*
ratus. LINN. *Spec.* 2. *Juncus laevis panicula ni-*
sparsa. B. *Pin.* 12. In locis vdis.

Juncus culmo aphylo panicula magis spar-
sa. HALL. *l. c.* n. 1. *J. effusus*. LINN. *Sp.* 3. *Junci*
laevis panicula sparsa maior. B. *Pin.* 12. In vdi

Juncus foliis angulatis, culmo dichotomo, flori-
bus solitariis sessilibus. *bufonius*. LINN. *Spec.* 1.
HALL. n. 8. Gr. *iunceum bufonium*. TABERN. p. 22
In locis inundatis.

Juncus foliis mollioribus, carinatis, panicu-
lae multiplici & ramosa. HALL. *Helv.* n. 10. *Junci*
parvus cum pericarpis rotundis. I. B. Ad *Linan*

Juncus foliis duris, carinatis, floribus cartilag-
inosus. HALL. *l. c.* n. 9. *Juncus squarrosus*. LINN. S.
7. Gram. *iunceum*, foliis & spica iunci. B. *Pin.*
In torfaceis circa Oderbrücke.

Juncus foliis articulatis, ramosae paniculae
ramis semel ramosis. HALL. 255. n. 11. *Junci*
foli

foliis nodoso-articulatis. ^{articulatis.} LINN. Sp. 8. Gr. *iuncum*, folio articulato, aquaticum. β. idem cum *utriculis*. γ. idem *sylvaticum*. BAVH. Prodr. 12. In pratis. v. g. bey Herperbausen.

Juncus foliis planis, pilosis, panicula in spicas congesta. HALL. l. c. 256. n. 14. *Juncus campestris*. LINN. Spec. 14. Gram. *birsutum*, capitulis *Psyllii*. C. B. Pin. 7. In sylvis.

Juncus foliis planis, latis, panicula ramosa, sparsa. HALL. n. 14. ^{pilosis.} Gram. *birsutum latifolium minus*. C. B.

Juncus foliis planis latissimis, panicula sparsa, duplicato-ramosa. HALL. 15. ^{ramosa.} Gram. *Luzulae maximum*. I. B. II. p. 493. In sylvis circa Oderbrücke. ^{Helv. 258. n. 26.}

Juncus foliis planis, panicula umbellata, floribus minimis. HALL. 258. n. 26. Gr. *birsutum angustifolium maius*. C. B. Prodr. 16. In sylvis.

^{graminis minus.} TRIGLOCHIN. LINN. 409. HALL.

Helv. 258.

Juncus similis staminibus, & calyce; filiculae autem distinctae tres aut sex, totidemque tubae plumosae.

Triglochin capsulis trilocularibus sublinearibus. *palustre*. LINN. Sp. 1. *Triglochin fructu tenui*. HALL. Helv. 258. In pratis vdis.

Triglochin capsulis sexlocularibus ovatis. *maritima*. LINN. Spec. 2. Gramen *spicatum marit. altissimum*. LOR. ic. 17. In pratis salis prope Harste.

ACORVS. LINN. 392. HALL. 259.

Clava cylindrica testâ floribus sessilibus. Calyx & stamina vt in priori. Capsula trilocularis, tuba brevissima.

Acorus. *Calamus*. LINN. *Sp. 1. Mat. med.* 18. Acorus verus f. *Calamus aromaticus officinarum*. C. B. *Pin.* 34. In fossis. In Ducatu Cellenfi. Circa Eimbeccam. Noribergam.

IV. Monophytæ, flores in receptaculo communibus, staminibus tribus, tuba simplici, calyce triglobo.

TYPHA. LINN. 924.

In suprema parte clavae cylindrica flores mares, in ima feminae femine papposo.

Typha clava crassissima, cylindrica. HALL. *Hist.* 260. *latifolia*. LINN. *Sp. 1. Typha palustris maior*. BAVH. *Pin.* 20. In fossis suburbanis & stagnis ad Denckerhausen.

SPARGANIUM. LINN. 925.

Flores in capitula collecti, masculini in superiori, feminini stylo bicorni in inferiori parte plantae.

Sparganium foliis erectis triquetris. *erectum*. LINN. *Sp. 1. HALL. H.* 259. n. 1. *α. Sparganium ramosum*, *β. non ramosum*. BAVH. *Pin.* 15. In fossis v. g. circa Bovenenden.

Sparganium foliis planis natantibus. *natans*. LINN. *Sp. 2. HALL, n. 2. 260. Sparganium minimum* C. B. RAIL. *Hist.* 1910. Circa Berolinum.

SAV-

PALMAE.

83

SAVRVRVS.

Flores in iulum digesti, ad singulas squamas stamina sex, & bacca tetrasperma, tubis quatuor. HALL. Gott. 9. bacca monosperma, tubis tribus. LINN. gen. 414.

H. Saururus foliis cordatis, petiolatis, spicis solitariis recurvis. LINN. Sp. 1. *Serpentaria repens*, floribus stamineis spicatis, *bryoniae nigrae folio ampliore pingui*, Virgin. PLUK. T. 117. f. 4. In Virginia. Aro proxime affinis. P.

PETALOIDEAE.

CLASSIS. I.

PALMAE.

Caulis (*Caudex*) simplex. Fructificationes in spadice cum Spatha.

ARVM. LINN. 915. HALL. 261.

In spatha cucullata spadix clavae formae superne nudus; inferne germina, in baccas polyspermas abeuntia; in medio antherae sessiles, interiectae duplici ordini papillarum cirrhosarum.

Arum foliis sagittatis, cucullo recto. HALL. 261. *Arum acaule*, foliis hastatis integerrimis, spadice clavato, *maculatum*. LINN. Sp. 12. *Arum vulg. a. maculatum* β. *non maculatum*, B. Pin. 195. BLACKW. T. 218. In sylvis.

H. *Arum acaule*, foliis peltatis ovatis repandis, basi semibifidis. *Colocasia*. LINN. Sp. 6. *Colocasia*. CLVS, Hist. 2. p. 75. *Arum maximum*

Aegyptiarum. α. caule viridi, β. ex atro purpureo.
B. Pin. 195. In *Hispania, Aegypto, Syria.*

CALLA. LINN. 917. Provenzalia. PET.

In spatha ovato cordata spadix digitiformis, vndique fere tecta antheris sessilibus, & germinibus in baccas abeuntibus.

H. *Calla* foliis sagittato cordatis, spatha cucullata, spadice superne masculino. *Aethiopica.* LINN. *Sp.* 1. *Arum Aethiopicum, flore odorato Moschum redolente.* COMMEL. *Hort.* 1. p. 95. T. 50. Antherae & germina sine certo ordine digesta mihi visa.

Calla palustris

in Quercu ciliata, Afr. Ind.

PALMA. PHOENIX. LINN. 1089,

LVDW. 908.

In spatha alari spadix ramosus. Corolla tripartita. Diphyta. *Mas.* Stamina sex. *Fem.* Bacca unilocularis, stylo simplici. Caudex apice frondosus.

H. *Phoenix* frondibus pinnatis; foliolis complicatis. *dactylifera.* LINN. *Sp.* 1. p. 1188. *Palma hortenſis, f. dactylifera.* BLACKW. T. 202. & auctor. In *India, P.*

MVSA. LINN. 1010. LVDW. 370.

Caudex apice frondosus. In spatha alari flores multi, corolla dipetala, staminibus sex, fructu carnoso longo, stylo cylindraceo. In flor. superioribus ovaria abortiunt.

H. *Musa* racemo simplicissimo. LINN. *H. Cliff.* 467. *Mus. Cliff.* 1. T. 1. spadice nutante, *paradisaca,*

faci. Sp. 1. TREW. Dec. Pl. rar. 11. & Commenc.
lit. 1739, 40, 41. In India.

CLASSIS. II.

ORCHIDEAE.

Calculus simplex, foliis simplicibus vaginantibus
 tinctus. Capsulae trivalves striatae flores sustinent.
 Petala quinque aut quatuor superiora in cucullum
 conniventia, sextum inferius cavum, nectariferum,
 interdum calcaratum *labelli* aut *nectarii* nomine,
 quod gerit machinam duorum staminum, sub
 qua stigma concavum. Vid. HALLER. *Comment.*
Soc. Sc. Gott. Tom. V.

CALCEOLVS. HALLER. l. c. Cypripedium.

LINN. 906.

Filamenta staminum duo distincta. Petala supe-
 riora quatuor, inferius amplissimum, cavum. Radix
 ramosa.

Calceolus. HALL. *Helv.* 276. *Gott.* 65. Cypripe-
dium *radicibus fibrosis, foliis ovato lanceolatis cau-*
linis. LINN. *Sp. 1. Calceolus Marianus.* DOD. *Mari-*
enshub. Circa *Detersen.*

EPIPACTIS. HALLER. Serapias. Ophrys.

LINN.

Machina staminum est scapus superne bilabiatus,
 abium superius inferne duas antheras nudas; labio
 inferiori incumbentes, fovet. Calcar nullum. Cap-
 su ovata. Radix ut plurimum ramosa.

1. *Helleborine*. labellum cavum, ex ovato-cordatum.

ep. longifolia. Epipactis (*Helleborine*) floribus obsolete carneis, raris, labello obtuso. HALL. Enum. Helv. 275. *Helleborine palustris nostras*. RAIL. hist. 1231. *angustifolia pratensis*. B. Pin. 87. *Serapias*. LINN. Sp. 1. y. Ad lacum prope Seeburg. Im Landolfsbäusen Sumpf. *Trope sylvan. prope Seeburg.*

latifolia. Epipactis (*Helleborine*) amplexicaulis multiflora. Helv. 275. *Helleborine latifolia montana*. C. B. a. flore atro rubente. B. flore pallido. C. B. Pin. 186. *Serapias*. LINN. Spec. 1. a. In sylvis v. g. circa Pless. Göttinger Wald.

angustifolia. Epipactis (*Helleborine*) latifolia floribus magnis paucis albisque. En. Helv. 276. *Helleborine f. Damalonium montanum latifolium*. C. B. In sylvis v. g. Göttinger Wald, circa Pless. &c.

longifolia. B. *Helleborine longifolia*, floribus magnis, paucis & albis. Helv. 275. Auf dem Bocksbühl. Epipactis (*Helleborine*) *longifolia*, paucis pureis floribus. Helv. 274. *Helleborine mont. angustifolia purpurascens*. B. Pin. 187. In sylvis v. g. circa Pless.

Epipactis (*Helleborine*) bifolia radice ovali, cucullo sparso. Comm. Soc. Sc. Gott. Tom. I. T. XIV. p. 215. Circa Dersfen.

2. *Ophrys*. Labellum longum, bicornes.

Epipactis (*Ophrys*) foliis ovatis. LINN. Cliff. 429. ROYEN, 14. HALL. Helv. 277. *Ophrys ovata*. LINN.

LINN *Sp. 5.* *Ophrys bifolia*. B. *Pin.* 87. In pratis & sylvis.

Epipactis (*Ophrys*) *aphylla*. HALL. *Helv.* 278.

Ophrys. *Nidus avis*. LINN. *Sp. 1.* *Orchis abortiva fusca*. B. *Pin.* 86. *Nidus avis*. LÖB. *ic.* 195. 1. R.

H. RAL. Omnes plantae partes lignei coloris. In locis vmbrosis nemorum.

ORCHIS. HALLER. *Orchis*. *Satyrion*.

LINN. 900. 901.

Machina staminea. In duobus loculis styli brevioris membrana clausis antherae duae ex filamentis convolutis factae. Calcar modo adest, modo deest.

Orchis radicibus subrotundis, cucullo purpureo, maculato, labello albo. HALL. *Gott. p.* 61. *Helv. n. 3.* *Orchis* LINN. *Sp. 10. 6.* *Orch. militaris minor*. B. *Pin.* 81. In M. *Hengst.*

Orchis radicibus subrotundis, brachiolis labelli brevioribus, spinula ex divisione medii trunculi eminente. HALL. *Helv.* 262. *Gott. l. c.* *Orchis* LINN. *Sp. 10. α.* *Orchis galea & alis fere cinereis*. 1. B. II. *p.* 757. In *Suecia*, rarius supra *Bovenden*.

Orchis radicibus subrotundis, spica brevissima labello punctato, serrato, mucico. HALL. *Gott. p.* 61. *Orchis* LINN. 10. γ. *Orchis militaris minor*. 1. B. RVPP. III. T. VI. In gramineis ficcis.

H. *Orchis* radicibus subrotundis, bracteis ovario longioribus, albo flore. *Gott.* 62. *Orchis foetida, sylvatica, alba, praecox, barba luteola*. RVPP. III. *p.* 297. Circa *Jenam*. In horto adest.

Orchis radicibus subrotundis, floribus obsoletis, cimicem olentibus, labello retro reflexo. *Enum. Helv.* 264. Orchis ^{*conopsea*} LINN. *Sp.* 6. Orchis *hirci* odore minor. B. *Pin.* 82. In pratis circa Dorsten ad viam versus Osterode, circa Hardeggen, ad lacum See-burgensem. *ad omni agris sub sylvis Raps. & Chaulis.*

Orchis radicibus subrotundis, cucullo sparso, labello ampliter trifido, crenato, emarginato. H. *Helv.* 265. Orch. ^{*Morio*} LINN. *Sp.* 8. *δ.* Orchis *morio mas.* B. *Pin.* 81. *Satyrium mas.* BLACKW. *T.* 53. In sylvis.

Satyrium mas. - bulb. & dimidi. nectaris latius quod dicitur, in sylvis, circa Dorsten, & in pratis, circa Osterode, & in sylvis, circa Seeburg.
Orchis radicibus subrotundis, cucullo clauso, lineato, labello ampliter trifido, crenato. HALL. *Helv.* 266. Orchis LINN. *Sp.* 8. *α.* Orchis *morio femina.* B. *Pin.* 82. In pratis siccis & caeduis sylvarum.

Orchis conopsea - fructus & semina, circa Dorsten.
Orchis radicibus conicis, labello lingulato, longissimo. HALL. *Helv.* 266. Orchis *bifolia.* LINN. *Sp.* 2. Orchis *alba bifolia minor, calcare oblongo.* BAVH. *Pin.* 83. SEGV. *Veron.* *T.* 15, f. 10. In sylvis.

Orchis radicibus subrotundis, calcare nullo, labello latiusculo, bisulco, brachiolis brevibus. HALL. *Helv.* 268. *Ophrys insectifera.* LINN. *Sp.* 15, *α.* *Orchis muscam referens.* B. *Pin.* 83. Rarius in Göttinger Wald bey der Brück, circa Pless, &c.

Orchis radice subrotunda, calcare nullo, labello trifido. HALL. *Gott.* 63. *Helv.* 269. *n.* 20. *Ophrys.* LINN. *Sp.* 11. *Herminium Fl. Lapp.* 317. *Monorchis montana minima, flore obsoleto, vix conspicua* MICHEL.

CHEL. p. 30. T. 26. Monorchis. c. B. In pratis
 & Dorsten & Osterode. Ad viam inter Hesse-
 fch & Roßbringen. *Epipactis atrorubens* L. *Epipactis atrorubens* L. *Epipactis atrorubens* L.

Orchis radicibus palmatis, calcare ovario lon-
 ge, labello aequaliter trifido, unicolor. HALL.
 v. 271. *Orchis conopsea*. LINN. 13. *Orchis*
nata minor, calcaribus oblongis. c. B. Pin. 85.
 pratis sylvarum.

Orchis radicibus palmatis, calcare crasso, foliis
 sulcatis, caule fistuloso. HALL. Gott. 63. Helv. n.

Orchis latifolia. LINN. Sp. 11. *Orchis palma-*
tratenfis α. *maculata*. β. *non maculata*. γ. *lati-*
a. &c. c. B. In pratis vbique.

Orchis radicibus palmatis, calcare crasso, labello
 funde trifido, ferrato, caule solido. HALL. Gott.

Orchis maculata. LINN. Sp. 12. *Orchis palma-*
montana, maculata. c. B. Pin. 86. In sylvis
 natis.

Satyrium

Orchis radicibus multis cylindricis, calcare bre-
 vi. HALL. Helv. 270. *Satyrium albidum*.
 LINN. Sp. 4. *Pseudorchis alpina, flore herbacea*
 CHEL. gen. 32. T. 26. In summo M. Brüttera.

Satyrium *Epipactis*

CLASSIS. III.

LILIACEAE.

Calyx nullus, cuius loco in variis spatia membra-
 ea, marcescens. Corolla tripetala, aut bis tri-
 ala, aut monopetala limbo bis trifido. Stamina
 aut sex. Stylus simplex, stigmate trifido, rarius

stylis tres. Fructus trigonus trilocularis, trivalvis. Folia simplicia, integra, nervis parallelis. Caulis aut nullus, aut foliis vaginantibus cinctus, simplex, herbaceus. Radix bulbosa, vel tuberosa, perennis.

I. Fructu sub flore posito,

α. flore tristemente. His fere omnibus folia ensata, & spathae sub flore.

COMMELINA. LINN. 58.

Corolla bis tripetala, tria petala calycem fermentientia. Filamentis tribus corpuscula cruciformia insistent.

Commelina corollis inaequalibus, foliis ovato-lanceolatis acutis, caule procumbente glabro. *communis*, LINN. *Sp.* 1. *Commelina procumbens americana*, *saponariae folia*, DILL. *Elth.* T. 78. f. 89. In America. A.

Commelina corollis inaequalibus, foliis lanceolatis glabris obtusis, caule repente. *africana*, LINN. *Sp.* 2. *Commelina procumbens flore luteo*, ROYEN. 538. HALL. *Gott.* 66. In Aethiopia.

Commelina corollis aequalibus, foliis ovato lanceolatis subciliatis. *tuberosa*, LINN. *Sp.* 5. *Commelina radice Anacampteros*, DILL. *Elth.* T. 79. f. 90. In Mexico.

TRADESCANTIA. LINN.

Spatha foliacea flores continens. Calyx trifolius. Petala tria. Stamina villosa.

H. *Tradescantia*. *Virginiana*, LINN. *Sp.* 1. *Ephemerum*, TOURN. 123. In Virginia.

BRO-

LILIACEAE.

91

BROMELIA. LINN. 356. Ananas.

TOURN.

Flores in nostra specie in conum foliis coronarum collecti, spathis intermixtis. Calyx trigonus. Petala tria. Bacca umbilicata.

H. *Bromelia* foliis ciliato spinosis mucronatis, spica comosa. *Ananas*. LINN. *Sp.* 1. *Ananas aculeatus*, fructu pyramidato, carne aurea. I. R. H. 653. In *America* meridionali.

HAEMANTHVS. LINN. 394.

Involucrum sexfolium umbelliferum. Corolla profunde sexpartita. Bacca subrotunda.

H. *Haemanthus* foliis linguiformibus planis. *occidentalis*. LINN. *Sp.* 1. *Haemanthus Africanus*. COMM. H. II. p. 127. T. 64. Ad C. bon. Spei.

H. *Haemanthus* foliis lanceolatis, *punicus*. LINN. *Sp.* 2. *Haemanthus colchici foliis*, perianthio herbaceo. DILL. *Elth.* 167. f. 140. In *Africa*.

GLADIOLVS. LINN. 55.

HALL. 279.

Corolla sexpartita, bilabiata. Stamina adscendentia; stigmata undulata.

H. *Gladiolus* foliis ensiformibus, floribus distantibus. *communis*. LINN. *Sp.* 1. HALL. *Helv.* 279. *Gladiolus floribus uno versu dispositis* C. B. Pin. 41. In *Marchia* & *Europa* australi.

IRIS.

IRIS. LINN. 57. HALL. 279. Xiphium.
Sisyrinchium.

Corolla sexpartita, petala tria reflexa. Stigmata
lata, petaloidea, antheras deprimentia.

1. Petalis reflexis barbatis.

H. Iris corollis barbatis, caule foliis longiore uni-
floro. *Susiana*. LINN. *Sp.* 1. Iris *Chalcedonica latifolia*.
I. B. II. p. 721. Iris *Susiana*, flore maximo ex al-
bo nigricante. C. B. *Pin.* 31. In Oriente.

H. Iris corollis barbatis caule foliis longiore,
multifloro. *Germanica*. LINN. *Spec.* 2. Iris nostras
bortenfis. BLACKW. *T.* 69. Iris *vulg.* *Germanica* f.
sylvestris C. BAVH. *Pin.* 30. a. petalis reflexis vio-
laceis (coeruleis), erectis aquate coeruleis (albidis).
In Germaniae editis. ^{in Germania}
^{in Germania}

H. Iris petalis reflexis fulcopicis, erectis luteis.
Iris *lutea variegata*. LOBEL. *hist.* 34. Iris ^{variegata} LINN.
Sp. 4. In Pannonia. Num Varietas?

H. Iris corollis barbatis, caule foliis brevioris uni-
floro. *pumila*. LINN. *Sp.* 6. Iris *humilis latifolia*
minor, flore purpureo. I. R. H. 361. In Austria.
Pannonia, Sibiria.

2. Petalis reflexis imberbibus.

Iris corollis imberbibus, petalis interioribus sti-
gmate minoribus, foliis ensiformibus. *Pseudacorus*.
LINN. *Sp.* 7. Iris *imberbis lutea*, nervo folii emi-
nente. HALL. *Helv.* 281. *Acorus adulterinus*.
BLACKW. *T.* 361. C. BAVH. *Pin.* 34. *Gelbe Wasser-*
Lilien. Ad ripas fossarum.

. Iris corollis imberbibus, petalis internis longine stigmatis, foliis ensiformibus. *foetidissima*. LINN. 8. Iris *foetidissima* f. *Xyris*. TOURN.
 . Spatula *foetida* f. *Xyris*. BLACKW. T. 158. In *lia, Gallia, Hetruria*.

. Iris corollis imberbibus, germinibus trigonis, tereti, foliis linearibus. LINN. Sp. 9. HALL. 1. 280. Iris *angustifolia non foetida altior*, C. H. Pin. 32. In *Marchia*. Nobis hortensis.

. Iris sylvestris peramoena. LEB. 16, 69.

. Iris corollis imberbibus, germinibus sexangulis, caule ancipiti, foliis linearibus. *graminea*. N. Sp. 13. ROYEN. 18. In *Austria*.

. Iris corollis imberbibus, foliis subulato canatis caule brevioribus. *Xiphium*. LINN. Spec. 16. *bulbosa latifolia, caule donata*. C. B. Pin. 38. *hispania*.

SISYRINCHIVM. LINN. 908.

patha diphylla, vniversalis. Antherae stylo ad stigmatis affixae.

. *Sisyrinchium* caule foliisque ancipitibus. N. Sp. 1. *Bermudiana iridis folio, fibrosa radi-* MILL, *Eltb*, T. 41. f. 48. In *Bermudis*.

IXIA. LINN. 54.

etala sex aequalia. Stigmata convoluta. Stamina. Spathae vagae.

. *Ixia* foliis ensiformibus, floribus remotis. Spec. 2. *Bermudiana flore aurantio macula-*

to. HALL. *Cott.* 68. ROLOF. *hort. Krauf.* p. 25.
ic. In India.

CROCVS. LINN. 53.

Corolla sexpartita, aequalis, ex spatha mono-
 phylla. Stigmata convoluta ferrata.

H. *Crocus* spathâ univalvi radicali, corollae tubo
 longissimo. *lacteus*. LINN. *Sp.* I. BLACKW. *T.* 144.
Crocus α. *vernus*. β. *autumnalis*. HALL. *Helv.* 2812
variorum colorum. Sponte in alpinis *Helveticis*,
Pyrenaeis.

β. Flore hexastemone, ut plurimum spathaceâ

COLCHICVM. LINN. 415.

Habitus fere Croci. Tubus floris radicans. Stylis
 tres, stigmata canaliculata.

Colchicum foliis planis lanceolatis erectis. *autu-*
mnale. LINN. *Sp.* I. *Colchicum*. HALL. *Helv.* 283.
 α. purpureum β. album γ. ex albo & purpureo
 varium. Ubique in pratis; germen intra radicem
 fovet, dum autumnis floret, altero vere proruptu-
 rum & in capsulam inflatam abiturum.

GALANTHVS. LINN. 362.

Petala tria maiora patula, tria dimidio minora;
 cylindrum formantia. Stigma simplex.

H. *Galanthus* *nivalis*. LINN. *Sp.* I. ROY. 35.
Leucoium bulbosum trifolium minus. C. B. *Pin.* 56.
 In hortis; spontanea fere in pomario arcis *Cattlen-*
burg. Patria alpina.

LEVCOIVM. LINN. 363.

Corolla campaniformis, petalis apice incrassatis.
Stigma fetaceum, simplex.

Leucoium spatha uniflora, stylo clavato. vernum.
LINN. Spec. 1. Leucolum petalis aequalibus. HALL.
Helv. 284. Narcisso-Leucoium vulgare. 1. R. H.
In sylvis. v. g. über Geismar. im Eulenloch.

NARCISSVS. LINN. 364. HALL. 284.

Ex tubo longo petala sex plana, inserta nectario cylindraceo monophyllo. Stigma breviter trifidum.

h. Narcissus spatha uniflora, nectarii limbo rotato brevissimo, (foliis ensiformibus). poeticus. LINN.
Sp. 1. Narcissus albus circulo purpureo & croceo.
C. B. Pin. 48. 1. R. H. In Helvetia. Gallia.

h. Narcissus spatha uniflora, nectarii limbo campanulato erecto petalo aequali. Pseudo-Narcissus.
LINN. Sp. 2. Narcissus sylvestris pallidus, calyce luteo. C. B. Pin. 52. In Angliae, Italiae nemoribus.

h. Narcissus spatha multiflora, nectario campanulato brevi, foliis subulatis. Fonquilla. LINN. Sp. 5.
Narcissus uncifolius luteus minor C. B. Pin. 51.
Sponte Monspelii.

AMARYLLIS. LINN. 367.

Petala sex lanceolata. Stigma trifidum tenue
Corolla in aliis aequalis, in aliis inaequalis, bilabiata.

H. *Amaryllis* spatha uniflora, corolla aequali, pistillo declinato. *Atamasca*. LINN. Sp. 2. Lilionarcissus ind. flore albo exterius rubente. I. R. H. In Virginia.

H. *Amaryllis* spatha uniflora, corolla inaequali, genitalibus declinatis. *formosissima*. LINN. Sp. 3. Lilio-Narcissus *Jacobaeus*, flore sanguineo nutante. DILL. Elb. T. 162. f. 196. In America meridionali.

H. *Amaryllis* spatha multiflora, corollis revolutis, genitalibus strictis. *Sarniensis*. LINN. Sp. 5. Narcissus *Japonicus*, rutilo flore. I. R. H. Guernsey-Lilly. DOUGLASS in proprio libello. Sponte in Japonia, nunc in insula Guernsey.

ALOE. LINN. 389.

Corolla monopetala in fundo nectarifera ventricosa, aut limbo parvo sexfido, aut profundius sexpartita, modo simplex aequalis, modo difformis. Stamina receptaculo inserta, stigma modo simplex, modo trifidum. Capsula oblonga, trifurca.

H. Aloe floribus pedunculatis cernuis corymbosis subcylindricis *perfoliata*. LINN. Sp. 1. Aloe foliis caulinis dentatis, amplexicaulibus vaginantibus. Cliff. 132. Ad hanc speciem tanquam varietates refert LINNAEVS sequentes, quae sunt.

α. Aloe Afr. caulescens, foliis glaucis brevissimis. COMMEL. Prael. T. 22. rarior. T. 45.

β. Aloe Africana caulescens, foliis glaucis caulem amplexantibus. COMMEL. *H. Amst.* II. T. 14. HALL. *Gott.* 70.

γ. Aloe Afr. foliis glaucis margine & dorso superiore spinosis, flore rubro. COMMEL. *Prael.* 24. *H. Amstel.* II. T. 12.

δ. Aloe Afr. caulescens, foliis glaucis, longioribus, in apice dorso etiam spinosis. HALL. *Gott.* 72.

ε. Aloe Afr. foliis glaucis, margine & dorso integro spinosis. COMMEL. *Prael.* T. 20. *H. Amst.* II. T. 13. *Gott.* 71.

ζ. Aloe caulescens foliis minus glaucis, caulem amplexantibus. COMMEL. *Prael.* 18.

η. Aloe Afr. caulescens, foliis spinosis, maculis ab utraque parte albicantibus notata. COMM. *H. Amst.* II. T. 5. HALL. *Gott.* 72.

θ. Aloe Africana maculata spinosa maior: DILL. *Eth.* T. 15.

ι. Aloe Africana humilis, spinis & verrucis obsita. COMM. *Prael.* T. 16. *Pl. rar.* T. 46. Conveniunt quidem stigmate simpliciter, sed differunt rubo, imbo & colore floris.

κ. Aloe floribus pedunculatis cernuis racemosis prismaticis, ore patulo. *variegata*, LINN. *Sp.* 2. Aloe foliis canaliculatis, trifariam imbricatis radicatis rectis: angulis cartilagineis ternis. *Hort. Cliff.* 132. Aloe Afr. humilis foliis ex albo viridi variegatis. COMM. *Prael.* 79. T. 28. *Pl. rar.* T. 27.

h. Aloe floribus pedunculatis ovato cylindricis curvis. *disticha*. LINN. *Sp.* 3. Aloe *foliis linguiformibus patulis distichis*. Cliff. 132.

α. Aloe Afr. flore rubro, folio triangulari, maculis ab utraque parte albicantibus notato. COMM. *Amst.* II. f. 8. HALL. *Gott.* 75.

β. Aloe Africana sessilis, foliis carinatis verrucosis. DILL. *Elth.* T. 16.

γ. Aloe Afr. foliis planis coniugatis, carinatis verrucosis, caule & flore corallii colore. BOERH. II. p. 131. MART. *Cent.* II. T. 11.

δ. Aloe Afr. arborescens, montana, non spinosa, folio longissimo, plicatili, flore rubro. COMM. *H. Amst.* II. T. 3.

h. Aloe floribus sessilibus horizontalibus infundibuliformibus aequalibus, limbo revolutis. *hyacinthoides*. LINN. *Spec.* 4. Aloe *foliis lanceolatis planis rectis radicatis*. Cliff. 32.

α. Aloe Ceylanica pumila, foliis variegatis. COMM. *Amst.* II. T. 21.

β. Aloe Guineensis, radice geniculata, folio variegato. COMM. *Prael.* T. 33. *H. Amst.* II. f. 20.

h. Aloe floribus sessilibus ovatis crenatis, segmentis interioribus conniventibus. *spiralis*. LINN. *Sp.* 5. Aloe *foliis ovatis acuminatis caulinis quinquefaris imbricatis*. Cliff. 132. Aloe Afr. *erecta*, folio *rotundo*, *acuminato*. COMMEL. *Prael.* 32. HALL. *Gott.* 73.

h. Aloe floribus sessilibus triquetris, bilabiatis labio inferiori revoluti. *retusa*. LINN. *Sp.* 6. Aloe
foliis

foliis rhomboidalibus ventricosis quinquesariam positis radicatis triquetris plano-extantibus. Cliff. 132. Aloe Africana, brevissimo crassissimoque folio, flore viridi. COMM. Amst. II. T. 6.

h. Aloe floribus sessilibus infundibuliformibus bilabiatis, laciniis quinque revolutis: summa erecta, viscosa. LINN. Sp. 7. Aloe foliis canaliculatis triariam imbricatis, caulinis apice reflexo-patulis. Cliff. 132. Aloe Afr. erecta triangularis & triangulari folio viscoso. COMM. Praed. T. 31. DILL. Elsb. T. 13. f. 13.

h. Aloe floribus sessilibus bilabiatis: labio superiore erecto, inferiore patente. *pumila*. LINN. Sp. 8. Aloe foliis ovato subulatis acuminatis tuberculis cartilagineis undique adspersis. Cliff. 131.

α. Aloe Afr. margaritifera minor. COMM. Amst. II. T. 11. DILL. Elsb. T. 16. f. 17.

β. Aloe Afr. margaritifera minima, DILL. Elsb. T. 16. f. 18.

γ. Aloe Afr. humilis arachnoidea. COMM. Praed. T. 27.

δ. Aloe Afr. minima atroviridis, spinis herbaceis numerosis ornata. BOERH. II. T. 131.

Conveniunt petalis sex fere distinctis conglutatis, stigmate trifido.

h. Aloe floribus sessilibus reflexis imbricatis prismaticis. *Vuaria*. LINN. Sp. 9. Aloe foliis lineari-bus radicalibus membranaceis. Cliff. 133. Aloe Afr. folio triangulari longissimo & angustissimo, floribus luteis foetidis. COMM. II. T. 15.

AGAVE. LINN. 390.

Corolla campaniformis. Stamina flore duplo longiora. Stigma trigonum.

H. Agave foliis dentato-spinosis, scapo ramoso. Americana. LINN. *Sp.* 1. Aloe folio in oblongum aculeum abeunte. C. B. *Pin.* 286. Aloe Americana. HALL. *Gott.* 69. TREW. *Comm. litt.* 1744. T. 3. In America.

H. Agave folio margini lobis foliis p. 172

II. Fructu intra florem posito.

α. solitariis nudis floribus.

1. floribus hexapetalis, aut ad basin divisis, campanulatis.

YUCCA. LINN. 388. Cordylina.

ROYEN.

Corolla vnguibus cohaerens. Stylus nullus. Filamenta staminum superne crassiora.

H. Yucca foliis integerrimis. gloriosa. LINN. *Sp.* 1. Yucca filiiis Aloes. BAVH. *Pin.* 91. In Canada, Peru.

H. Yucca foliis crenatis nutantibus. draconis. LINN. *Spec.* 1. Yucca Draconis folio serrato. DILL. *Eltham.* T. 324. f. 417. Cordylina foliis pungentibus crenatis. ROY. *Lugd.* 22. In America.

HEMEROCALLIS. LINN. 391.

Ex tubo brevi corolla fere infundibuliformis. Stamina declinata. Stylus filiformis.

H. Hemerocallis scapo ramoso, corollis monopetalis. Liliosphodelus. LINN. *Sp.* 1. α. Liliosphodelus

delus, luteus. CLVS. *Hist.* 1. p. 137. In *Sibiria Italia.*

Memorialis folia 2. p. 462.

β. *Lilioasphodelus phoeniceus.* CLVS. l. c. LOB. 16. 93. In *India.*

LILIVM. LINN. 371.

Petala reflexa, ad basin singulorum linea nectarifera. Stylus stigmate triangulari.

H. *Lilium* foliis sparsis, corollis campanulatis, intus glabris. *candidum.* LINN. *Sp.* 1. *Lilium album.* C. B. *Piq.* 76. BLACKW. *T.* 11. & *offic.* In *Paestina.*

H. *Lilium* foliis sparsis, corollis campanulatis erectis, intus scabris, *bulbiferum.* LINN. *Sp.* 2. *Lilium purpureo-croceum.* C. B. *Pin.* 77. In *Sibiria Austria. Italia.*

Lilium foliis verticillatis, floribus reflexis corollis revolutis. *Martagon.* LINN. *Sp.* 5. HALL. *Hely.* 289. *Lilium montanum f. sylvestre.* In sylvis.

H. *Lilium* foliis verticillatis, floribus reflexis, corollis campanulatis. *Canadense.* LINN. *Sp.* 6. *Lilium angustifolium, flore flavo maculis nigris distincto.* I. R. H. 371. In *Canada.*

FRITILLARIA. LINN. 372. Corona Imperialis
TOURN. 197.

Fovea nectarifera ad basin petalorum. Stigma triplex. Flores penduli.

H. *Fritillaria* foliis omnibus alternis. *Meleagris.* LINN. *Sp.* *Fritill. v. foliorum alis florens.* Cliff. 119.

Fritillaria α . *alba*. β . *purpureo variegata*. C. B. *Pin.* 64. In *Helvetia*. *Gallia*.

H. *Fritillaria* racemo comoso inferne nudo, foliis integerrimis. *imperialis*. LINN. *Spec.* 1. *Corona imperialis*. I. R. H. C. B. & auctor. In *Oriente*.

ERYTHRONIVM. LINN. 375.

Ad basin trium petalorum interiorum tubercula duo. Stylus stigmatibus triplici.

H. *Erythronium*. *Dens Canis*. LINN. *Sp.* 1. *Dens Canis* BAVH. *Pin.* 87. In *Italia*. *Sibiria*. *Virginia*.

TULIPA. LINN. 376.

Stylus nullus. Stigma triangulare, trilobum, angulis bifidis. Capsula trigono-teres.

H. *Tulipa* flore erecto, foliis ovato-lanceolatis. *Gesneriana*. LINN. *Sp.* 2. *Tulipa* BAVH. & auctorum. In *Oriente*.

H. *Tulipa* flore subnutante, foliis lanceolatis. *sylvestris*. LINN. *Sp.* 1. *Tulipa minor lutea italica gallica*. B. *Pin.* 63. *Monspeli*. In pomario *Cattlenburgensis* praefecturae quasi spontanea.

2. floribus monopetalis.

HYACINTHVS. LINN. 385.

Corolla monopetala, in aliis sexpartita, in aliis limbo breviter sexfido. Ad apicem germinis tres pori nectariferi.

H. *Hyacinthus* corollis campanulatis, sexpartitis apice revolutis. *non scriptus*. LINN. *Sp.* 1. *Hyacinthus*

Hyacinthus oblongo flore coeruleus maior. C. B. Pin. 43.
In *Anglia, Gallia, Helvetia.*

H. *Hyacinthus* corollis infundibuliformibus semis-
fexfidis basi ventricosus. *orientalis.* LINN. Sp. 5.
Hyacinthus orientalis, C. B. Pin. 44. DOD. Pempt.
216. Variæ flore pleno in hortis. In *Asia, Africa.*

H. *Hyacinthus* corollis ovatis. *Muscari.* LINN.
Sp. 6. *Hyac. racemosus muscibatus.* C. B. Pin. 43.
Muscari obsoletiori flore. CLVS. Hist. I. p. 178.
In *Syria.* H. *Hyac. botryoides.* L. p. 92. p. 1

H. *Hyacinthus* corollis subovatis. *monstrosus.*
LINN. Spec. 7. *Hyac. Sannesius panicula comosa.*
COL. Ecpbr. 2. p. 10. T. 12.

H. *Hyacinthus* corollis angulato-cylindricis, sum-
mis sterilibus longius pedunculatis. *comosus.* LINN.
Spec. 8. *Muscari latifolium, summis floribus in lon-
gis petiolis erectis.* HALL. Helv. 285. n. 1. *Hyac.
max. botryoides, coma coerulea.* 1. B. II. p. 574.

POLYANTHES. LINN. 384.

Corolla infundibuliformis. Stamina ex faucibus
corollae orta.

H. *Polyanthes* floribus alternis. *tuberosa.* LINN.
Sp. 1. Cliff. 127. *Hyacinthus indic. tuberosa radice.*
CLVS. Hist. I. p. 176. *Tuberosa hortulanorum.* In
India.

POLYGONATVM. HALL. 286. Convallariæ LINN. spec.

Corollae campanulatae limbus breviter fexfidus.
Bacca globosa immatura maculata.

Polygonatum bifolium, scapo florifero aphyllō.
HALL. *Helv.* 286. *Convallaria maialis*, LINN. *Sp.*
1. *Lilium convallium*, C. B. *Pin.* 304. BLACKW.
T. 70. *Mayblume*. In sylvis.

Polygonatum foliis alternis, pedunculis pendulis
multifloris. HALL. *l. c.* 287, *n.* 4. *Convallaria multi-*
flora, LINN. *Spec.* 4. *Polygonatum maximum lati-*
folium, C. B. *Pin.* 303. BLACKW. *T.* 251. vulgo *Si-*
gillum Salomonis. In dumetis.

Polygonatum foliis alternis, pedunculis pendu-
lis unifloris. *En. Helv.* 287. *Convallaria Polygo-*
natum, LINN. *Sp.* 3. *Polygonatum flore odoro-*
C. B. *l. c.* In praecipitiis circa arcem *Pless*.

Polygonatum foliis verticillatis, HALL. *l. c.* 288.
Convallaria verticillata, LINN. *Sp.* 2. *Polygonatum*
angustifolium non ramosum, C. B. *Pin.* 303. In
sylvis.

(VNIFOLIUM, HALL. *Helv.* 288. *Convallariae*
Sp. LINN.

Priori proxime vicinum. Flos autem quadrifidus.
Stamina quatuor. Fructus bilocularis.

Vnifolium, HALL. *l. c.* 288. *Convallaria foliis*
cordatis, LINN. *Sp.* 3. *Lilium convallium minus*,
B. *Pin.* 304. In sylvis.

ASPARAGVS. LINN. 382.

Petala vnguibus cohaerentia. Stamina petalis in-
serta. Bacca globosa umbilicata.

h. Aspa-

H. *Asparagus* caule herbaceo erecto, foliis setaceis, stipulis paribus. *officinalis*. LINN. *Sp.* 1. *Asparagus sativus*. C. B. *Pin.* 489. BLACKW. *T.* 332. In arenosis. v. g. circa *Berolinum*: ad *Rhenum*.

H. *Asparagus* aculeis solitariis, ramis flexuosis, foliis fasciculatis angulatis muticis deciduis. *albus*. LINN. *Sp.* 6. *Asparagus aculeatus*, *spinis horridus*, C. B. *Pin.* 490. In *Hispania*.

3. floribus nudis planis.

ASPHODELVS. LINN. 379.

Corolla sexpartita. Valvulae sex, basi petaliferae circa germen. Filamenta staminum ex valvulis orta.

H. *Asphodelus* caule folioso, foliis triquetris fistulosis. *luteus*. LINN. *Spec.* 1. *Asphodelus luteus*. C. B. *Pin.* 28. *Hasta regia*. BLACKW. *T.* 233. In *Sicilia*.

H. *Asphodelus* caule nudo, foliis strictis subulatis striatis subfistulosis. *fistulosus*. LINN. *Sp.* 2. *Asphodelus minor*. CLVS. *Hist.* 1. p. 197. In *Europa australi*, *Syria*, *Creta*.

ANTHERICVM. LINN. 380. *Phalangium*.

HALL. TOURN.

Corolla hexapetala. Filamenta staminum subulata. Stigma obtusum.

H. *Anthericum* foliis carnosis subulatis planiusculis. *aloides*. LINN. *Sp.* 5. *Phalangium capense sessile foliis aloeformibus pulposis*. DILL. *Elsh.* *T.*

232. f. 300. Stamina plumosa. Ad Caput bonae Spei.

h. Anthericum foliis carnosis teretibus, caule fruticoso. *frutescens*. LINN. *Spec.* 4. Phalangium Afr. foliis cepaceis, floribus spicatis aureis. BOERH. *Ind. Pl.* II. p. 133. Phalangium Capense caulescens, foliis cepaceis succosis. DILL. *T.* 231. f. 298. Stamina barbata.

^{h. Anthericum annuum}
Anthericum foliis planis, scapo simplicissimo. *Liliago*. LINN. *Sp.* 3. Phalangium radicibus teretibus, caule nudo, foliis gramineis, petiolis florum non ramosis. HALL. *Helv.* 292. n. 1. Phalangium parvo flore non ramosum. C. B. *Pin.* 29. Filamenta glabra. In asperis lapidosis circa arcem Plesi.

Anthericum foliis planis, scapo ramoso, corollis planis. *ramosum*. LINN. *Spec.* 2. Phalangium priori simile, petiolis ramosis. HALL. *l. c.* n. 2. Phalangium parvo flore ramosum. C. B. *l. c.* Flores minores & viridibus lineis notati. In M. *Altenstollberg*.

β. Ornithogalum spicis florum longissimis ramosis. GMEL. *Fl. Sib.* T. I. Tab. 8. Flos totus albus.

ORNITHOGALVM. LINN. 377.

Corolla hexapetala. Filamenta staminum plana, in aliis omnia, in aliis alterna apice trifido.

Ornithogalum scapo anguloso diphylo, pedunculis umbellatis simplicibus, filamentis omnibus subulatis. *luteum*. LINN. *Sp.* 1. Phalangium radice sub-
rotun-

rotunda, foliis & floribus umbellatis. HALL. *Helv.* 193. n. 5. *Ornithogalum luteum.* C. B. *Pin.* 1. In agris circa *Göttingam*. Omnis habitus hanc plantam sic retinet, etsi stamina a definitione generica differant.

H. *Ornithogalum floribus corymbosis, pedunculis scapo altioribus, filamentis alternis emarginatis umbellatum.* LINN. *Sp.* 9. *Ornithog. pedunculis florum imis supremos superantibus.* HALL. *Helv.* 194. *Ornithog. umbellatum medium angustifolium.* C. B. *Pin.* 294. Sponte in *Helvetia* in agris.

H. *Ornithogalum floribus corymbosis, pedunculis scapo humilioribus, filamentis emarginatis. arabicum.* LINN. *Sp.* 8. *Ornithogalum arabicum.* CLVS. *Hist.* II. p. 186. In *Aegypto. Arabia.*

β. floribus aggregatis vagina communi exceptis.

H. *Ornithog. nutans* n. 2. *pag.* 10.

ALLIVM. HALL. LINN. 370. Ceba.

Porrum. Moly.

Petala concavo-erecta. Stigma acutum. Capsula revis, trigona. Spatha bifolia.

I. Staminibus alterne trifidis.

H. *Allium staminibus alterne trifidis, foliis gramineis, capite bulbifero, nucleis in bulbum connatis.* HALL. *opusc. bot.* 331. *de allio.* n. 1. *Allium. sativum.* LINN. *Sp.* 10. C. BAVH. *Pin.* 73. n. *Sicilia.*

H. *Allium staminibus alterne trifidis, capite bulbifero, scapo ante maturitatem contorto.* HALL. *ib.* n. 2. *Allium Scorodoprasum.* LINN. *Sp.* 11. β. *Allium*

Allii genus *Ophioscorodon* dictum quibusdam. I. B. *Hist.* II. p. 539.

Allium staminibus alterne trifidis, capite bulbifero, foliis gramineis leviter ferratis. HALL. *All.* n. 3. RVPP. *Jen.* T. 2. f. 2. Allium *arenarium*. LINN. *Sp.* 3. Allium *mont. latifolium*. CLVS. *Pann.* p. 212. In M. Kishhäuser.

Allium staminibus alterne trifidis, foliis fistulosis, capite sphaerico, bulbifero, radice multiplici. HALL. *All.* n. 4. Allium *vineale*. LINN. *Sp.* 18. Allium *sylvestre*. C. B. *Pin.* 74. Porrum *sylv. vinearum*. C. B. 72. In agris circa *Gottingam*.

H. Allium staminibus alterne trifidis, foliis gramineis, floribus sphaerice congestis, radice sobolifera. HALL. *All.* 5. Allium *Ampeloprasum*. LINN. *Sp.* 1. Allium *sphaerico capite, folio latiore*. C. B. *Pin.* 74. In *Minorca*. In inf. *Holms*.

H. Allium staminibus alterne trifidis, foliis latissimis, capite sphaerico, caule ancipiti. HALL. *All.* n. 6. Allium *nutans*. LINN. *Sp.* 20. Porrum *scapo nudo, ancipiti, antequam floruerit, nutante, foliis ensiformibus, hinc paulo convexioribus*. GMEL. *Fl. Sib.* T. I. p. 55. T. 12. In *Sibiria*.

H. Allium staminibus alterne trifidis, foliis gramineis floribus sphaerice congestis, radice tunicata, cauli circumnata. HALL. *All.* n. 7. Allium *Porrum*. LINN. *Sp.* 2. Porrum *commune*. C. B. *Pin.* 72.

H. Allium staminibus alterne trifidis praelongis capite sphaerico, atropurpureo. HALL. *All.* n. 11.

An

An *Allium sphaerocephalum*. LINN. *Sp.* 14. *Moly atropurpureum*. SWEERT. T. 60. In *Sibiria*.

h. *Allium* staminibus alterne trifidis, caule ad terram ventricoso. HALL. *All. n.* 12. *Allium scapo nudo inferne ventricoso longiore, foliis teretibus*. (Cepa.) LINN. *Sp.* 25. *Cepa vulgaris*. C. B. *Pin.* 71.

2. *Staminibus omnibus simplicibus, subulatis.*

h. *Allium* staminibus simplicibus, caule medio ventricoso. HALL. *All. n.* 13. *Allium scapo nudo, foliis adaequante, foliis teretibus ventricosis*. (*fistulosum*.) LINN. *Sp.* 27. *Cepa oblonga*. C. B. *Pin.* 71.

h. *Allium umbellatum* foliis teretibus radicalibus, scapi longitudine, floribus tubulosis. HALL. *All. n.* 14. *Allium Schoenoprasum*. LINN. *Sp.* 28. *Cepa scutellaria*. TABERN. *ic. p.* 486.

h. *Allium* foliis gramineis, umbella sphaerica, bicolore. HALL. *All. n.* 16. *Comment. Soc. Gott.* T. 11.

Allium radice senescente lignosa transversa. HALL. *All. n.* 19. *Allium scapo nudo ancipiti, foliis linearibus canaliculatis subtus subangulatis, umbella fastigiata*. (*angulosum*.) LINN. *Spec.* 22. *Allium s. Moly foliis Narcissinis*. CLYS. *Pann.* 221, 222. In *M. Altenstullberg*.

β. *Allium senescens*. LINN. *Sp.* 21. HALL. *Helv.* 297. n. 2. *Allium scapo ancipiti, teretiufculo, foliis infirmis, binc paulo convexioribus*. Gmel. *Sib.* p. 53. T. XI. f. 2. In *Sibiria*. Cf. *Helv.* 1227.

Allium

Allium foliis radicalibus latissimis petiolatis, floribus umbellatis. HALL. *All. n.* 21. *Enum. Helv. p.* 297. *Allium ursinum*. LINN. *Sp.* 23. LOB. *ic. p.* 159. *A. sylv. latifolium*. G. B. *Pin.* 74. Abunde in sylvis.

H. *Allium* foliis latissimis, radice oblonga, obducta reticulo. HALL. *All. n.* 20. *Allium Victorialis*. LINN. *Sp.* 4. *A. montanum latifolium maculatum*. G. BAVH. *Pin.* 74. *A. anguinum* CAM. *epit.* 329. In M. Kifhäuser. In Alpibus Helv. & Silviacis.

* H. *Allium* umbellatum, foliis fistulosis, compressis, radice obducta reticulo. HALL. *Comm. Gott. T. II. Tab. IX. p.* 337. In Sibiria.

H. *Allium* foliis caulinis lanceolatis, floribus umbellatis, ex ala bulbiferum. HALL. *All. 22. p.* 382. *Allium*. LINN. *Sp.* 6. *Moly latifolium liliflorum*. G. B. *Pin.* 75.

H. *Allium* foliis radicalibus sessilibus, lanceolatis, floribus umbellatis. HALL. *All. n.* 23. ROYEN. *Lugdb. p.* 57. *Allium Moly*. LINN. *Sp.* 26. *Moly latifolium luteum odore Allii*. G. B. *Pin.* 75. In Hungaria. M. Pyrenaicis.

H. *Allium* foliis teretibus, vagina bicorni, umbella pendula suavepurpurea. HALL. *All. 386. n.* 25. *cum icone*. In Sibiria. Prope Veronam.

Allium inodorum, umbella bulbifera, vagina bicorni, foliis carinatis. HALL. *All. 391. n.* 27. *Tab. 2.* *Allium carinatum*. LINN. *Spec.* 13. *A. sylvestre bicorni, flore ex herbaceo albicante, cum triplici*
in

CHELIDONIVM. HALL. 305.

LVDW. 447.

Omnia prioris, sed stamina non connata, filiqua unilocularis, stigma capitatum, planta vndique succo aureo manat.

Chelidonium foliis subrotundis. HALL. *Helv.* 305.
Chelidonium, *pedunculis umbellatis. minus.* LINN. *Sp.*
1. Chelid. *maius vulgare.* BLACKW. *T. 91.* β. Chelid.
foliis lactiniatis. CLVS. *Hist.* II. p. 203. Vbique
in ruderatis. A.

CHRISTOPHORIANA. HALL. 306.

Actaea LINN. 568.

Calyx quadrifolius, caducus. Petala aliquot exigua, quasi degenerantia stamina. Bacca globosa, stigmate oblique depresso.

Christopheriana. DOD. *Purg.* p. 231. CLVS. *Hist.*
2. p. 86. Actaea racemo ovato, fructibusque baccatis. *spicata.* LINN. *Sp.* 1. Aconitum *bacciferum.*
C. B. *Pin.* 182. In sylvis & ad sepes. P.

CLASSIS. VI:

MVLTISILIQVAE.

In vniversum huius classis plantis stamina sunt multo plura, ac petala, & post florem filiculae, in aliis polyspermae dehiscentes, in aliis monospermae, semine non discedentes.

III. BVTOMVS, LINN. 455. HALL. 299.

Involucrum trifolium umbelliferum, Calycis folia petalis similia colorata. Stamina novem. Siliquae sex distinctae, tubae apice bifido. Folia graminea.

Butomus ^{umbellatus} LINN. *Spec.* 1. HALL. *l. c.* *Juncus floridus maior*. C. B. *Pin.* 112. *Gladiolus aquatilis*. DOD. *Pempt.* 600. In fossis. P.

ALISMA, LINN. 418. Damafonium.

Involucrum trifolium flores colligens. Stamina sex. Capsulae plures in trigonum posita tuba simplici.

Alisma foliis ovatis acutis, fructibus obtuse trigonis. ^{Plantago} LINN. *Spec.* 1. *Damafonium panicula verticillata, petiolis ramosis, siliquis brevissimis*. HALL. *Helv.* 300. *Plantago aquatica*. C. B. *Pin.* 190. In fossis. P.

Alisma foliis gramineis & ovatis, caule repente, umbella pauciflora. *Alisma natans*. LINN. *Spec.* 4. *Damafonium repens, potamogetonis rotundifolii folio*. VAILL. *Act. Par.* 1719, p. 29. T. 4. f. 8. *Damafonium scapo geniculato, reptante, uniflora*. BOEHM. *Fl. Lips.* p. 108. Rarius in inundatis circa *Berolinum*. Capsulae circiter octo. Flores ex spatha nonnullos prodeuntes semper vidi. P.

SAGITTARIA, LINN. 946. HALL. 300.

Monophyta. *Mares*. Stamina viginti quatuor in capitula congestae. *Fem.* intra mares. Siliculae plurimae in globum collectae, Spathae trifoliae.

Sagit-

TRIPETALOIDEAE.

213

Sagittaria foliis sagittatis, acutis ^{*sagittifolia*} LINN. *Spec.* 1. *Sagittaria aquatica. a. latifolia, angustifolia.* C. B. P. 194. Caulis verticillatus. In lacu ad *Seeburg.* P.

STRATIOTES. *Comm. Soc. Gott. T. III. p. 425.*

LINN. 607. *Aloides* BOERH. II.

p. 172.

Diphyta. *Fl. Masc.* Spatha bifolia flores plures colligens. Stamina duodecim, antheris erectis, corona filamentorum lanceolatorum cincta. *Fem.* Spatha bifolia uniflora. Corolla filamentorum circa tubas, ut in mare. Tubae sex triangulares bifidae. Capsula oblonga, sexlocularis.

Stratiotes. *Comm. Gott. l. c. p. 426. sq. Tab. XLX.* Stratiotes *foliis ensiformi-triangularis ciliato-aeuleatis.* *Aloides.* LINN. *Sp.* 1. *Aloe palustris.* C. B. P. 286. In paludibus & stagnis profundioribus circa *Berolinum.* P.

HYDROCHARIS. *Comm. Soc. Gott. T. III.*

p. 431. LINN. 999.

Diphyta. *Mas.* Spatha bifolia triflora. Stamina novem, quorum tria interiora stylum e basi emittunt. *Fem.* Spatha nulla. Filamenta sex, lanceolata, ut in Aloide, posita circa tubas sex bifidas. Capsula nuda sexlocularis. Caulis per flagella repit. Folia reniformia.

Hydrocharis. Morsus ranar. LINN. *Spec.* 1. HALL. *Helv.* 301. *Comm. Gott. l. c. T. XI. f. 1-9.* Nymphaea *alba minor.* C. B. P. 193. *Microleuconymphaea.* BOERH. *H. Lugd.* 11. p. 232. In lacu ad *Seeburg.* P.

H

GLAS-

ORDO. II.

Multifliquae siliculis polyspermis dehiscentibus.

α. Floribus nectario instructis.

FRAXINELLA. HALL. 311. Dictamnus.

LINN. 468.

Calyx quinquefolius. Petala quinque. Filamentis decem staminum glandulae adspersae. Siliculae quinque connatae, rubra vnica.

Fraxinella. HALL. 311. Dictamnus albus. LINN.

Sp. 1. Mat. med. 208. BLACKW. T. 75. α. flore rubro β. flore albo. In *Germania, Gallia, Italia*. P.

AQVILEGIA. LINN. 605.

Petala quinque patentia, totidem nectaria alterne cum petalis posita, infundibuliformia. Stamina plurima. Siliculae quinque.

Aquilegia nectarialis incurvis. vulgaris. LINN. Sp. 1.

Aquil. foliis rotunde lobatis. HALL. H. 310. Aquilegia. BLACKW. T. 409. & auctorum. In sylvis & pomariis. P. *pl. alba, capite purpureo, in minoribus prope folia*
pubescentia

H. Aquilegia nectarialis rectis, staminibus corolla longioribus. Canadensis. LINN. Sp. 3. Aquilegia pumila praecox Canadensis. LOB. ic. 26. In *Virginia, Canada*. P.

ACONITVM. LINN. 603.

Petala quinque inaequalia, supremum galeatum, sub quo in longis pedunculis nectaria duo corniculata. Siliculae tres & quinque.

Aconi-

Aconitum foliis peltatis multifidis villosis, petalo supremo cylindraceo. LINN. *Spec.* 1. HALL. *H.* 312. n. 1. *Aconitum lycoctonum luteum*. C. B. *Pin.* 183. In sylvis v. g. circa Lützelengen. P.

H. *Aconitum* foliis multipartitis, laciniis linearibus incumbentibus, squarrosis. *pyrenaicum*. LINN. *Sp.* 4. HORT. *Vpf.* 152. In Sibiria. Pyrenaeis. P.

H. *Aconitum* foliorum laciniis linearibus superne latioribus linea exaratis. *Napellus*. LINN. *Spec.* 3. HALL. *l. c. n.* 2. *Aconitum coeruleum* f. *Napellus*. B. *Pin.* 183. In Helvetia. Gallia. *In Domo huiusmodi hinc inde f. 2. in maximo hinc inde*

H. *Aconitum* foliis multifidis, laciniis semipartitis, superne latis. *variegatum*. LINN. *Sp.* 2. *Acon. coeruleum minus*. B. *Pin.* *l. c.* CLVS. *Aconit. lycoct.* IX. *Judenbergense*. p. XCVII. In Italia, Bobemia. P.

H. *Aconitum* floribus pentagynis. LINN. *Sp.* 1. *Acon. foliorum laciniis linearibus distinctis ubique eiusdem latitudinis*. LINN. *Mat. med.* 270. HALL. *H.* 313. *Aconit. salutarium* f. *Anthora*. LINN. *Sp.* 5. In alpidibus Helveticis. M. Thury prope Genevam. P.

DELPHINIVM. LINN. *Sp.* 602.

Calyx nullus. Petala quinque inaequalia, supremum in conicum calcar abit, quod simile fere calcar replet modo simplex modo duplex, intra petala in bracteam ovatam abiturum, Silicula una, & tres.

Delphinium nectariis monophyllis, caule subdiviso. *Consolida*. LINN. *Sp.* 1. *Delphinium flore unifiliquo, foliis tenuissime divis.* HALL. *Helv.* 314.

Delphinium segetum. I. R. H. *Consolida regalis arvensis*. B. Pin. 142. Inter segetes A.

H. *Delphinium nectariis monophyllis*, caule simpliciter. *Aiacis*. LINN. *Spec.* 2. *Delphinium sativum*. RIV. T. 123. *Consolida regalis hortensis*. B. Pin. 142. Siliquae villosae. A.

H. *Delphinium nectariis diphyllis*, labellis integris, floribus subfoliariis, foliis compositis linearimultipartitis. *grandiflorum*. LINN. *Sp.* 4. *Hort. Vpf.* 150. *Delphin. perenne*, foliis tenuissime divisis, siliquis villosis. HALL. *Gott.* 94. *Delphin. elatius subincanum perenne*, floribus amplis azureis. AMM. *Ruth. n.* 175. ROLFE. *Hort.* 61. T. 3. In *Siberia*. P.

Delphinium nectariis diphyllis, labellis bifidis apice barbatis, foliis semiquinquelobis, subhirsutis, lobis latis, serratis. *Delphinium elatum*. LINN. *Sp.* 5. *Hort. Vpf.* 151. *Delphinium perenne*, aconiti folio ampliori, floribus coeruleis longioribus pedunculis insidentibus, minus villosum. AMM. n. 174. HALL. *Gott.* 93. *Aconitum lycoctonum flore Delph. I. Siles.* CLVS. XCIV. Folia semiseptiloba, lobis latis, serratis ciliatis. In *Siberia*.

H. *Delphinium nectariis diphyllis*, foliis multipartitis, glabris, laciniis linearibus superne latioribus, pinnatifidis. Pro distincta specie haberi meretur. LINNAEVS autem in *diff. de plantis hybridis*, *Amoen. Acad. Vol. III. p.* illam esse credit plantam hybridam enatam ex *Delphinio elato*, matre, & *Aconito Napello*, patre. Flos prioris speciei. P.

H. Del-

H. Delphinium nectariis diphyllis, foliis palmatis, lobis integris. LINN. Sp. 6. Mat. med. 269. Delphinium orient. platani folio. HALL. Gott. 93. Staphis-agria. B. Pin. 324. BLACKW. T. 265. In Hungaria. Italia. circa Monspolium. P.

Delphinium amblyum

RESEDA. LINN. 207. Luteola.

HALL. 315.

Petala digitata, fissa, ad basin supremi squama nectarifera. Capsula unilocularis inter stylos aperta.

Reseda foliis lanceolatis integris, calycibus quadrifidis. LINN. Sp. 1. Luteola. HALL. l. c. BLACKW. T. 283. Luteola herba salicis folio. BAVH. Pin. 10. Petala tria, Styli tres, stamina plurima. In gramineis. v. g. circa Pless. A.

H. Reseda foliis omnibus trifidis, inferioribus pinnatis. lutea. LINN. Sp. 7. HALL. H. l. c. Reseda foliis rariter pinnatis undulatis. HALL. Gott. 94. Reseda vulgaris. B. Pin. 100. Calyx sexfidus, petala sex, styli tres. In Thuringia & arenosis Europae australioris. A.

H. Reseda foliis pinnatis, floribus tetragynis. alba. LINN. Sp. 6. Reseda foliis pinnatis planis, flore specioso. HALL. Gott. 95. Reseda maxima. B. Pin. 100. Calyx sexfidus, petala sex omnia trifida, stamina undecim. Monspelii. A.

H. Reseda foliis integris, floribus odoratis, calyce corollam vix superante. HALL. Gott. 95. Reseda Aegyptiaca odoratissima. Calyx sexfidus, petala sex, stamina plurima alba. In Aegypto. A. odorata
2. Linn. Spec. 11.

H. Reseda foliis integris trilobisque, calycibus maximis hexaphyllis. *Phyteuma*, LINN. Sp. 8. Reseda minor vulgaris. 1. R. H. 423. Resedae affinis *Phyteuma*. B. Prodr. T. 42. Calyx corolla multo maior, petala sex, styli tres, stamina pauciora, rubella. In Gallia. Italia. A.

PARNASSIA. LINN. 133. HALL. 316.

Squamae quinque intra petala cordatae, singulae 15 stylis globuliferis instructae, Stamina quinque. Capsula quadrangularis, ex apice hians.

Parnassia. ^{par/is/1733} LINN. Sp. 1. ROYEN. 420. HALL. l.c. Gramen Parnassii. LOB. ic. 603. Gramen Parnassii albo simplici flore. C. B. Pin. 309. In pratis viginosis. P.

NIGELLA. LINN. 606. HALL. 316.

Flos nudus. Petala quinque. Squamae octo, petiolatae, bifidae, lineatae; in singularum origine fovea nectarifera, cui lamina lanceolata applicatur. Capsulae quinque aut decem, liberae modo, modo connatae in totidem stylos revolutos abeunt.

Nigella flore foliis nudo, pistillis corollam aequantibus. HALL. Helv. 316. Nigella pistillis quinis, petalis integris, capsulis turbinatis, arvensis. LINN. Sp. 3. Nigella arvensis cornuta. 1. R. H. B. Pin. 145. Inter segetes. v. g. circa Lützelengen. A.

H. Nigella petalis subtricuspidatis, foliis subpilosiss. sativa. LINN. Sp. 2. H. Vps. 154. Nigella capite ♂ folio minore. 1. B. III. 1. p. 208. In Palaestina. Aegypto. A.

H. *Nigella* floribus involucro folioso exceptis. *Damascena*. LINN. *Sp. I. H. Vpf.* 53. *Melanthium sylvestre*. LOB. *ic.* 740. β . Var. flore pleno. Inter *segetes Europae australis*. A.

H. *Nigella* pistillis dens corolla longioribus. *orientalis*. LINN. *Sp.* 5. HALL. *Gott.* 96. *Nigella orientalis*, flore flavo, semine alato plano. TOURN. *Cor.* 19. Capsulae octo ad decem, Circa *Alepum*. A.

GARIDELLA, LINN. 507.

Nigellae proxime adfinis. Sed stamina decem, capsulae tres, styli vix vili, petala exigua.

H. *Garidella*. LINN. *Sp. I.* GARIDEL. *Prov. T.* 39. *Nigellastrum raris & foeniculaeae foliis*. MAGN. *Hort. T.* 143. *Nigella cretica*, folio foeniculi. C.B. *Prodr.* 75. In *Provincia. Creta*. A.

HELLEBORVS. LINN. *gen. ed. prior.* 462. HALL. 317. *Trollius Isopyrum*. LINN.

Flos nudus. Intra orbem petalorum neetaria conica, tubulata. Capsulae plurimae, compressae, liberae.

H. *Helleborus* vaginis pallidis, vix serratis, flore virente. HALL. *Helv.* 317. *n.* 2. *Helleb. caule multiflora, foliis fo, foliis pedatis. (foetidus.)* LINN. *Sp.* 4. *Helleb. niger foetidus*. B. *Pin.* 185, *Helleboraster*. BLACKW. *T.* 75. In *Germania meridionali. Helvetia*. P.

Helleborus vaginis foliaceis, digitatis, flore virente. HALL. *l. c. n.* 1. *Helleb. caule multiflora, foliis fo, foliis digitatis. (viridis.)* LINN. *Sp.* 3. *Hel-leb.*

leb. *niger bortensis, flore viridi.* B. Pin. 185. Circa Grubenbagen. Viennam. P.

H. Helleborus scapofubunifloro subnudo, foliis pedatis. *niger.* LINN. Sp. 2. Helleb. *niger flore roseo.* B. Pin. 186. Elleborus *niger, legitimus* GLVS. Hist. I. p. 275. In Austria. Italia. P.

H. Helleborus caule plerumque unifloro, petalis et filiquis plurimis. HALL. Gott. 97. Helv. 218. *Trollius corollis conniventibus, nectariis longitudine flaminum.* LINN. Sp. 1. Ranunculus *flore globofo.* DOD. Purg. 287. Trollius flos GESN. Hort. 144. Petala circiter 14. nectaria novem. In montibus Helvetiae. P. *n. viridis*

H. Helleborus foliis repetito multifidis, nectariis planis. HALL. Gott. 97. Helleborus *folio fumariae* AMM. Ruth. 74. T. 12. *Isopyrum stipulis subulatis, petalis acutis.* LINN. Sp. 1. In Sibiria. A.

β. nectario carentibus.

PAEONIA. LINN. 600.

Calyx quinquefolius, inaequalis, quinque petala; capsulae duae, tres, ovatae.

H. Paeonia foliis difformiter lobatis. HALL. Helv. 311. Paeonia *officinalis.* LINN. Sp. 1. Paeonia *femina* C.B. BLACKW. T. 65.

β. Varietas *flore pleno hort.* Sponte in montibus Helvetiae.

H. *Paeonia laciniis foliorum linearibus*. Folia decomposita, pinnulis ultimis lineari-lanceolatis, nervo depresso; Flos nido folioso excipitur. Petala egregie phoenicea. Planta a priori diversissima & MARTINO missa, in *Vkrainia* habitat. P. *lanceolata, foliis linearibus multipartitis* L. *sup. p. 748*.
CALTHA. LINN. 623. **Populago** HALL. TOWN.

Petala quinque sine calyce. Capsulae stellatim dispositae.

Calcha palustris. LINN. *Spec. 1. C. B. Pin. 276*.
Populago. HALL. *H. 219. TABERN. 750. c. 16*.
 I. R. H. Ubique in vdis. Varietas *fl. pleno bott.*

ORDO. III.

Multisiliquae siliculis non dehiscentibus monospermis, receptaculo communi exceptis. *Gymnospermiae*.

a. Petalis & staminibus ex ambitu receptaculi ortis.

LIRIODENDRON. LINN. 609. **Tulipifera**.

Calyx trifolius petala sex. Stamina viginti quatuor in sinibus squamarum, stigmate globoso, in receptaculo conico.

H. *Liriodendron*. LINN. *Sp. 1. H. Vps. 154*.
Tulipifera arbor Virginiana HERM. *Lugdb. 613*.
 T. 613. CATESB. *Car. 1. p. 48. T. 48*. BUTTNER. *H. Canon. In America septentrionali. Fr.*

MYOSVRVS. LINN. 355.

Calycis folia quinque ex vngue angusto repanda. Petala totidem fere similia, minora. Stamina ad duodecim. Semina in receptaculo longissimo.

Myosurus. LINN. *Sp. p.* 284. HALL. *Gott.* 98. Cauda muris. LOBEL. *ic.* 40. DODON. *Pempt.* 112. In collibus apricis v. g. prope *Göttingam* in *M. Hagen*, A.

ADONIS. LINN. 618.

Petala octo aut duodecim. Semina in receptaculo longissimo.

Adonis floribus octopetalis. fructibus subcylindricis. ^{*autimralis*} LINN. *2 Sp. p.* 12. HALL. *H.* 318. *α. flore phoeniceo. β. flore citrino.* Inter segetes *γ. petalis brevioribus flore rubro, maior.* hort. A.

H. *Adonis flore dodecapetalo*, fructu ovato. *vernalis* LINN. *Sp. 2.* *Adonis perennis dodecapetalos.* HALL. *H.* 319. *Helleborus niger tenuifolius, Bupthalmi flore.* C. B. *Pin.* 186. Circa *Halberstadt*. In *Borussia*, P.

RANUNCVLVS. LINN.

Calycis folia & petala quinque, ad singulorum basin fovea nectarifera.

Ranunculus foliis ovato lanceolatis, petiolatis, caule declinato. *flammula*. LINN. *Sp. 1.* HALL. *H.* 322. *Ranunculus longifolius palustris minor.* B. P. 180. *Flammula-Ranunculus.* DOD. *Pempt.* 432. In vdis vbique. P.

Ranun=

Ranunculus foliis lanceolatis, sessilibus, caule erecto. *Lingua*, LINN. Sp. 3. HALL. H. l. c. n. 2. Ranunculus *longifolius palustris maior*. C. B. Pin. 34. In lacubus ad Seeburg & Denckershausen. P.

Ranunculus foliis inferioribus palmatis, summis digitatis, fructibus oblongis. *sceleratus*, LINN. Sp. 15. HALL. Helv. 323. n. 4. Ranunculus *palustris apertifolius, laevis*. B. Pin. 180. BLACKW. T. 259. In offis & vdis. A.

Ranunculus foliis radicalibus reniformibus crenatis incisus: caulinis digitatis, linearibus, caule multifloro. *auricomus*, LINN. Sp. 13. Ranunc. *nemorosus folio subrotundo*. B. Pin. 178. In sylvis. P.

Ranunculus foliis omnibus quinatis lanceolatis inciso-serratis. *aconitifolius*, LINN. Sp. 16. Ranunc. *mont. aconiti folio albus*. C. B. Pin. 182. In sylva Hercynica circa M. Bructetum. P.

h. Ranunculus foliis ternatis biternatisque, foliis trifidis incisus, caule inferne ramoso: *asiaticus*, LINN. Sp. 18. Ranunculus *grumosa radice, ramosus*. B. Pin. 181. Varietates plurimae ex colore flore pleno in hortis. In Asia. P.

h. Ranunculus foliis compositis, caule simplicifloro subnudo unifloro, radice tuberosa. An Ranunculus *mont. leptophyllus, asphodeli radice*. COL. Ephr. 1. p. 312. T. 311. Icon egregie conveniens in horto flore pleno adest. P. *charophyllus* ^{*bulbosus*} *sp. 262. 30.*

Ranunculus radice simplici globosa. ^{*bulbosus*} LINN. Cliff. 30. HALL Helv. 323. Gott. 101. Ranunculus *lycibus retroflexis, pedunculis sulcatis, caule erecto*

Ho, foliis compositis. LINN. *Sp.* 25. *Ranunc. bulbosus.* LOB. *ic.* 666. P.

(β.) *Ranunculus* caule ex geniculis radicato, foliorum lobis petiolatis. HALL. *Helv* 324. n. 6. *Ranunculus calycibus patulis, pedunculis sulcatis, folionibus repentibus, foliis compositis.* LINN. *Sp.* 26. *Ranunc. prat. repens hirsutus.* B. *Pin.* 179. Num Varietas? Uterque in pratis & gramineis. P.

Ranunculus caule erecto multifloro, folio pentagono, lobis sessilibus ferratis. HALL. *H.* 324. n. 7. *Ranunc. acris.* LINN. *Sp.* 28. *Ranunc. pratensis, erectus, acris.* B. *Pin.* 178. In pratis. P.

(β.) *Ranunculus* priori simillimus, foliorum lobis sessilibus laciniatis. HALL. *Gott.* 101. *Ranunculus polyanthemor.* LINN. *Sp.* 27. LOB. *ic.* 666. P.

(γ.) *Ranunculus* foliis tripartitis acute sinuatis tomentosis ^{lanuginosis}. LINN. *Sp.* 29. *Ranunc. magnus valde hirsutus, flore luteo.* 1. B. *Hist.* 3. p. 417. In sylvis. Num varietates? P.

Ranunculus feminibus aculeatis, foliis superioribus decompositis linearibus. *arvensis.* LINN. *Sp.* 31. HALL. *H.* 323. n. 5. *Ranunculus arvensis echinatus.* B. *Pin.* 179. In agris. A.

H. *Ranunculus* feminibus aculeatis, foliis simplicibus palmatis incis. *muricatus.* LINN. *Sp.* 32. *Ranunc. creticus echinatus latifolius.* P. ALP. *Exot.* 262. B. *Pin.* 180. In Europa meridionali. A.

H. *Ranunculus* ramis unifloris, foliis trifidis, filiquis incurvis. HALL. *Gott.* 99. *Ranunc. foliis filiformi-ramosis, calycibus appendiculatis.* LINN. *Sp.*

35. *Ranunc. ceraticeus*, *feminibus falcatis*, in spicam adactis. MORIS. II. 8. 4. T. 28. f. 22. Stamina pauca ad decem vsque. In Europa australi. A.

Ranunculus foliis submersis capillaceis, emeris peltatis. *aquatilis*. LINN. Spec. 37. HALL. Helv. 328. n. 16. *Ranunc. aquat. folio rotundo & capillaceo*. B. Pin. 180. In stagnis & inundatis.

β. *Ranunculus foliis capillaceis*, circumscriptione vaga. HALL. t. v. n. 17. In aquis parum velocibus v. g. im Rasemühlen-Wasser.

γ. *Ranunculus foliis capillatis*, circumscriptione rotunda. HALL. l. c. n. 18. *Ranunculoides folio circinato, tenuissime diviso*. VAILL. In aquis quietis. v. g. bey Schürzenhof.

δ. *Ranunculus foliis capillaceis*, circumscriptione oblonga. HALL. l. c. n. 19. In aquis rapidioribus. v. g. in der alten Leine. Varietates esse videntur LINNAEO, LUDWIGIO, BOEHMERO, nec HALLERVS plane dissentit.

Ranunculus foliis linearibus, brevissime bifurcatis. HALL. Rupp. 104. MAPP. Atlas. 263. cum &c. Num satis a prioribus differt?

Ranunculus foliis cordatis angulatis petiolatis. LINN. Spec. 9. *Ficaria*. HALL. Helv. 321. *Chelidonium minus*. BLACKW. T. 54. *Chelidonium rotundifolia minor*. BAVH. Pin. 167. Ad sepes & in umbrosis. Calyx trifolius, petala octo. P.

Ranunculus foliis trifoliis. *Chelidonium rotundifolia minor*. BAVH. Pin. 167. 2 p.

HEPATICÆ. DILL. LINN. ed. pr. 457. Trinitas

HALL. 320. *Anemones spec.* LINN.

ed. alt.

Calyx trifolius. Flos bis tripetalos. Receptaculum papposum.

Hepatica. LINN. *Vps.* 155. *Mat. med.* 277.

BLACKW. *T.* 207. Trinitas. HALL. *l. c.* *Anemone*

foliis trilobis integerrimis. ^{Hepatica} LINN. *Sp.* 1. Vbique

in dumetis. β . flore pleno, albo, rubro, coeruleo hortens.

ANEMONE. LINN. 614. Pulsatilla.

Involucrum foliaceum a flore remotum. Petala duorum triumve ordinum.

Anemone femine levi, petalis quinis, subrotundis, luteis. HALL. *Helv.* 329. *n. 1.* *Anemone ranunculoides*. LINN. *Spec.* 17. Ranunculus *nemorosus luteus*. B. *Pin.* 178. Primo vere in sylvis. P.

Anemone femine levi, petalis senis ellipticis. HALL. *H.* 330. *n. 2.* *Anemone nemorosa*. LINN. *Spec.* 15. Ranunculus *phragmites, vernus, albus*. I. R. H. In sylvis. P.

Anemone pedunculo nudo, feminibus seta plumosa terminatis. LINN. *Cliff.* 224. *sylvestris*. *Sp.* 10. *Anemone sylvestris alba maior*. B. *Pin.* 176. In sylva prope *Hessendreisoh* & in monte super *Lützelengen*. P.

H. *Anemone* caule ramoso, petalis lanceolatis. ROYEN. *Lugdb.* 488. *Anemone pedunculis alternis longissimis, fructibus cylindricis, seminibus*
hiv-

hirsutis muticis. (Virginiana.) LINN. Sp. II. HERM. Parad. T. 18. In Virginia. P.

β. Priori similis, sed constanter omnibus partibus altera duplo maior, femina villosissima, serius semper floret.

Anemone tubis caudatis, foliis omnibus duplicato pinnatis: pinnulis remotis. HALL. Helv. 331. n. 5. Anemone foliis caulinis ternis connatis supradecompositis multifidis, seminibus hirsutis caudatis. (*alpina*.) LINN. Sp. 8. Pulsatilla flore albo. B. Pin. 177. In M. Bruclero. P.

h. Anemone tubis caudatis, foliis tenuissime divisis, flore connivente nutante. HALL. Gott. 103. Pulsatilla foliis decompositis pinnatis, flore pendulo, limbo reflexo. LINN. Cliff. 223. Anemone LINN. Sp. 6. Pulsatilla flore minore nigricante. B. Pin. 177. In campis arenosis v. g. circa Berolinum. P.

h. Anemone (*Pulsatilla*) foliis simpliciter pinnatis, foliolis lobatis, flore erecto. LINN. Fl. Suec. edit. prior. 448. Anemone *vernalis*. LINN. Sp. 4. Pulsatilla involucri piloso nitido Fl. Lapp. 223. Pulsatilla *apii folio*, *vernalis*. B. Pin. 177. HALL. Helv. 332. cum icone. Petala villosa, tubae caudatae. P.

CLEMATIS. LINN. 616.

Petala quatuor. Antherae lateri filamentorum adnatae. Styli subulati. Folia coniugata.

Clematis foliis pinnatis, foliolis cordatis, inaequaliter incis, serratis. *Vitalba*. LINN. Sp. 5. HALL. H. 334. Clematis *sylvestris latifolia*. B. Pin. 300. In sylvis & sepibus. P.

H. Clematis foliis compositis decompositisque, foliolis quibusdam trifidis, *Viorna*. LINN. *Sp.* 2. Flammula *audens*, flore violaceo, clauso. DILL. *Elth.* 144. T. 118. f. 444. Clematis *purpurea repens*, petalis florum coriaceis. RAII. *Hist.* 1928. 3. Variet. flore pleno. In *Virginia*. P.

H. Clematis foliis pinnatis, foliolis ovato lanceolatis integerrimis, caule erecto. *recta*. LINN. *Sp.* 8. Clematis f. Flammula *surrecta alba*. I. B. II. p. 127. Flammula. CLVS. *Hist.* I. p. 124. In *Austria*, *Montspeli*. P.

H. Clematis foliis simplicibus ovato lanceolatis, *integrifolia*. Clematis *coerulea erecta*. B. *Pin.* 380. Clematis *coerulea Pannonica*. CLVS. *Hist.* I. p. 123. P.

ORD. IV

SENTICOSAE.

In univcrsum his omnibus, quas fructus similitudo cum prioribus iungit, petala quinque, stamina plurima ex calyce orta, radix perennis, & vires adstringentes. Praeterea Geum, Comarum, Potentilla, Fragaria, & Alchimilla habent calycem semidecempfidum aut semioctofidum, laciniis alternis minoribus, omnes autem stylos lateri germinis insertos.

GEVM. LINN. 561. Caryophyllata. TOVRN.

151. HALL. 335.

Styli longi, geniculati, pilosi.

Geum floribus erectis, fructu globoso, aristis vncinatis nudis, foliis lyratis. *urbanum*. LINN. *Sp.* 2.

Caryo-

*Caryophyllata pinnis paucioribus rhomboideis, tunc
terum sine uncinato.* HALL. l. c. n. 1. *Caryoph.
vulgaris.* B. Pin. 321. BLACKW. T. 253. Ad se-
pes. P.

β. *Caryophyllata Canad.* foliis acutioribus flore
luteo maiori. JVSSIEV.

H. Geum floribus erectis fructu globoso, aristis
uncinatis nudis, foliis ternatis. *Virginianum.* LINN.
Sp. 1. *Caryophyllata Virginiana, albo flore minore, ra-
dice inodora.* HERM. Parad. T. III. In Virginia. P.

Geum floribus nutantibus, fructu oblongo, ari-
stis plumosis. *rivale.* LINN. Sp. 3. *Caryophyllata
pinnis frequentioribus, extrema semisrisida, flore
connivente nutante.* HALL. Helv. 336. *Caryoph.
aquatica, nutante flore.* B. Pin. 321. I. R. H. In
pratis humidis & ad rivulos. β. flore pleno in
horto. P. H.

COMARVM. LINN. 563.

Petala segmentis calycis rubri triplo minora. Sty-
li breves. Receptaculum carnosum.

*Comarum, ^{palustre} LINN. Sp. 1. HALL. H. 337. Quin-
quefolium palustre rubrum.* B. Pin. 325. In vdis v.
g. ad lacum prope Seeburg. P. in Sylv. Cur. repr. And.

POTENTILLA. LINN. 559. HALL. 338.

Tormentilla LINN. Quinquefolium. T.

Calycis quinque segmenta patentia, quinque
circa fructum maturum se complicantia. Styli
breves.

H. *Potentilla foliis pinnatis, caule fruticoso. fruticosa.* LINN. *Sp.* 1. *Potentilla fruticosa, foliis ternatis, medio trifido.* HALL. *Gott.* 106. *Pentaphylloides rectum frutescens.* WALTH. *T.* 17. In Anglia, Sibiria. Fr.

H. *Potentilla foliis pinnatis, pinnis perangustis, quinquefidis.* HALL. *Gott.* 105. *Potentilla foliis bipinnatis, segmentis integerrimis, distantibus, subtus tomentosis, caule decumbente. (multifida.)* LINN. *Sp.* 4. In Sibiria. Fr.

Potentilla foliis pinnatis serratis, caule repente. Anserina. LINN. *Sp.* 2. *Potentilla repens, scapo nudo uniflora, pinnis plurimis confertis, minoribus intermixtis.* HALL. *H.* 338. n. 2. *Anserina* BLACKW. *T.* 6. Vbiq̃ ad vias. P.

H. *Potentilla foliis pinnatis, hirsutis, pinnis tredecim ovatis, crenatis, decurrentibus.* HALL. *Gott.* 106. P.

H. *Potentilla foliis pinnatis, pinnis septenis, subrotundis, crenatis, albo flore.* HALL. *l. c.* *Potentilla foliis pinnatis alternis, foliolis quinis ovatis crenatis, caule erecto. (rupestris.)* LINN. *Sp.* 6. *Pentaphylloides erectum.* 1. B. *Hist.* II. p. 398. In Marchia. In monte Hundsrück. P.

H. *Potentilla foliis novenis palmatis, in apice serratis.* HALL. *l. c.* Folia glabra, nervosa, in solo apice acute serrata, cuneiformia. P.

H. *Potentilla foliis radicalibus septenatis cuneiformibus serratis, caulinis ternatis, ramis filiformibus decumbentibus.* LINN. *Cent. I. Plant. n.* 35. ? Folia

ex stipulis amplexicaulibus digitatis palmata, ampla, media biuncialia, obverse ovata, in toto ambitu magnis dentibus profunde serrata, pilosa. Calyx villosus segmentis alterne integris & profunde trifidis, corollae magnitudine. Petala flava. P.

H. *Potentilla foliis digitatis lanceolatis, serratis, utrinque subpilosis, caule erecto*. ^{recta} LINN. *Sp.* 10. *Quinquefolium maius*. I. TABERN. 351. *Quinquefolium rectum luteum*. B. *Pin.* 325. *Quinquefolium mont. erectum, hirsutum, luteum*. GARIDEL. *T.* 33. Folia quinata, petala sulphurea. P.

H. *Potentilla foliis quinatis, palmatis, sericeis, serratis*. HALL. *Gott.* 107. *Potentilla alba*. LINN. *Sp.* 16. *Quinquefolium album maius*. B. *Prodr.* 325. Pedunculi florum ex radice. In *Marchia Brandeburgensi*. P.

Potentilla foliis quinatis, longe petiolatis, caule repente. HALL. *H.* 340. n. 9. *Potentilla reptans*. LINN. *Sp.* 17. *Quinquefolium maius repens*. B. *Pin.* 325. BLACKW. *T.* 454. Caulis ex intervallis radículas emittit. In muris aliisque locis argillofis. P.

Potentilla foliis quinatis, obtuse in apice serratis, petalis cordatis. HALL. *H.* 341. n. 6. *Quinquefolium minus repens luteum*. I. R. H. C. B. l. c. *Quinquefolium minus*. TABERN. p. 123. Primo vere locis ficcis. P.

Potentilla foliis quinatis angustissimis, subtus tomentosis, caule erecto. HALL. *H.* 341. n. 10. *Potentilla argentea*. LINN. *Sp.* 11. *Quinquefolium folio argenteo*. B. *Pin.* l. c. *Quinquefolium petraeo-*

um maius. TABERN. p. 122. In muris & locis
ficcis. P.

Potentilla foliis quinatis, flore tetrapetalo, cau-
le recto. HALL. H. n. 11. Tormentilla *caule erecto*,
LINN. Sp. 1. Tormentilla ^{apocla}. LINN. Mat. med. 249.
BLACKW. T. 445. officin. In sylvis & locis steri-
libus. P.

H. Potentilla foliis ternatis hirsutis, caule erecto
umbellifero. HALL. Gott. 108. An Pentaphylloides
maius erectum, fragariae folio, flore luteo. MOR-
RIS. Hist. 2. p. 193. S. 11. T. 20. f. 2. ? Icon sa-
tis convenit. Flores parvi inter folia latentes, in um-
bellas dense congesti. Caulis & folia pilosa. A.

H. Potentilla procumbens, foliis ternatis, hirsu-
tis. HALL. Gott. l. c. Potentilla *Norwegica*. LINN. Sp.
19. Potentilla *foliis ternatis incis, caule diffus*.
CLIFF. 193. In Norwegia & Canada. P.

H. Potentilla foliis ternatis, ampliter crenatis,
subtus tomentosis. HALL. Gott. l. c. Potentilla fo-
liis *ternatis incis, subtus tomentosis, caule adscen-*
^{nica}*dente*. LINN. Sp. 20. P.

FRAGARIA. LINN. 558.

Stamina circiter viginti. Receptaculum pulpo-
sum, molle, esculentum.

Fragaria flagellis reptans. *vesca*. LINN. Sp. 1.
Mat. med. 245. HALL. H. 243. Fragaria vulgaris.
B. P. 326. BLACKW. T. 77. Vbique in sylvis.
Varietas α . frequentibus in placenta pilis & peduncu-
lis lanuginosis in Saxonia; β . *bortensis* fructu maiori.

Fragaria foliis; caule decumbente prostrato. ALCHI-
fing. Lexis. Ann. 1797. 12. 216. *Fragaria*
fragella nulla.

ALCHIMILLA. HALL. DENN. 153.

Aphanes, LINN. 154.

Calyx semioctofidus. Petala nulla. Stamina quatuor. Styli ad basin germinis inserti. Semen vnum aut duo in fundo calycis.

Alchimilla foliis integris stellatis polygoniis. HALL. *Helv.* 184. Alchimilla foliis lobatis, (*vulgaris*.) LINN. *Sp.* 1. Alchimilla. BLACKW. *T.* 72. *vulgaris*. B. *Pin.* 319. In sylvis, P.

Alchimilla foliis mollibus semitrilobis, lobis incis. HALL. *Gott.* p. 31. Aphanes. LINN. *Spec.* 1. Perchepier. HALL. *Helv.* p. 184. In arvis abunde. A.

RVBVS. LINN. 557.

Calyx quinquepartitus. Styli lateri germinis inserti. Acini multi monospermi in baccam compositam congesti.

Rubus foliis ternatis nudis, caule aculeato. *caesius*. LINN. *Sp.* 4. HALL. *H.* 344. n. 2. Rubus repens, fructu *caesio*. B. *Pin.* 479. In agris & dumetis v. g. circa *Graezelii* fabricas, P.

Rubus foliis ternatis & quinatis, subrus tomentos; fructu rubro, villoso. HALL. *H. l. c.* n. 3. Rubus Idaeus. LINN. *Sp.* 1. BLACKW. *T.* 289. Rubus Idaeus α. *spinosus* β. *laevis*. B. *Pin.* 479. In sylvis v. g. circa *Pless*, P.

Rubus foliis ternatis & quinatis, costa spinosa, fructu nigro & levi. HALL. *H. n.* 4. Rubus fruticosus. LINN. *Sp.* 5. Rubus vulgaris fructu nigro. B. *Pin.* 479. BLACKW. *T.* 45. β. Varietas *fructu albo*. Ubique in dumetis & sylvis, P.

• *Rubus laxifolius*, foliis ternatis nudis, ROSA. *fragilis* septentrionalis habundans. Lin. 151. n. 17.

ROSA. LINN. 556.

Calyx carnosus, quinque segmentis coronatus.
Germina multa in fundo calycis, qui in fructum
mollem, coloratum abit.

Rosa aculeata, foliis odoratis subtus rubiginosis,
sp. nov. in v. g. 175. in 475.
 WALL. *Helv.* 350. (*eglantaria*). LINN. *Sp.* 2. Rosa
sylvestris, foliis odoratis. B. *Pin.* 483. In sterilibus
 ad vias v. g. versus *Pless.* Fr.

Rosa *Eglensis* β . Varietas flore flavo, hort. *Distincta* *species*: *Rosa*
sp. nov. 14. 1729 *fructu* *vide* *Rich.* *Ed. 8.* *p. 11.* *the* *May* *v. 4.* *p. 476.* *249 F*
 Rosa foliis utrinque villosis, fructu spinoso

Rosa foliis vtrique villosis, fructu spinoso.
HALL. H. 350. n. 8. *villosa*. LINN. Sp. 3. Rosa
sylvestris pomifera maior. B. Pin. 484. Passim in
sepibus. Fr.

Rosa fructu levi, calycis foliolis divisis. HALL, H.
349. Rosa caule aculeato, petiolis inermibus, caly-
cibus semipinnatis. (canina). LINN Sp. 4. Rosa
canina sylvestris. BLACKW. T. 8. Vbique ad se-
pes. Fr.

H. Rosa caule aculeato, pedunculis hispida, calycibus semipinnatis glabris. centifolia. LINN. Sp. 6. Rosa *multiplax media*. B. Pin. 482. Fr.

H. Rosa caule aculeato, pedunculis laevibus, calycibus semipinnatis glabris. alba. LINN. Spec. 7.
Rosa alba vulgaris maior. B. L. c. In hortis. Fr.

CLASSIS. VII

POMIFERAE.

ORDO I.

Pomiferae umbilicatae.

Petala & stamina, ut in priori classe, ex calyce oriuntur. Germen infra receptaculum floris. Con-
ceptaculum seminum pulposum ex apice calycem
persistentem semiquinquefidum praefert.

RIBES. LINN. 247.

Petala calycis segmentis minora. Stamina quin-
que. Tuba bifida. Bacca unilocularis.

Ribes inerme racemis glabris pendulis, floribus
planiusculis. *rubrum*. LINN. *Sp.* 1. *Mat. med.* 99.
HALL. *H.* 346. Ribes *vulgare acidum*. BLACKW.
285. *Johannisbeeren*. Sponte in *Helvetia*, circa
Altdorfium, & regionibus septentrionalibus. β. Va-
rietas hortensis fructu rubro & albido. Fr.

Ribes inerme floribus planiusculis, racemis ere-
ctis. HALL. *l. c.* n. 2. Ribes *alpinum*. LINN. *Spec.* 2.
Ribes *alpinum dulce*. 1. B. *Hist.* 2. p. 98. In sylvis
Hercynicis super *Stollberg*. Fr.

Ribes inerme, foliis & nigro fructu maximis,
olentibus. HALL. *Helv.* 347. n. 4. Ribes *inerme*,
racemis pilosis, floribus oblongis. (nigrum.) LINN. *Sp.*
1. *Mat. med.* 100. *Grossularia non spinosa, fructu*
nigro. B. *Pin.* 455. In pago *Rösdorf* ad rivum. Fr.

Ribes ramis aculeatis erectis, fructu glabro. LINN.
Hist. 82. HALL. *H.* 346. n. 1. Ribes *una crispa*,
LINN.

LINN. *Sp.* 7. *Grossularia sylvestris*. C. B. BLACKW. T. 277. Variet. hort. Sponte rarius ad sepes. Fr.
fr. Rubus Grossularis

MYRTVS. LINN. 543.

Calyx erectus. Tuba simplex. Bacca trilocularis.

H. Myrtus floribus solitariis involucro diphylo. *communis*. LINN. *Sp.* 1. *Mat. med.* 224. α . Myrtus minor vulgaris. Tarentina. β . Myrtus latifolia Baetica. B. Pin. 469. BLACKW. 114. In Europa australi. Fr.

SORBVS. HALL. LINN. 548. *Crataegus* L. 547. *Mespilus*. L. 549.

Stamina viginti circiter. Tuba duae, tres, quinque, & totidem semina in bacca rotunda umbilicata.

Sorbus foliis pinnatis vtrinque glabris fructu minimo. *aucuparia*. LINN. *Sp.* 1. HALL. *Helv.* 250. *Sorbus sylvestris*. Tab. 1020. *Sorbus torminalis*. BLACKW. 173. *Vogelbeerbaum*. Ubique in sylvis. Tubae tres. Fr.

Sorbus (*Crataegus*) folio septangulo, subtus hirtuto. HALL. *H.* 354. *Crataegus foliis cordatis acutis*, lacinulis acutis serratis. LINN. *Cliff.* 187. *torminalis*. LINN. *Sp.* 2. C. B. & *varior. auctorum*. In sylvis passim. Tubae duae. Fr.

Sorbus (*Crataegus*) spinosa, foliis obtusis, saepe trifidis. HALL. *H.* 354. *Crataegus foliis obtusis, bitrifidis, serratis*. (*Oxyacantha*.) LINN. *Sp.* 8. *Mespilus sylv. apii folio*, spinosa. B. Pin. 454. Spina alba.

alba. BLACKW. T. 149. Ad sepes. Tubae duae & tres. Fr. — *etiam Tubae 1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100.*

H. Sorbus (*Mespilus*) inermis, foliis lanceolatis integerrimis subtus tomentosis, calycibus acuminatis *Germanica*. LINN. *Sp.* 1. *Mespilus Germanica folio laurino non ferrato*. B. *Pin.* 453. *Mespilus* BLACKW. 154. In hortis hic colitur. Tubae quinque. Fr.

H. Sorbus (*Mespilus*), foliis ovatis integerrimis. *Cotoneaster*. LINN. *Spec.* 6. HALL. *H.* 352. n. 2. *Chamaemespilus Gesneri*. CLVS. *Hist.* 1. p. Tubae duae, tres, & quinque. Fr.

PYRVS. LINN. 550. Malus. T. 466.

Cydonia. 405.

Stamina viginti. Tubae quinque. Pomum carnosum in quinque loculis membranaceis semina oblonga fovet.

Pyrus foliis ferratis, pomis basi productis. LINN. *Sp.* 1. HALL. *H.* 351. *Pyrus sylvestris*. B. *Pin.* 439. BLACKW. 453. In sylvis. Fr.

β. Pyri domesticæ varietates *bortenses* plurimae.

Pyrus foliis ferratis, pomis basi concavis. *Malus*. LINN. *Sp.* 2. HALL. *H.* 351. *Malus sylvestris*. B. *Pin.* 435. BLACKW. 178. In sylvis. Fr.

β. Mali domesticæ varietates *bortenses*.

H. Pyrus foliis integerrimis, (calyce ferrato). *Cydonia*. LINN. *Sp.* 4. *Malus Cotonea*. Fr.

PVNICA. LINN. 544. T. 407.

Calyx coloratus. Tuba simplex. Pomum magnum nouem loculis instructum.

H. *Punica*. *Granatum*. LINN. *Sp.* 1. *Malus Punica*. B. *Pin.* 438. *Granata*. BLACKW. 145. β . *Balaustia fl. pleno* B. *l. c.* BLACKW. 97. In *Europa australi*. Fr.

PSIDIUM. LINN. 541. Guaiava.
T. 443.

Calycis segmenta ovata. Stylus subulatus, longissimus. Bacca maxima unilocularis.

H. *Psidium*. *Guaiava*. LINN. *Sp.* 1. *Psidium ramis tetragonis*. H. *Vpf.* 122. *Guaiava alba a. acida, β . dulcis*. In *India*. Fr.

ORDO. II.

Pomiferae non umbilicatae.

Omnia prioris. Sed ovarium intra calycem cavitatem intermedia remotum, deciduum. In fructu carnosus nux unilocularis. Tuba simplex.

PRVNVS. LINN. 546. Armeniaca. Cerasus.
Laurocerasus. Padus. T.

Calyx campanulatus. Stylus filiformis. Fructus oblongus aut subrotundus simili nuce.

Prunus spinosa foliis lanceolatis. LINN. *Sp.* 10. HALL. *Helv.* 355. *Prunus Sylvestris*. B. *Pin.* 444. BLACKW. *T.* 305. Vbique in collibus apricis. Fr.

H. Pru-

H. Prunus pedunculis simplicibus foliis lanceolato-ovatis convolutis, domestica. LINN. Sp. 9. Chiff. 186. Mat. med. 232.

β. Pruni domesticae varietates *bortenses*. In hac & priori fructus ovatis. Fr.

Prunus pedunculis simplicibus, foliis ovato-lanceolatis, conduplicatis. Cerasus. LINN. Sp. 8. Cerasus foliis ovato lanceolatis. HALL. Fluv. 156. n. 1. Cerasus *sylvestris*. C. B. Sponte in sylvis. Fr.

β. Cerasi *bortensis* varietates. Fructus subrotundus.

H. Prunus floribus sessilibus, foliis subcordatis, Armeniaca. LINN. Sp. 6. Mala Armeniaca. B. Pin. 482. Fr.

H. Prunus floribus racemosis, foliis sempervirentibus dorso biglandulosis. Laurocerasus. LINN. Sp. 4. Padus *sempervirens*, foliis solidis. HALL. Gott. 112. Laurocerasus. CLVS. Hist. 1. p. 4. Ex Oriente ad nos advenit. Fructus subrotundus, calyx non deciduus. Fr.

Prunus floribus racemosis, foliis deciduis basi subtribus biglandulosis. Padus. LINN. Sp. 1. Padus HALL. H. 357. Cerasus *racemosa sylvestris*, fructu non eduli. I. R. H. Elzbeeren. Circa *Northeim*, *Noribergam*, & in *Holvetia*. Fructus & calyx similis prioris. Fr.

AMYGDALVS. LINN. 548. Persica. T. 400.

Habitus prioris. Fructus subrotundus, villosus. Nux compressa, sulcis reticulata.

H. *Amygdalus* foliorum ferraturis omnibus acutis. *Perfica*. LINN. *Sp.* 1. *Perfica* Malus. BLACKW. *T.* 101. B. *Pin.* 440. *Pferfigbaum*. Fr.

ORDO. III.

Pomiferae *Hesperides*.

CITRVS. LINN. 807. *Aurantium*. *Citreum*.
Limon. TOVRN. 393-397.

Calyx minimus, quinquedentatus. Stamina multa filamenta inter se connata, ex thalamo orta. Stylus stigmate globoso. Fructus pulposus, novemlocularis.

H. *Citrus* petiolis linearibus. *Medica*. LINN. *Sp.* 1. ROYEN. 266. *Malus Medica*. B. *Pin.* 435. *Citria* Malus. BLACKW. *T.* 361. Fr.

H. β . *Limon vulgaris*. FERRAR. *Hesper.* 193. *Limonia* Malus. BLACKW. 362. Fr.

H. *Citrus* petiolis alatis. *Aurantium*. LINN. *Sp.* 2. *Malus Aurantia*. B. *Pin.* 346. *Aurantia* BLACKW. *T.* 349. Omnes sponte in vtraque *India*. Fr.

CLASSIS. VII.

MULTILOCVLARES.

Similes prioribus numero staminum, petalis liberis, patentibus. Fructus autem est capsula sicca, aut unilocularis, aut multilocularis.

PHILADELPHVS. LINN. Sp. 440. Syringa.

T. 389.

Petala quatuor. Tuba quadrifido stigmate. Capsula quadrilocularis calyce cincta.

1. *Philadelphus foliis subdentatis. coronarius.* LINN. Sp. 1. *Syringa alba.* 1. *Philadelphus Atbei.* I. R. H. B. Pin. 399. *Frutex coronarius.* 18. *Hist.* I. p. 55. Ad sepes & in hortis, spontanea, Fr.

CAPPARIS. LINN. 567.

Calycis folia quatuor, totidemque petala maxima. Capsula carnosâ unilocularis.

1. *Capparis aculeata. spinosa.* LINN. Spec. 1. med. 259. *Capparis spinosa, fructu minore, rotundo.* B. Pin. 480. In Europa Australi. Fr.

TILIA. LINN. 487. HALL: 357.

Calycis folia quinque colorata, totidemque petala. Tuba stigmate pentagono. Capsula quinquelocularis, sed unicum semen plerumque adolescit.

Tilia floribus nectario destitutis. LINN. Sp. 1. a *foliis amplioribus, mollibus subhirsutis.* HALL: v. 357. *Tilia femina, folio maiori.* B. Pin. 426.

1. *Tilia folio glabro, duriori, minori.* HALL: l. n. 2. *Tilia femina folio minori.* BAVH. l. c. 1. s & ad vias. Fr.

CORCHORVS. LINN. 599.

Calycis folia & petala quinque. Tuba crassa. Sinus cylindraceus, quinquelocularis, maxima.

H. *Corchorus* capsulis oblongis ventricosis, foliorum infimis serraturis setaceis reflexis. *olitorius* LINN. *Sp.* 2. *Corchorus* PLINII. B. P. 317. CAMER. *Hort.* 47. T. 12. In *Asia, Africa, America.* A.

CISTVS. LINN. 598.

Calyx quinquefolius, duo folia alterna minora. Petala quinque fugacia. Stilus stigmate orbiculato. Capsula subrotunda.

H. *Cistus* arborescens, foliis lanceolatis supra laevibus, petiolis basi coalitis vaginantibus. *Ladanifera*. LINN. *Sp.* 3. *Cistus ladanifera hispanica incana*. B. *Pin.* 467. *Cistus Ledon* L. *angustifolius*. CLVS. *Hist.* I. p. 77. In *Hispania*. Fr.

H. *Cistus* arborescens, foliis lanceolatis, sessilibus, vtrinque villosis, trinerviis, alis nudis. *Monspeliensis*. LINN. *Sp.* 5. *Cliff.* 205. *Cistus Ledon*. V. CLVS. *Hist.* I. p. 79. *Cistus ladanifera Monspeliensium*. B. *Pin.* 467. Folia sicciora, ac in sequente, margine reflexa, inferne exalbida, vix pilosa, viscida, odore forti, basi vaginantia, nervis lateralibus per maximam folii longitudinem decurrentibus. Petioli & calycis folioli pilis albis dense obsita. In *Gallia Narbonensi* & *Hispania*. Fr.

H. *Cistus* arborescens, foliis sessilibus basi coalitis vtrinque villosis, rugosis, ellipticis. *Cistus Ledon* IV. CLVS II p. 78. nostram plantam exacte refert, etsi *Cistus hirta*. LINN. *Cent.* I. n. 37., quo ille iconem *Chusianam* revocat, a nostra sit diversissima. Num *Cistus* HALL. *Gott.* n. 2. ? Viderur. Folia vtrinque pilosa, sine obtuso, qualia CLVSIVS pingit,

git, molliora ac in priori, viscida, nervis lateralibus
vix ad dimidiam longitudinem folii decurrenti-
bus. Fr.

H. Cistus arborescens foliis sessilibus vtrinque
villosis, rugosis, obverse ovatis, margine undula-
tis. Cistus *mas. folio rotundo hirsutissimo*. C. B. Pin.
465. HALL. Gott. 114. Cistus *mas. folio rotun-
diore*. 1. B. Hist. II. p. 2. Icon, ad quam DV HA-
MEL, *Traité des arbres & arbustes* T. 1. T. 64.
Synonymum 1. B. citat, nostrae plantae egregie
convenit, nisi quod petioli iusto longiores pin-
gantur. Folia basi coalita, cum ramulis villosa, sub-
tus incana, ad dimidium lata ac longa, rugosa, non
trinervia, neque viscida, ut in prioribus. Flos pur-
pureus, solitarius. Fr.

H. Cistus arborescens, foliis petiolatis, cordato-
subrotundis, mollibus, vtrinque tomentosis, crenat-
is. Num Cistus *mas.* III. CLVS. Hist. 1. p. Icon
DV HAMEL. lib. cit. T. 68. sic satis ob figuram
foliorum ad nostram accedit. Arbuscula trunco
firmiori. Folia tactu villi sericei instar pubescentia ad
insertionem petioli hamulis sagittata. Ramuli incani.
Nondum floruit. Cistum totus habitus prodit. Fr.

Tres species hic omisas vide sub fine libri.

Cistus helianthemum L. p. 440.

HYPERICVM. LINN. 808.

Calyx quinquepartitus, quinque petala. Fila-
menta staminum multa, saepe coalita. Capsula sub-
rotunda, tot loculis, quot styli adsunt.

Hypericum floribus trigynis, caule ancipiti, foliis
obtusis pellucido-punctatis. *perforatum*. LINN. Sp.

14. HALL. *H.* 360. n. 1. *Hypericum vulgare*. B. *Pin.* 279. *Hypericum*, BLACKW. *T.* 15. Vbique in agris & sylvis. P.

Hypericum humifusum, foliis glabris perforatis, punctis nigris in margine notatis. HALL. *l. c.* n. 2. *Hypericum floribus trigynis, caulibus ancipitibus prostratis*. LINN. *Sp.* 15. *Hypericum minus supinum*. B. *Pin.* 279. I. R. H. In arenosis v. g. M. Hagen; jenseit Rheinbausen im Wald. P. ^{dam. in junc.} ^{in der Nahe d. Rhyn. Berg.}

Hypericum foliorum ora punctata, calycibus ferrato-glandulosis. HALL. *H.* 361. n. 4. *Gott.* 115. *Hypericum hirsutum*. LINN. *Sp.* 16. *Var. α.* Ascyron. f. *Hypericum bifolium glabrum, non perforatum*. B. *Pin.* 280. *Androsaemum Campoclaurense*. COLUMN. *Ephr.* 73. *T.* 74. Tubae tres. Circa Rheinbausen, in monte ad Bovenden, ad viam sub M. Pless. In M. Altenstollberg. P.

Hypericum villosum, calycibus ferrato-glandulosis, foliis perforatis. HALL. *H.* n. 5. *Gott.* 116. *Hypericum* LINN. *Sp.* 16. *Var. β.* *Hyp. villosum erectum, caule rotundo*. I. R. H. *Androsaemum alterum hirsutum*. COL. *l. c.* p. 75. Rarius ad sepes. P.

Hypericum floribus trigynis, calycibus ferrato-glandulosis, foliis cordatis glabris. *pulchrum*. LINN. *Sp.* 18. *Hypericum calyce ferrato, foliis connatis*. HALL. *Gott.* 116. *Helv.* 3. *Hyp. pulchrum* Tragi. I. R. III. p. 383. In sylva super Hardeggen. P.

Hypericum caule quadrangulo, foliis perforatis. HALL. *H.* n. 6. *Gott.* 116. *Hyp. quadrangulum*. LINN. *Sp.* 13. I. R. H. In fossis. P.

H. *Hypericum floribus trigynis, staminibus corolla longioribus, caule ancipiti fruticoso. hircinum.* LINN. *Sp.* 8. *Hypericum foetidum frutescens.* 1. R. H. DILL. *Elth.* 182. T. 151. In *Europa australi.* Fr.

H. *Hypericum floribus trigynis, fructu baccato, caule fruticoso ancipiti. Androsæum.* LINN. *Sp.* 5. *Androsæum maximum frutescens.* B. Pin. 280. *Hypericum bacciferum.* BLACKW. T. 94. In *Anglia Italia.* Fr.

H. *Hypericum floribus pentagynis, caule fruticoso, foliis ramisque cicatritatis. Bulearicum.* LINN. *Sp.* 1. *Myrto-Cistus Pennæi.* CLVS. *Hist.* I. p. 68. In *Maiorca.* Fr.

CLASSIS. VIII.

COLVMNIFERAE.

Calyx saepe duplex, interior quinquesidus. Petala quinque obcordata, vnguibus connata cum tubo, ex filamentis staminum coalitis composito. Stylus per tubum stamineum adscendens fissus in tot stigmata, quot loculi in fructu circa axin. Omnibus vires emollientes.

HIBISCVS. LINN. 756. *Ketmia.* T. 26.

Calyx exterior multifidus; interior quinquesidus. Stigmata quinque crassa. Fructus loculi quinque, polyspermi.

H. *Hibiscus foliis ovato-lanceolatis; superne inciso-serratis, caule arboreo. Syriacus.* LINN. *Sp.* 9.

Alcea arborescens. CAMER. H. T. 3. 4. *Syriaca*. B. Pin. 316. In Syria. Fr.

H. *Hibiscus aculeatus*, foliis trilobis & quinquelobis, inaequaliter ferratis. HALL. Gott. 117. *Hibiscus*. LINN. Sp. 13. Var. δ . *Ketmia Indica*, foliis digitatis, flore magno sulphureo, umbone atropurpureo, petiolis spinosis. EHRET, T. 6. f. 1. *Alcea Benghalensis*. Comm. H. I. p. 35. T. 18. In India. A.

H. *Hibiscus foliis palmato-digitatis septempartitis*, *Manibot*. LINN. Sp. 13. *Ketmia folio Manibot ferrato*, flore amplo sulphureo. DILL. Elth. 189. T. 156. f. 189. In India. A.

H. *Hibiscus foliis palmatis ferratis*, fructu pyramidalis, longissimo. HALL. G. 118. An *Ketmia Zeylanica Fici folio*, perianthio oblongo integro. DILL. Elth. T. 157. f. 190. ? *Hibiscus ficulneus*. LINN. Sp. 10. ? Folia conveniunt, sed in nostra flores non nutant. An potius huc referenda *Alcea Americana* flore flavo, fructu pyramidalis sulcato. COMM. H. Amst. 1. p. 537. T. 19. *Hibiscus foliis quinquepartito-pedatis*, calycibus interioribus latere rumpentibus. LINN. Sp. 15. ? Caulis undique asper aculeis sparsis mitibus. Folia inferiora triloba, superiora quinqueloba, lobis ferratis latioribus. Flos prior multo minor simili colore. Calyx interior spathaeformis brevissime quinquefendendus. Fructus pyramidalis undique strigosus & aculeis echinatus. A.

H. *Hibiscus foliis subpeltato-cordatis septemangularibus ferratis hispidis*. *Abelmoschus*. LINN. Sp. 14. *Ketmia Aegyptiaca*, semine moscato. I. R. H.

100. BVRM. *Cycl.* 134. *Abelmosch vulgo.* HERM. *Lugdb.* 25. In *Indiis.* P.

H. Hibiscus foliis imis cordato-lanceolatis, superioribus pentagoniis ferratis. HALL. *Gott.* 118. Hibiscus foliis inferioribus cordatis angulatis, superioribus subbistatis, floribus subnutantibus, pistilla cernuo. (*Pentacarpus.*) LINN. *Sp.* 19. cum descr. *Ketmia palustris minor.* ZANICH. *Venet.* 155. T. 91. In paludosis *Venetiae.* Corolla purpurea. P.

H. Hibiscus foliis tripartitis incis, calycibus inflatis. *Trionum.* LINN. *Spec.* 20. *Aloea vesicaria.* B. *Pin.* 317. β. *Ketmia vesicaria Africana.* TOURN. J. R. H. 101. In *Italia, Africa.* A.

ALTHAEA. HALL. LINN. 749.

Alcea. LINN. 750.

Calyx duplex, exterior in Althaea inaequaliter novemfidus, in Alcea semisextidus. Capsulae monospermae, fructum depressum faciunt.

H. Althaea foliis simplicibus tomentosis. *officinalis.* LINN. *Sp.* 1. Althaea *Dioscoridis & Plinij.* B. *Pin.* 315. Althaea BLACKW. T. 90. In subhumidis. v. g. ad fluvium *Vnstrut.* P.

H. Althaea foliis scabris, acutis, semitrilobis. HALL. *Gott.* 119. Althaea foliis inferioribus palmatis, superioribus digitatis, caule fruticoso. (*cannabin.*) LINN. *Sp.* 2. B. *Pin.* 316. In *Hungaria, Italia.* P.

H. Althaea foliis sinuato angulosis. HALL. l. c. (*Alcea rosea.*) LINN. *Spec.* 1. *Malva rosea folio*

subrotundo. B. Pin. 315. *bortensis*. DOD. *Pempt.* 652. floribus varii coloris. In *Hungaria*. *Palæstina*. B.

H. *Althaea* foliis palmatis. HALL. l. c. (*Alcea*) *ficifolia*. LINN. Sp. 2. *Malva rosea bortensis*. *folio ficus*. B. l. c. I. R. H. floribus varii coloris. B.

MALVA. HALL. LINN. 751.

Lavatera. LINN. 752.

Calyx in *Lavateris* semitrifidus, in *Malvis* trifolius. Fructus prioris.

H. *Malva* caule arboreo, foliis palmatis, obtusis, plicatis. HALL. *Goit.* 120. *Lavatera caule arboreo*, foliis septemangularibus tomentosis plicatis, pedunculis confertis axillaribus ^{arbores}. LINN. Sp. 1. *Malva arborea Veneta dicta*, parvo flore. B. Pin. 315. Ad ripas Maris mediterranei. Fr.

H. *Malva* (*Lavatera*) foliis quinqueangularibus, acutis crenatis, lacinia media productiore. LINN. *Cliff.* 348. HALL. G. 120. *Lavatera caule fruticoso*, foliis quinquelobo-hastatis. (*Olbia*.) LINN. Sp. 3. *Althaea frutescens*, folio acuto, molli, virente, flore specioso. PLUKNET. T. 8. f. 1. *Althaea arborea olbia* in Gallo Provincia. LOB. ic. 653. Pedunculi axillares solitarii brevissimi. Ad ripas maris Mediterranei circa *Monspelim*. Fr.

H. *Malva* caule arboreo, foliis tomentosis, obtusissimis micantibus. HALL. l. c. *Lavatera micans*. LINN. Spec. 2. *Malva foliis mollibus undulatis, superius micis sulphureis ad solem splendentibus donatis*. MORIS. *Hist.* 1. p. 523. S. 5. T. VI. f. 9. *Folia*

in tomentosa, subrotunda, plicata. Stipulae magnae ovatae. Pedunculi axillares solitarii, petiolis breviores. Calyx exterior trifolius. Habitus prioris. In *Hispania*. Fr.

H. *Malva* (*Lavatera*) caule fruticoso, foliis subcordatis subtrilobis, rotundatis crenatis, stipulis cordatis, pedunculis unisporis. *triloba*. LINN. *Sp.* 4. *Althaea frutescens, folia rotundiora incano*. B. *Pin.* 316. Pedunculi tres, pluresve ex alis foliorum, petiolis breviores. Stipulae cordatae. Calyx exterior trifolius. In *Lusitania*. Fr.

H. *Malva* (*Lavatera*) foliis inferioribus cordato-subrotundis, superioribus lobatis crenatis, calycibus utrinque incis, caule herbaceo. ROY. *Lugdb.* 355. *Lavatera Thuringiaca*. LINN. *Sp.* 6. *Althaea Thuringiaca grandiflora*. DILL. *Elth.* 9. T. 8. f. 8. CAMER. *Hort.* 12. T. 6. In *Thuringia, Pannonia*.

H. *Malva* foliis subrotundis, capsulis scuto subpositis. HALL. l. c. *Lavatera foliis glabris, caule scabro herbaceo, pedunculis unisporis, fructibus orbiculo cectis. (trimestris.)* LINN. *Sp.* 8. *Lavatera foliis subcordatis obtusis angulatis serratis, caule herbaceo*. ROY. 355. *Malva folio vario*. B. *Prodr.* T. 137. flore rubro & albo. In *Hispania, Gallia Narbonensi*. A.

H. *Malva* caule erecto, foliis semiorbiculatis crenatis, calyce exteriori diphylo. *Hispanica*. LINN. *Sp.* 11. *Malva rotundifolia glabra ex Hispania, flore amplo*. PLVKNET. *Alm.* 238. T. 44. f. 3. flore rubro & albo. In *Hispania*. A.

Malva caule erecto, foliis pene palmatis obtuse ferratis. HALL. *Helv.* 362. Malva caule erecto herbaceo, foliis lobatis obtusis, pedunculis petiolisque pilosis. (*sylvestris*.) LINN. *Sp.* 9. *M. sylvestris*, folio sinuato. B. *Pin.* 314. vulgaris. BLACKW. *T.* 22. Vbique in gramineis. A.

Malva caule repente, foliis cordato-orbiculatis obsolete quinquelobis. *rotundifolia*. LINN. *Sp.* 8. HALL. *H.* 363. Malva *sylvestris* folio rotundo. B. *Pin.* 314. Vbique ad vias. A.

h. Malva floribus albis, parvis, conglomeratis. HALL. *Gott.* 121. Malva caule erecto, foliis angulatis floribus axillaribus glomeratis. (*verticillata*) LINN. *Sp.* 12.

Malva *lin.* *p. ed. 2. p. 110*
β. Malva foliis crispis. C. B. 315. MORIS. *S. V.* *T.* 15. In Syria. A.

γ. Malva humifusa, minima SHERARDI. TILLII *Hort. Pis.* *T.* 35.

η. Malva foliis subcordatis laciniatis glabris, caule arborecente. *Capensis*. LINN. *Sp.* 6. Malva *capensis frutescens grossulariae folio maiori hirsuto*. DILL. *Elth.* 209. *T.* 169. f. 207. In Africa. Fr.

Malva caule erecto, foliis multipartitis scabriuiculis. *Alcea*. LINN. *Spec.* 13. HALL. *H.* 363. η. 3. *Alcea vulgaris maior*. C. B. 315. I. R. H. BLACKW. *T.* 309. Ad viam versus *Bovenden* in dumetis, ad rivum prope *Langlern*. P.

θ. Malva foliis palmatis scabris, caule tomentoso fruticoso, pedunculis multifloris. *bryonifolia*. LINN.

LINN. *Sp.* 5. *Althaea frutescens, bryoniae folio.* B. *Prodr.* 138. In *Hispania*. Fr.

H. Malva caule erecto, foliis imis subrotundis, superioribus tenuiter divisis. HALL. *H.* 364. Malva foliis radicalibus reniformibus incis, caulinis quinquepartitis pinnato-multifidis, ~~moscata~~. LINN. *Sp.* 14. Malva montana, f. *Alcea rotundifolia laciniata*. COL. *Ecpbr.* 1. p. 148. T. 147. In *Alsacia, Helvetia, Italia*. P.

H. Malva foliis palmatis dentatis, corollis calyci aequalibus. An Malva aegyptia. LINN. *Sp.* 15. c. *descr.*? Pedunculi saepe biflori in nostra semper foliis longiores, ex alis solitarii, in summitate plures congesti. Corollae coerulescentes, calyci aequales. *Alceam minorem maritimam tenuifoliam procumbentem* HERM. *Parad.* 9. T. 9. quae est Malva Tournefortiana. LINN. *Cent.* 1. n. 63. huc referrem ob figuram foliorum, nisi corollae calyce nimis maiores pingerentur. A.

H. *Malva Peruviana* Linn. 8 spec. 6.

SIDA, LINN, 747. Abutilon.

TOVRN. 25.

Calyx ut plurimum simplex. Fructus capsulis cornutis, cristatis.

H. *Sida* caule repente, foliis multifidis. Malva *Caroliniana* LINN. *Sp.* 7. Abutilon *Carolinianum repens, alceae foliis, flore belvolo*. DILL. *Elth.* 5. T. 4. f. 4. MARTYN. *Cent.* T. 34. Calyx exterior trifolius. Capsulae aristatae. In *Carolina*. A.

H. *Sida* foliis cordatis subrotundis crenatis tomentosis, pericarpis inflatis multilocularibus crehatis

tis repandis. *crispa*. LINN. *Sp.* 8. *Abutilon vesca-*
rium crispum, floribus melinis parvis. DILL. *Elth.*
6. T. 5. f. 2. In Carolina, Babama.

H. Sida foliis cordato-lanceolatis integerrimis,
periplocifolia. LINN. *Sp.* 5. *Abutilon periplocae*
acutioris folio, fructu stellato. DILL. *Elth.* 4.
T. 3. f. 3. In Ceylona.

* H. Sida foliis inferioribus cordato-angulatis, su-
perioribus hastatis, petalis integerrimis. ROYEN.
Lugdb. 349. LINN. *Vps.* 199. *Abutilon america-*
num, flore coeruleo. HALL. *Gott.* 122. Caules de-
biles procumbentes. Anguli foliorum inprimis su-
periorum acuti, ora fusca, rarius dentata. Stigma-
ta septem. A.

H. Sida foliis crenatis, inferioribus dentatis ob-
tusis, superioribus acuminatis. ROYEN. 349. Sida
cristata. LINN. *Spec.* 9. *Abutilon Lavaterae flore*,
fructu cristato. DILL. *Elth.* 3. T. 2. f. 2. Caules
erecti. Folia omnia crenata, inferiora obtusiora,
subcordata, superiora saepius hastata, angulis poste-
rioribus rotundatis. Flos priori multo maior. Sti-
mata viginti. Planta a priori diversissima. A.

H. Sida foliis subrotundo-cordatis acuminatis
crenatis. *Abutilon*. LINN. *Sp.* 7. *Althaea Theophras-*
ti, flore luteo. B. *Pin.* 316. In Indiis. A.

H. Sida foliis ovatis ferratis, floribus ad alas con-
glomeratis. *Abutilon Vlmifolium*, floribus congloba-
tis ad foliorum alas. HALL. *Gott.* 123.

H. Sida foliis lanceolato-rhomboidibus ferratis
rhombifolia. LINN. *Sp.* 2. *Malvinda unicornis*, fo-
lio

*la rhomboid. perenn. TALL. Fl. 216. T. 102.
f. 12. Arch. carn. T. 2. f. 10. f. 10. f. 10. f. 10. f. 10.
St. 11. T. 6. f. 2. Pedunculi petioli longiores. A.*

*H. Sida foliis cordato-ovatis serratis, stipulis serratis axillis truncatis. (Sida) LINN. Sp. 1. Mal-
vacea bicolor, sanguinea. TALL. Fl. 216. f. 10. f. 10. f. 10. f. 10. f. 10.
DILL. Herb. L. 12. T. 1. f. 1. f. 1. f. 1. f. 1. f. 1. Pedunculi pe-
toli sere breviores. A.*

H. Sida foliis inferioribus subrotundis, superioribus, tribus, crenatis, glandulis exiguis aggregatis, folio spathuliformi exceptis. Ita interim definire licet plantam, cum flores plures in involucrio spathulaceo compresso aggregati latent. Corolla flava calyce piliis paulo maior. Ad utrumque latus singulorum florum adstant duae ligulae lineares, capillaceae. Stamina pauci. Capsulae quinque monospermae, seminibus oblongis, masculinis. Verius hanc plantam alibi proxime describam. (Sida) LINN. Sp. 1. f. 1. f. 1. f. 1. f. 1. f. 1.

VRENA. LINN. 754.

Calyx exterior semiquinquedus, interior quinquefolius. Capsulae echinatae.

*H. Vrena foliis angulatis. LINN. Sp. 1. Vre-
na Sinica, Xanthii facie. DILL. Herb. 340. T. 319.
f. 412. In China. Fr.*

NAPAEA. LINN. 748.

Calyx simplex. teres. Petala oblonga. Stamina plurima. Capsulae muticae.

*H. Napaea pedunculis involucriatis angulatis, foliis scabris, floribus dioicis. dioica, LINN. Sp. 1. Abu-
tilon*

tilon folio profunde dissecto, pedunculis multifloris, femina & mas. EHRET. T. 7. f. 1. & T. 8. HALL. Gott. 124. Flores feminae staminibus effoetis, abortivis gaudent. In Virginia. P.

HERMANNIA. LINN. 742.

Calyx simplex. Stamina quinque. Stigma simplex. Fructus quinquelocularis.

H. Hermannia foliis lanceolatis obtusis, ferratis. Hyssopifolia, LINN. Sp. 4. Althaea Africana frutescens, floribus aureis cochleatis pendulis minoribus. VOLKAM. Norib. T. 24. In Africa. Fr.

PENTAPETES. LINN. 757.

Calyx simplex. Stamina decem fertilia, binatim alterne posita cum filamentis quinque sterilibus, coloratis.

H. Pentapetes foliis hastato-lanceolatis ferratis. phoenicea. LINN. Sp. 1. Alcea Indica, lucido hastato folio, flore blattariae phoeniceo. PLUK. Phyt. T. 255. f. 3. A.

GOSSYPIMUM. LINN. 755.

Calyx duplex. Capsula quadrilocularis. Semina lana involuta.

H. Gossypium foliis palmatis, lobis lanceolatis, caule fruticoso. arborescens. LINN. Sp. 3. Gossypium arborescens, caule larvi. BAYH. Pin. 430. In India. Fr.

CLASSIS IX.

SVCCULENTAE.

caulis & folia succo aqueo turgida & carnosâ.

ORDO. I.

CACTI.

Calyx concavus monophyllus. Corolla in pluripetala profundissime fissa, parieti calycis affixa. Nina plurima calyci inserta. Fructus carnosus subcô.

CACTVS. LINN. 539. Cereus. Opuntia.

Melocactus. Tuna. Peireskia.

Planta aphyllor. Calyx tubulato-cavus. Stylus odraceus stigmate multifido. Fructus carnosus, ocularis. Sola Peireskia foliis instructa.

1. Cactus subrotundus testis tuberculis, ovatis, natis. *mammillaris*. LINN. Sp. 1. Echino-Melocactus minor lactescens, tuberculis sc. *mammillis* natis. HERM. Parad. 136. T. 136. Ficoides Americana sphaerica, tuberculata lactescens, flore al. COMM. H. 1. p. 105. T. 55.

2. Cactus quadrangularis longus erectus: angustis compressis. *tetragonus*. LINN. Sp. 4. H. Vps.

3. Cactus erectus sexangularis longus. *hexagonus*. LINN. Sp. 5. Cereus altissimus Surinamensis. RM. Par. 116.

4. Cactus erectus longus subnovemangularis; angustis obsoletis, spinis latis brevioribus. *laevigatus*. LINN.

LINN. Sp. 8. An *Cereus Curassavicus erectus, lanugine flavescente*. HERM. Parad. T. 115.?

H. Cactus repens subquinqueangularis. *grandiflorus*. LINN. Sp. 11. Cactus *scandens angulis quinque pluribusque obtusis*. H. Cliff. 182. c. descr. *Cereus Americ. maior articulatus, flore maximo, nocte se aperiente*. VOLK. Hesp. 1. T. 234. TREWII Icon. plant. rar.

H. Cactus repens decemangularis. *flagelliformis*. LINN. Sp. 12. Cactus *scandens angulis decem obsoletis*. H. Vpf. 121. *Cereus minor scandens polygonus spinosissimus, flore purpureo*. TREW. rar. pl. Dec.

1. Cactus triangularis scandens articulatus. ^{*triangularis*} LINN. Sp. 13. *Cereus erectus, foliis triangularibus profunde canaliculatis*. PLUK. Alm. 147. T. 29. f. 3.

H. Cactus compressus, articulatus, ramosissimus, articulis ovatis, spinis setaceis. LINN. Cliff. 183. Cactus *Opuntia*. LINN. Sp. 15. *Opuntia vulgo barbariorum*. 1. B. H. 1. p. 154. In *America calidiori, & Virginia*.

H. Cactus compressus articulatus, ramosus, articulis ovato-oblongis, spinis setaceis. LINN. Cliff. 183. Cactus *Ficus indica*. LINN. Sp. 16.

H. Cactus compressus articulatus ramosus, articulis ovato-oblongis, spinis subulatis. LINN. Cliff. 183. Cactus *Tuna*. LINN. Sp. 17. *Tuna maior spinis validis flavicantibus, flore gilvo*. DILL. Elth. 396. T. 295. f. 238.

H. Cactus tereti-compressus articulatus ramosus. LINN. Cliff. 182. Cactus *Curassavicus*. LINN. Sp.

19. *Ficus indica* f. *Opuntia Curassavica minima*.
COMM. H. 1. p. 107. T. 56.

H. *Cactus* prolifer ensiformi-compressus serrato-
repandus. *Phyllanthus*. LINN. Sp. 20. *Cactus foliis*
anisformibus obtuse serratis. H. Cliff. l. c. *Cereus colo-*
pandrii folio brachiato. DILL. *Elth.* 73. T. 64. f. 74.

H. *Cactus* caule tereti arboreo, spinoso, foliis
lanceolato-ovatis. *Peirekia*. LINN. Sp. 21. *Peire-*
kia aculeata, flore albo, fructu flavescente. DILL.
Elth. 305. T. 227. f. 294.

MESEMBRYANTHEMUM. LINN. 552. DILL.

Elth. p. 159. Ficoides TOURN.

Calyx semiquinquefidus carnosus. Fructus carno-
sus tot loculis, quot styli adfunt. Folia in plerisque
opposita.

H. *Mesembryanthemum* foliis alternis ovatis obtu-
sis undulatis. *crystallinum*. LINN. Sp. 2. *Mesembryan-*
themum crystallinum, plantaginis folio undulato (mi-
sis argenteis asperso). DILL. *Elth.* 231. T. 180. f. 211.

H. *Mesembryanthemum* acaule, foliis linguiformi-
bus, altero margine crassioribus. *linguiforme*. LINN.
Sp. 34. *Mesembr. folio linguiformi latiore*. DILL.
Elth. T. 184. f. 224. Corolla flava.

H. *Mesembryanthemum* foliis acinaciformibus
connatis, angulo carinali scabris, ramis angulatis,
Mesembr. acinaciforme, flore amplissimo purpureo.
DILL. *Elth.* T. 211. f. 271.

β. *Mesembryanthemum falcatum maius, flore*
amplio luteo. DILL. T. 212. f. 271.

H. Mesembryanthemum foliis acinaciformibus distinctis laevibus, ramis teretibus. ^{falcatum} LINN. Sp. 20. Mesembr. *falcatum*, *minimum*, *flore purpureo parvo*. DILL. Elth. T. 213. f. 276.

H. Mesembryanthemum acaule, foliis semicylindricis connatis externe tuberculatis. *rostratum*. LINN. Sp. 26. Mesembr. *rostrum ardeae referens*. DILL. Elth. T. 186. f. 229.

H. Mesembryanthemum caulescens, foliis subulatis semicylindricis recurvis connatis longis. *lorcum*. LINN. Sp. 29. Mesembr. *foliis corniculatis longioribus*. DILL. Elth. T. 199. f. 253.

H. Mesembryanthemum foliis alternis subulatis triquetris longissimis. LINN. Sp. 35. Mesembr. *folio pugioniformi, flore amplo stramineo*. DILL. Elth. T. 210. f. 269.

H. Mesembryanthemum foliis subcylindricis acutis connatis arcuatis laevibus. *veruculatum*. LINN. Sp. 27. Mesembr. *foliis veruculiformibus, floribus melinis umbellatis*. DILL. Elth. T. 203. f. 259.

H. Mesembryanthemum acaule, foliis subteretibus connatis, floribus octogynis. LINN. Sp. 6. Mesembr. *calamiforme*. DILL. Elth. T. 186. f. 228.

H. Mesembryanthemum ^{Quila Cris. f. 32} foliis dolabriformibus. LINN. Sp. 32. DILL. Elth. T. 191. f. 237. Corolla intus aurea micans, extus aurantia.

H. Mesembryanthemum foliis subulatis triquetris strictis acutis: punctis pellucidis obsoletis sparsis. *glaucum*. LINN. Sp. 28. Mesembr. *scabrum, flore sulphureo convexo*. DILL. Elth. T. 196. f. 248.

H. Me-

γ. *Mesembryanthemum* acaule, foliis triquetris linearibus apice trifariam dentatis. LINN. *Sp.* 18. *Mesembr. bellidiflorum*. DILL. *Elth. T.* 189. f. 233.

η. *Mesembryanthemum* caulescens, foliis deltoidibus triquetris dentatis. LINN. *Sp.* 9. *Mesembr. deltoides*. & dorso & lateribus muricatis. DILL. *Elth. T.* 195. f. 246.

β. *Mesembryanthemum* deltoides, foliis non dorso, sed lateribus muricatis. DILL. *Elth. T.* 195. f. 243.

η. *Mesembryanthemum* foliis linearibus obsolete triquetris distinctis: summis imbricatis laevibus, *splendens*. LINN. *Sp.* 25. *Mesembr. foliis confertis splendidibus, flore pallido*. DILL. *Elth. T.* 204. f. 260.

η. *Mesembryanthemum* foliis planis oppositis ovatis acuminatis connatis integerrimis. *tortuosum*. LINN. *Sp.* 30. *Mesembr. tortuosum, foliis semper-vivi expansis*. DILL. *Elth. T.* 182. f. 223. β. Simile prioris sed foliis congestis. *T.* 181. f. 222.

η. *Mesembryanthemum* articulis caulinis terminatis in folia acuminata subtus dentata. *vincinatum*. LINN. *Sp.* 14. *Mesembr. perfoliatum, foliis maioribus triacanthis*. DILL. *Elth. T.* 193. f. 240.

η. *Mesembryanthemum* foliis semicylindraceis, calycibus quadrifidis. *noctiflorum*. LINN. *Sp.* 4. *Mesembr. noctiflorum, flore intus candido, extus stramineo odoratissimo*. DILL. *Elth. T.* 206. f. 263. Styli quatuor.

H. Mesembryanthemum foliis apice barbatis. barbatum. LINN. Sp. 10. Mesembr. radiatum, humile, floribus minoribus. DILL. Elth. T. 190. f. 235.

β. Mesembryanthemum radiatum, ramulis prolixis, decumbentibus. DILL. Elth. T. 190. f. 234.

γ. Mesembryanthemum radiatum, humile, foliis maioribus. DILL. Elth. T. 190. f. 236.

H. Mesembryanthemum foliis semiteretibus, floribus sessilibus axillaribus. geniculiflorum. LINN. Sp.

3. Mesembr. Capense geniculiflorum, Neapolitanum creditum. DILL. Elth. T. 205. f. 261.

H. Mesembryanthemum foliis subtriquetris scabris, corollis bicoloribus. bicolor. LINN. Sp. 22.

H. Mesembryanthemum tenuifolium, frutescens, flore croceo. DILL. Elth. T. 202. f. 258. Petala intus flava, extus rubella.

H. Mesembryanthemum foliis subulatis semiteretibus glabris internodio longioribus. tenuifolium. LINN. Sp. 17. Mesembr. tenuifolium procumbens, flore coccineo. DILL. Elth. T. 201. f. 256.

AIZOON. LINN. 553. Ficoidea. DILL.

Calyx quinquepartitus. Corolla nulla. Stamina per phalanges calyci inserta.

H. Aizoon foliis cuneiformi-ovatis, floribus sessilibus, Canariense. LINN. Spec. 1. Kali aizoides Canariense, procumbens, portulacae foliis, aspergine rorida perpetuo madidis. VOLK. Norib. T. 236. A.

H. Aizoon foliis lanceolatis, floribus sessilibus.
Hispanicum. LINN. *Sp.* 2. Ficoidea *Hispanica annua*, folio longiore. DILL. *Elth.* T. 117. f. 143. A.

ORDO. II.

Petalis liberis & staminibus receptaculo insertis.

PORTULACA. LINN. 531.

Calyx bifolius. Petala quinque. Germen sub flore. Stigmata quinque. Capsula ovata, unilocularis.

H. *Portulaca* foliis cuneiformibus, floribus sessilibus. *oleracea*. LINN. *Spec.* 1. HALL. *Helv.* 392. *Portulaca sylvestris*. LOB. *ic.* 388. In *Europa australi*. v. g. ad ripam *Rheni*; in utraque *India*. A.

H. *Portulaca* foliis subulatis alternis, axillis pilosis, floribus sessilibus. *pilosa*. LINN. *Sp.* 2. *Portulaca Curassavica*, lanuginosa procumbens. HERM. *Parad.* T. 215. COMM. H. I. p. 2. T. 5. In *America meridionali*.

TETRAGONIA. LINN. 551.

Calyx quadrifolius, coloratus. Corolla nulla. Styli quatuor. Capsula quadrangularis sub flore.

H. *Tetragonia* foliis linearibus. *fruticosa*. LINN. *Sp.* 1. *Tetragonocarpus Afr. rutilans*, foliis longis & angustis. COMM. H. II. p. 205. T. 103. Fr.

CRASSULA. LINN. 352. DILL. Elth. 96.

Calyx quinquefolius. Petala quinque unguibus connata. Stamina quinque, totidemque capsulae, ad basin nectariferae.

H. *Crassula* foliis planis cartilagineo-ciliatis basi connato-vaginantibus. *coccinea*. LINN. Sp. 1. *Cotyledon Afr. frutescens, flore umbellato coccinea*. COMM. rar. 24. T. 24. Fr.

H. *Crassula* foliis lanceolato-subulatis sessilibus connatis, canaliculatis subtus convexis. *perfoliata*. LINN. Sp. 2. *Crassula altissima, perfoliata*. DILL. Elth. T. 96. f. 113. *Aloe Afr. caulescens perfoliata glauca & non spinosa*. COMMEL. Prael. 74. T. 23. Fr.

H. *Crassula* foliis subulatis obsolete tetragonis. *tetragona*. LINN. Sp. 3. ROY. Lugdb. 455. Fr.

H. *Crassula* foliis oppositis obtuse ovatis integerrimis: hinc angustioribus. *cultrata*. LINN. Sp. 4. *Crassula Anacampserotis folio*. DILL. Elth. T. 97. f. 114. Fr.

H. *Crassula* Portulacae facie, arborescens. DILL. Elth. T. 101. f. 120. *Portulaca arborescens* vulgo. Genus satis certe nondum definiri potest, cum flores a nullo Botanico observati fuerint.

ORDO III.

SVCCULENTAE *dyplostemones*.

Dyplostemones dicuntur, quibus stamina duplicatione ad petala aut segmenta petali sunt; ob summam autem analogiam variae plantae *isostemones* huc vocari debent.

COTYLEDON. LINN. 512. DILL.

Elth. 95.

Calyx & Corolla femiquinquefida. Capsulae quinque, singulis ad basin squama nectarifera.

H. ^{Cotyledon linearis} Cotyledon foliis lineari-lanceolatis, integerrimis, floribus pendulis. Cotyledon Afr. frutescens, folio longo & angusto, flore flavescente. COMMEL. var. T. 23. β. Cotyledon foliis depressis, cruciatis. WALTH. H. p. 16. c. ic.

H. Cotyledon foliis semiglobosis. baemispbaetica, LINN. Sp. 2. DILL. Elth. 112. T. 95. f. 111. Fr.

H. Cotyledon foliis laciniatis, floribus quadrifidis, laciniata. LINN. Spec. 6. Cotyledon Afrq, folio grasso lato laciniato, flosculo aureo. BOERH. Lugdb. 1. T. 288. Fr.

H. ^{Cotyledon orbiculata} Cotyledon orbiculata Sp. pl. p. 614.

SEDVM, HALL. Helv. 392. LINN. 513.

Sempervivum. LINN. 538. Rhodiola.

LINN. 997.

Calycis folia quinque ad duodecim, totidemque petala. Stamina semper bis tot. Capsulae tot quot petala. In Sedis LINN. & Rhodiola squamae nectariferae ad basin capsularum.

H. Sedum caule ramoso, rofulis foliorum ciliatis obrito. HALL. Gutt. 133. Sempervivum caule arborescente laevi ramoso. LINN. Sp. 1. Sedum maius arborescens. B. Pin. 282. legitimum. CLVS. Hist. II. p. 58. In America & Europa australiori. Fr.

Sedum rofulis sessilibus liberis conniventibus, foliis ciliatis, in apice purpureis. HALL. H. 392. n. 1.

L 5

Sem-

Sempervivum foliis ciliatis, propaginibus patentibus. (teetorum.) LINN. Sp. 3. Sedum maius vulgare. B. Pin. 283. BLACKW. T. 366. In rupibus prope Rheinbausen & teetis: P.

H. Sedum rosulis liberis, foliis ciliatis, conniventibus, viridibus. HALL. Gott. 137. En. Helv. n.

2. Sempervivum foliis integerrimis, propaginibus patulis. (montanum.) LINN. Sp. 6. Sedum maius mont. flore rubenta. G. B. Pin. 284. In alpinis Helveticis. P.

H. Sedum rosulis foliorum tomento conjunctis. HALL. H. 393. Sempervivum foliis pilis intertextis, propaginibus globosis. (arachnoideum.) LINN. Sp. 5. Sedum mont. tomentosum, BAVH. Pin. 284. In alpinis Helveticis.

H. Sedum sexu distinctum, tetrapetalum, foliis ferratis. HALL. H. n. 4. Rhodiola. LINN. 1035. Radix Rhodia. B. Pin. 286. CLVS. Hist. II. p. 65. In alpinis. P.

Sedum foliis confertis ferratis ellipticis, umbellæ caulem terminante densissima. HALL. H. n. 5. Sedum Telephium. LINN. Sp. 2. Anacampteros vulgo faba crassa. I. B. III. p. 681. I. R. H. In muris, & agris, v. g. versus Lenglern. A.

• Sedum foliis ovalibus ferratis coniugatis, ex alis foliorum umbelliferum. HALL. H. n. 6. Telephium Hispanicum. I. CLVS. Hist. II. p. 66. Circa Rheinbausen. A. varietas Telephium maximum.

H. Sedum foliis cuneiformibus integerrimis, caulibus decumbentibus, floribus corymbosis. LINN.

Sp.

Sp. 3. *Telephium minus, sempervirens.* LOB, ic, 340. P.

H. *Sedum foliis ovato-lanceolatis, ferratis, floribus flavis umbellatis, caule erecto.* HALL. l. c. *Sedum aizoon.* LINN. Sp. 4. *Anacampseros flore flavo.* AMM. Ruth. 96. T. 11. In *Sibiria*,

Sedum foliis planiusculis angulatis, floribus lateralibus sessilibus solitariis. ^{*Helveticum*} ~~*apertatum*~~. LINN. Sp. 6, HALL. H. n. 7. *Sedum echinatum vel stellatum, flore albo.* 1. B. Hist. 3. p. 680. In *Helvetia, Italia*,

Sedum foliis teretibus, in spinulam terminatis, umbella sparsa. HALL. H. n. 15. *Sedum rupestre.* LINN. Sp. 9. *Sedum rupestre repens, foliis compressis.* DILL. Elth. T. 256. f. 333. *Sedum minus luteum, folio acuto.* B. Pin. 283. Circa *Blankenburg & Kelbra.* P.

Sedum acre, foliis conicis confertissimis, floribus in summis alis sessilibus, luteis. HALL. H. n. 13, LINN. ^{*acre*} Sp. 12. BLACKW. T. 232. *Sempervivum minus vermiculatum acre.* B. Pin. 283. In agris sterilibus & muris, P.

Sedum foliis cylindricis obtusis, sessilibus, patentibus, panicula ramosa. HALL. H. n. 18, LINN. ^{*album*} Sp. 10. *Sedum minus teretifolium album.* B. Pin. 283. *Sedum minus* L. CLVS. p. LIX. LIX. In muris & rupibus. v. g. circa *Rheinhausen*,

SAXIFRAGA. LINN. 494. Geum. TOURN.

Calyx quinquepartitus. Petala quinque. Capsula unilocularis birostris.

H. *Saxifraga* ora foliorum cartilaginea, acuta crenata. HALL. H. n. 8. *Saxifraga foliis radicalibus aggregatis lingularis cartilagineo-ferratis, caule paniculato.* (*Cotyledon.*) LINN. Sp. 1. *Saxifraga sedifolia* foliis subrotundis ferratis. I. R. H.

β. *Saxifraga Sedi foliis angustioribus ferratis.* I. R. H. SEGV. *Vexon. T. IX. f. 1.* In alpinis. P.

H. *Saxifraga foliis petiolatis, retusis, ora cartilaginea, ferrata, caule nudo paniculato.* HALL. H. n. 4. LINN. Spec. 11. *Geum rotundifolium minus.* I. R. H. In Alpibus Helvetiae. P.

H. *Saxifraga foliis reniformibus, acute crenatis, caule ramoso, folioso.* HALL. H. n. 5. *rotundifolia.* LINN. Sp. 19. *Sanicula alpina f. guttata quibusdam.* CAMER. Epit. p. 764. *Geum rotundifolium maius.* I. R. H. In montosis Helvetiae. P.

Saxifraga foliis caulinis reniformibus, lobatis, caule ramoso, radice granulata. *granulata.* LINN. Sp. 20. HALL. H. n. 6. *Saxifraga rotundifolia alba.* B. Pin. 309. BLACKW. T. 56. In collibus apricis, siccis. P.

Saxifraga rosulis sessilibus, foliis palmatis, hirsutis. HALL. Gott. p. 136. *Helv. n. 16. tridactylites.* LINN. Sp. 25. β. *Saxifraga alba petraea* PONAÉ M. Bald. apud GLVSIVM. CCCXXXVII. In Hercyniae rupibus *Christinen-Klippe* dictis prope *Elbingerode.* P. *Saxifraga petraea* Lin. 2. Sp. 25.

Saxifraga foliis late petiolatis, trifidis, ad caulem palmatis. HALL. H. n. 15. LINN. Sp. 25. α. *Saxifraga verna annua humilior.* I. R. H. In muris. A. DI-

DYPLOSTEMONES MVLTIOCVLARES. 173

DYPLOSTEMONES MVLTIOCVLARES.

In plurimis calyx quinquefolius, stamina bis tot, quot petala, styli aut stigmata tot, quot loculi, aut stylus stigmate simplici. Fructu quadri-aut quinquelocularis.

OXALIS. LINN. 515. Oxys. ТОВАН.

Petala vnguibus coälita. Styli quinque. Fructus pentagonus.

Oxalis scapo vñifloro, foliis ternatis, radice squamoso-articulata *Acetofella*. LINN. *Sp.* 1. HALL. 41. 364. *Trifolium acetosum vulg.* C. B. *Pén.* 330. *Oxys flore albo.* I. R. H. 88. Luiula. BLACKW. T. 305. Copiose in fylvis. P.

H *Oxalis* scapo vñifloro, foliis ternatis, radice bulbosa. *purpurea*. LINN. *Sp.* 2. *Oxys bulbosa, Africana, rotundifolia, caulibus & floribus purpureis amplis.* COMM. *Hort.* 1. p. 41. T. 21. In *Aethiopia*. P.

H. *Oxalis* pedunculis vñifloris, caule dichotomo: *incarnata*. LINN. *Sp.* 5. *Oxys bulbosa Aethiopica minor, folio cordato, flore albedo purpurascens.* COMM. *H.* 1. p. 43. T. 22. In *Aethiopia*. P.

GERANIVM. LINN. 746:

Calyx quinquefolius. Petala quinque. Filamenta laminum decem, quorum in aliis tria, in aliis quinque antheris carent, basi vix coälita. Capsulae quinque monospermae, longe caudatae & axi-
com-

communi applicatae. Stylus in quinque stigmata fissus.

I. Pedunculis vnifloris.

Geranium pedunculis vnifloris, foliis palmatis. HALL. H. 365. n. 1. Geranium pedunculis vnifloris, foliis quinquepartitis trifidis orbiculatis. (*Sanguineum*.) LINN. Sp. 38. In M. Altenstollberg. Prope specum Baumannianam. P.

H. Geranium pedunculis vnifloris, foliis triangularibus, quinquelobis, serratis. HALL. Comm. Gott. T. I. p. 225. Geranium pedunculis subvnifloris, foliis quinquepartitis, acutis, foliolis pinnatifidis. (*Sibiricum*.) LINN. Sp. 37. A.

II. Pedunculis bifloris, staminibus decem fertilibus.

Geranium pedunculis bifloris, petalis integris longitudine calycis, caule prostrato, foliis reniformibus incis. *rotundifolium*. LINN. Sp. 36. Geranium foliis reniformibus semiquinquelobis, lobis trifidis. HALL. Gott. 137. Helv. 365. n. 2. Ger. folio malvae rotundo. B. Pin. 318.

B. Geranium foliis reniformibus, semipinnatifidis, caule erecto, umbellato flore. HALL. H. 366. n. 3. Geran. columbinum maius, flore minore coerulesco. VAILL. Par. 79. T. 15 f. 1. Ad vias. A.

Geranium foliis mollibus, reniformibus, semiquinquefidis, lobis semitrifidis obtusis. HALL. H. 366. n. 5. Geranium pedunculis bifloris foliisque rameis alternis, caule ramoso diffuso, calycibus muticis.

tricus. (molle.) LINN. Sp. 33. Ger. columbinum vil-
losum, petalis bifidis purpureis. VAILL. Par. 79.
T. 15. f. 3. Ad vias & in agris. A.

Geranium foliis ad nervum quinquefidis, pedunculis brevioribus, caule erecto, HALL. J. c. n. 6. Geranium pedunculis bisloris, foliis quinquepartito-multifidis, caule glabro erectiusculo. (dissectum.) LINN. Cent. I. Plant. n. 62. VAILL. T. 15, f. 2. In agris A.

Geranium foliis ad nervum vsque quinquefidis,
pedunculo longissime, prostrato caule. HALL. H.
367. n. 7. columbinum. LINN. Sp. 35. VAILL. T.
15. f. 4. Circa Rafemühle, Herperbaufen, A.

Geranium pedunculis bifloris, foliis peltatis multipartitis rugosis pinnato-laciniatis acutis. HALL. H. n. 8. *pratense*. LINN. Sp. 29. Ger. *batrachioides* f. *Gratia Dei Germanorum*. B. Pin. 318. in pratis vulgaris, *Elizabet* *vel* *flavesc.*
P. flore albo & coccineo striato. In pratis. P.

Geranium foliis rugosis, semitrilobis, lobis dentatis, lateralibus bipartitis, petalis purpureis. HALL. *l. c. n. 9. sylvaticum*. LINN. *Sp.* 28. Ger. *batrachioide*. folio *Aconiti*. B. *Pin.* 317. (In aquosis, P.

H. *Geranium pedunculis bifloris*, foliis quinquelobis, lobis medio dilatatis, petalis emarginatis reticulatis. *versicolor*. LINN. *Cent. 1. n. 61. Ger. Romanum versicolor f. striatum*. RAIL *Hist. 1063. In Italia. P.*

H. Geranium pedunculis bifloris, foliis caulinis trilobis integris serratis, supremis subsessilibus. LINN. Sp. 27. *nodosum*. B. Pin. 318. In Gallia.

註。

H. *Geranium foliis quinquelobis serratis, floribus paniculatis atropurpureis*. HALL. *Helv.* 368. n. 10. Ger. *phaeum*. LINN. *Sp.* 26. Ger. *fusum petalis reflexis*. I. R. H. In *Helvetia*. P.

H. *Geranium pedunculis bifloris, calycibus inflatis, pistillo longissimo. macrorhizum*. LINN. *Sp.* 25. Ger. *batrachioides odoratum*. I. R. H. B. *Pin.* 318: P.

H. *Geranium foliis septilobis, lobis obtusis, dentatis, capsulis hirsutissimis*. HALL. *Gott.* 139. Ger. *Bohemicum*. LINN. *Cent.* II. n. 176. Ger. *batrachioides Bohemicum, capsulis nigris hirsutis*. DILL. *Elth.* 159, T. 133. f. 160. P.

Geranium foliis quinque-trivepartitis, lobis pinnatifidis, calycibus hirsutis decemangulatis. robertianum. LINN. *Sp.* 31. HALL. *H.* 369. n. 12. BLACKW. T. 480. Ad muros & rupes. P.

III. Pedunculis multifloris.

Geranium procumbens, pedunculis multifloris, calycibus pentaphyllis, foliis pinnatis, pinnis semipinnatis dentatis. cicutarium. LINN. *Sp.* 21. a. HALL. n. 14. Ger. *cicutae folio minus & supinum*. B. *Pin.* 319. In hortis & agris. A.

H. *Geranium procumbens pedunculis multifloris, calycibus pentaphyllis, foliis pinnatis, pinnis incisis dentatis*. HALL. *H.* n. 13. *Gott.* 140. LINN. *Sp.* 21. (β. Ger. *moschatum*. B. L. c. Riv.

H. *Geranium procumbens, pedunculis bi- & trifloris, foliis ovatis hirsutis crenato-incisis obtusis*. ROYEN. p. 451, n. 12. PLUK. *Phyt.* 301. f. 4.
Haec

Haec esse videtur species, quam descripsi in *Comment. Soc. Sc. Gott. T. V.* Planta supina. Petioli & pedunculi breves, crassiusculi. Corollae plurimae apetalae mihi sunt visae, in paucioribus vix vnum alterumve petalum album, calyce minus, occurrit. Capsulae hirsutae. Stamina quinque sterilia.

H. *Geranium pedunculis multifloris, foliis cordatis sublobatis, incis. malacoides.* LINN. *Sp.* 22. *Ger. folio Althaeae.* B. *Pin.* 318. *malacoides.* LOB. *ic.* 662. A.

β. *Geranium Cbium, vernal, Caryophyllatae folio.* T.

H. *Geranium pedunculis multifloris, foliis lobatis, lobis acutis serratis.* HALL. *Gott.* 140. *gruinum.* LINN. *Sp.* 23. *Ger. latifolium, acu longissima.* B. *Prodr.* 138. *Ger. Creticum.* RIV. In *Creta.*

H. *Geranium pedunculis multifloris, foliis pinnatis, pinnatifidis, obtusis, ciconium.* LINN. *Cent.* I. n. 60. HALL. *Helv.* n. 15. *Ger. apulum Coriandri-folium.* COL. *Ecphr.* L. p. 136. T. 135. In *Europa australi.* Pinnae minimae decurrentes maioribus intermixtae. Folia supina. Acus longissima. A.

IV. Pedunculis multifloris, calycibus monophyllis.

H. *Geranium foliis hirsutis, duplicato pinnatis, pinnulis incis, floribus maculosis, radice subrotunda.* HALL. *Gott.* 141. *triste.* LINN. *Sp.* 1. CORNVT. *Canad.* 109. T. 110. *Ger. noctu olens foliis Myrrhidis.* BREYN. *Cent.* 1. p. 126. T. 58.

H. *Geranium calycibus monophyllis, foliis duplicato pinnatis, pinnulis laxè multifidis.* HALL. *G.* 141. *myrrhifolium.* LINN. *Sp.* 5. *Ger. myrrhidis folio, flore magno striato.* BREVN. *Cent.* 1, p. 129. *T.* 59. COMMEL. *H.* II. p. 125. *T.* 63. P.

H. *Geranium calycibus monophyllis tetrapetalon, foliis bipinnatis.* HALL. *l. c.* *Ger. Afr. Coriandri folio, floribus incarnatis.* HERM. *Lugdb.* 279. *T.* 280. RIV. *T.* 106. Corolla fere papilionacea. P.

H. *Geranium calycibus monophyllis, foliis dentatis, trifidis, medio lobo semitrilobo, lateralibus minoribus incis.* HALL. *l. c.* *Ger. fulgidum.* LINN. *Sp.* 2. *Ger. Afr. Alceae folio, flore coccineo, fulgidissimo.* DILL. *H. Elth.* 156. *T.* 130. f. 137. TILL. *Pisf.* 68. *T.* 26. Fr.

H. *Geranium calycibus monophyllis, caule nodoso, foliis glaucis sinuatis.* HALL. p. 142. *Geran. gibbosum.* LINN. *Sp.* 8. *Ger. Afr. noctu olens Aquilegiae foliis.* HERM. *Lugdb.* 284. *T.* 285. RIV. *T.* 99. Fr.

H. *Geranium calycibus monophyllis, foliis cucullatis, dentatis.* *cucullatam.* LINN. *Sp.* 7. DILL. *Elth.* 155. *T.* 129. f. 156. *Geran. Afr. ibisci folio rotundo, carlinae odore.* HERM. *Lugdb.* 274. *T.* 275. Folia inferiora reniformia; non angulata. P.

H. *Geranium procumbens, calycibus monophyllis, foliis cordatis, glabris, obtusis, serratis.* HALL. *l. c.* *Geran. grossularioides.* LINN. *Sp.* 16. *Ger. Afr. vuæ crispæ folio, floribus exiguis rubellis.* HERM. *Lugdb.* 287. *T.* 289. RIV. *T.* 107. P.

H. Geranium calycibus monophyllis, foliis hirsutissimis, oblongis, semiquinquelobis, vndatis. HALL. l. c. Geran. *capitatum*. LINN. Spec. 13. Ger. Afr. *frutescens*, *malvae folio laciniato*, odorato. HERM. Lugdb. 277. T. 278. RIV. T. 98. Fr.

H. Geranium calycibus monophyllis, foliis oblongis, semiseptifidis, serratis, mollissimis. HALL. l. c. Ger. Afr. *Betonicae folio laciniato*, floribus incarnatis. HERM. Lugdb. 278. RIV. T. 105.

H. Geranium calycibus monophyllis, foliis cordiformibus semitrilobis, serratis, hirsutis. HALL. p. 143. Ger. *vitifolium*. LINN. Sp. 12. Ger. Afr. *arbor. vitis folio*, odore *melissae*. DILL. Elth. 152. T. 126. f. 153. Fr.

H. Geranium calycibus monophyllis, foliis cordiformibus, obtusis, serratis, mollissimis. HALL. l. c. Ger. *odoratissimum*. LINN. Sp. 15. Ger. Afr. *humile*, *folio fragantissimo molli*. DILL. Elth. 157. T. 131. f. 138.

H. Geranium calycibus monophyllis, foliis pinguibus tomentosis, reniformibus obtuse crenatis. HALL. l. c. Geran. *inquinans*. LINN. Spec. 3. Ger. Afr. *arbor. Malvae folio pingui*, flore *rocceae*. DILL. Elth. T. 125. f. 151. MART. Cent. 3. f. 3. RIV. T. 97. Fr.

H. Geranium calycibus monophyllis, foliis cordato-orbicularibus incis, zona notatis. *zonale*. LINN. Sp. 11. a. Ger. *foliis reniformibus obtuse dentatis*, *nigrum circulum inscriptis*. HALL. 144. Ger. Afr.

arborescens Alchimillae folio, flore rubro. COMM. Prael. 51. T. 1. Fr.

β. *Geranium simile prioris, foliis album circum inscriptis.* HALL. 143. *Geran. Afr. Alchimillae folio maculato.* RIV. T. 102. Fr.

h. *Geranium calycibus monophyllis, foliis pilosis reniformibus, quinquelobis, lobis tridentatis.* HALL. G. 144. *Ger. alchimilloides.* LINN. Sp. 14. *Ger. Afr. alchimillae hirsuto folio, flosculis albidis.* HERM. Lugdb. 282. T. 283. Fr.

h. *Geranicum calycibus monophyllis, foliis glabris subovatis carnosis crenatis. acetosum.* LINN. Sp. 10. *Ger. Afr. frutescens, folio crasso & glauco, acetosae sapore.* COMM. Prael. 54. T. 4.

h. *Geranium calycibus monophyllis, foliis pentagoniis integerrimis glabris peltatis. peltatum.* LINN. Sp. 10. *Geran. Afr. foliis inferioribus asari, superioribus staphidis agriae maculatis splendentibus, & acetosae sapore.* COMM. Prael. 52. T. 2. Fr.

LINVM. LINN. 349.

Calycis folia & petala, & stamina quinque, totidemque styli. Fructus fere globosus, decemlocularis.

Linum calycibus capsulisque mucronatis, petalis crenatis, foliis lanceolatis alternis, caule subsolitario. usitatissimum. LINN. Sp. 1. *Linum ramis foliisque alternis lineari-lanceolatis, radice annua.* HALL. H. 370 n. 1. L. *salivum.* B. Pin. 214. Vbi que colitur. A.

9. *Linum sativum, latifolium, Africanum, fructu maiore.* I. R. H. 339.

H. *Linum calycibus capsulisque obtusis, foliis alternis lanceolatis integerrimis. perenne.* LINN. *Sp.* 2. HALL. *H.* n. 2. *L. perenne maius, coerulcum, capitulo maiore.* MORIS. *H.* II. p. 573. I. R. H. In *Sibiria. Anglia. Helvetia.* P.

Linum calycibus acuminatis, foliis sparsis lineari-setaceis retrorsum scabris. tenuifolium. LINN. *Sp.* 5. *Lin. perenne, foliis angustissimis, calyce aristato.* HALL. *H.* 371. n. 3. *Linum sylv. angustifolium flore magno.* I. R. H. 340. In siccis apricis, v. g. über der *Papiermühle, am Wege nach Robringen.* &c. P.

H. *Linum foliis oppositis lineari-lanceolatis, floribus terminalibus pedunculatis. Africanum.* LINN. *Sp.* 15. ROYEN. 434. *Linum Afric. luteum, foliis coniugatis.* BOERH. p. 284. In *Africa.*

Linum foliis coniugatis, petalis albis acutis. HALL. *H.* 372. n. 4. *Linum foliis oppositis ovato-lanceolatis, caule dichotomo, corollis acutis.* LINN. *Sp.* 17. *Linum catharticum.* BLACKW. *T.* 268. In pratis & gramineis. A.

Linum ramosissimum, flore tetrapetalo. HALL. *G.* 147. *Linum foliis oppositis, caule dichotomo, floribus tetrandris tetragynis. (Radiola.)* LINN. *Sp.* 19. *Chamaelinum vulg.* VAILL. *Parif.* 33. *T.* 4. f. 6. In sabulosis, v. g. *Cellensibus & circa Bërolinum.*

DROSERA. LINN. 351. Rorella.

HALL.

Calyx semiquinquefidus, petala subovata. Stamina & styli quinque. Folia roriferis villis ciliata.

Drosera scapis radicatis, foliis orbiculatis, rotundifolia. LINN. *Sp.* 1. Rorella *caule simplici, foliis rotundis retusis*. HALL. *H.* 372. *n.* 2. Ros Solis, *folio rotundo*. B. *Pin.* 357.

Drosera scapis radicatis, foliis oblongis. longifolia. LINN. *Sp.* 2. Rorella *foliis ellipticis*. HALL. *L.c.n.* 1. Ros folis f. *Salifiora*. THAL. *Herc.* 116. *T. 9. f.* 2. Vtraque in paludosis circa M. *Bructerum*.

FAGONIA. LINN. 475.

H. *Fagonia* spinosa, foliis lanceolatis plauis linearibus. retusa. LINN. *Sp.* 1. Trifolium *spinosum creticum*. CLVS. CCXLII. c. B. *Prodr.* 142. In Oriente. P.

TRIBVLVS. LINN. 476.

Petala quinque. Stylus simplex. Fructus ex quinque capsulis aculeatis compositus.

H. *Tribulus* foliolis sexiugatis subaequalibus, terrestris. LINN. *Sp.* 3. *Tribulus terrestris*. LOR. *ic.* 2. p. 841. In Europa australi. A.

MELIA. LINN. 473. Azedarach. *T.* 387.

Calyx minimus. Nectario cylindrico decem stamina se inferunt. Stylus simplex.

H.

H. *Melia* foliis decempartitis. HALL. G. 161. *Melia foliis bipinnatis*. LINN. Sp. I. *Azedarach*. DOD. *Pempt.* 848. I. R. H. 616. In *Syria*. Fr.

ZYGOPHYLLVM. LINN. 474. *Fabago*.

T. 135.

Petala ovata. Circa stamina decem totidem squamae erectae; conniventes. Stylus simplex.

H. *Zygophyllum* foliis petiolatis, capsulis prismatiko-pentaëdriis. LINN. Sp. I. *Fabago Belgarum*. I. R. H. *Capparis Fabago*. DOD. *Pempt.* 471. In *Syria*. P.

DYPLOSTEMONES ODORAE.

Omnes, ut priores, dyplostemonas, flores in plerisque numero partium quinario- & quaternario ludunt. Loculi fructus tot, quot petala. Omnes graveolentes.

RUTA. LINN. 469.

Petala concava. Capsula punctata quadri- aut quinqueloba, cruce insculpta.

H. *Ruta* foliis decompositis. *graveolens*. LINN. Sp. I. *Ruta hortensis latifolia*. I. R. H. B. *Pin.* 336. BLACKW. T. 7. Petala oris laciniatis. P.

PEGANVM. LINN. 530. *Harmala*. TOURN.

Rutae affinis. Sed stamina quindecim, capsula trigona, trilocularis. Stigmata tria.

H. *Peganum* foliis multifidis. LINN. Sp. I. *Harmala* DOD. *Pempt.* 121. *Purg.* 420. LOB. p. 55. In *Europa* australiori & *Syria*. P.

MONOTROPA. LINN. 477. Hypopitys.

HALL. 411.

Habitus aphyllus. Flores terminales quinquepartiti, laterales quadripartiti.

Monotropa floribus lateralibus pctandris, terminali decandro. LINN. *Sp.* 1. Hypopitys *spica nutante*. HALL. *H.* p. 411. Orobanche *Hypopitys*. C. B. *Prodr.* 31. In umbrosis sylvis parasitica radicum.

ADOXA. LINN. 450. Moschatellina HALL. 412.

TOURN. T. 68.

Flores in spicas cubicas collecti.

Adoxa. LINN. *Sp.* p. 367. Moschatellina. HALL. *H.* 412. In dumetis versus *Eichenkrug*.

PARIS. LINN. 449. HALL. 412.

Omnes partes floris numerum quaternarium observant. Fr. bacca.

Paris foliis ^{quadrifolius} quaternis. LINN. *Spec.* p. 367. HALL. *H.* 412. Herba Paris. I. R. H. BLACKW. *T.* 286. Solanum *quadrifolium bacciferum*. B. *Pin.* 167. In umbrosis sylvis & dumetis.

DYPLOSTEMONES ALSINACEAE.

Calyx quinquefolius. Petala quinque patentia. Stamina his tot, paucioribus exceptis. Tubae tres, quatuor & quinque. Capsula unilocularis. Folia opposita.

ALSINE.

ALSINE. HALL. H. 385. *Arenaria*
spergula. *Sagina*. *Silene*. *Holcus*.
Capfula conica. Calycis folia obtusiora.

I. Petalis integris ; tubis trinis.

Alfine foliis lanceolatis, trinerviis, petalis integris. HALL. H. n. 18. *Arenaria foliis subovatis acutis petiolatis nervosis. (trinervia.)* LINN. Sp. 4. Alfine plantaginis folio. 1. B. Hist. III. 364. In sylvis montanis. A.

Alfine foliis lanceolatis, petalis integris calycis magnitudine. HALL. H. n. 17. *Arenaria serpyllifolia*. LINN. Sp. 16. Alfine minor multicaulis. B. Pin. 251. In vallis & sylvis macilentis. A.

Alfine foliis perangustis, ad basin lateſcentibus. HALL. H. n. 15. *Arenaria foliis subulatis, caule paniculato, petalis calyce brevioribus.* LINN. Sp. 8. Alfine tenuifolia. 1. B. VAILL. Par. T. 3. f. 1. In M. Heymberg. A.

Alfine foliis fasciculatis, petiolis simplicibus, petalis integris. HALL. H. n. 9. *Spergula foliis oppositis subulatis laevibus; caulibus simplicibus. (nodosa.)* LINN. Sp. 3. Alfine nodosa Germanica. B. Pin. 251. In vdis circa Seeburg. A.

Alfine foliis coniugatis, angustis, linearibus, petalis integris. HALL. H. n. 14. *Arenaria*. LINN. Sp. 7. Alfine saxatilis & multiflora, capillaceo folio. HALL. T. 2. f. 3. Circa Herzberg & in sylva Hercynica.

Alfine tetrapetala, foliis angustis, in origine lateſcentibus. HALL. H. n. 20. *Sagina ramis procum-*

^{monanthus}
bentibus. LINN. Sp. 1. *Alfne minima flore fugaci.*

I. R. H.

Sagina speciosa 1711-12, cap. 169.

Sagina creta
1711-12, p. 169.

Alfne foliis filiformibus, vaginis ad foliorum
ortum squalentibus. HALL. H. n. 8. *Arenaria ru-*
bra. LINN. Sp. 6. *Spergula purpurea*. I. B. III. p. 722.
In agris siccioribus.

β. Eadem pulposior & altior, locis salis. v. g. in
prato prope *Haxle*.

II. Petalis integris, tubis quinque.

Alfne foliis verticillatis, petiolis ramosis, petalis
integris. HALL. H. n. 7. *Spergula arvensis*. LINN.
Sp. 1. *Alfne Spergula dicta maior*. B. Pin. 251.
In arvis. A. *1712 Gladstet v. n. abh. II. 100.*

III. Pentastemones.

Alfne pentastemon, petalis ferratis, umbellifera.
HALL. H. n. 6. *Holotheum floribus umbellatis*.
LINN. Sp. 3. DILL. nov. gen. p. 130. T. 6. TABERN.
ic. p. 233. In vallo urbis, hortis & agris. A.

Alfne pentastemon, petalis bipartitis, foliis ova-
to-cordatis. HALL. H. n. 5. LINN. Sp. 1. *Alfne*
media. B. Pin. 250. A.

IV. Petalis bipartitis, decastemones.

Alfne foliis cordatis, petalis bipartitis. HALL. H.
n. 4. *Stellaria foliis cordatis petiolatis, pedunculis*
ramosis. (nemorum.) LINN. Sp. 1. *Alfne altissima*
nemorum. B. Pin. 251. *Alfne maior*. TABERN. p.
707. In fossis. A.

+ *Alfne* foliis ovato-lanceolatis, petalis bipartitis.
HALL. H. n. 1. *Alfne longifolia, virginifolia locis*
pro-

proveniens. I. B. III. p. 365. *Alfine aquatica media*.
I. R. H. Super *Hardeggen*; in palude inter *Harsto*
& *Baronsen*. *Jam. n. 10. H. n. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100.*

Alfine foliis gramineis, petalis bipartitis. HALL.
H. 2. *Stellaria foliis linearibus, integerrimis*. LINN.
Sp. 5. *Gramen floridum minus*. TABERN. ic. 232.
In agris & palustribus inter Juncos. A.

Alfine foliis gramineis, asperis, petalis semibifidis. HALL. H. n. 1. *Stellaria Holostea*. LINN. Sp. 4.
Gramen *Fuchsii*. I. B. III. p. 361. In sylvis & dumetis.
Holostea umbellata. V. p. 126.

CERASTIVM. LINN. 518. Myosotis.
HALL, 383.

Petala bifida, Capsula vtp plurimum longior, incurva.

Cerastium (*Myosotis*) foliis oblongis, obtusis, petalis calyce maioribus. HALL. H. n. 3. *arvense*. LINN. Sp. 5. *Myosotis arvensis hirsuta, flore maiore*. VAILL. Par. 141. T. 30. f. 4. I. R. H. Ad vias. A.

Cerastium (*Myosotis*) foliis ovatis, acuminatis, flore calyci aequali. HALL. H. n. 4. *Myosotis hirsuta, parvo, albo flore*. I. R. H. VAILL. Par. T. 30. f. 1. A.

Cerastium (*Myosotis*) floribus pentandris, petalis emarginatis. LINN. Sp. 3. *Myosotis exigua pentastemon*. HALL. H. 5. *Myosotis arvensis hirsuta minor*. VAILL. Par. 142. T. 30. f. 2. Filamenta quinque sterilia, quinque antherifera. Styli tres. A.

Cerastium (Myosotis) foliis lanceolatis, caule dichotomo ramosissimo, capsulis erectis. LINN. *Sp.* 6. *Myosotis Hispanica segetum.* I. R. H. *Alfine corniculata.* CLVS. *Hist.* II. p. 184. A. an?

Cerastium (Myosotis) viscosa, foliis ovatis. HALL. *H.n.* 6. *Cerastium erectum villoso-viscosum.* LINN. *Spec.* 2. *Myosotis hirsuta altera viscosa.* VAILL. *Par.* 142. T. 30. f. 1. In macilentis. A.

H. *Cerastium (Myosotis) foliis connatis.* LINN. *Sp.* 1. *Myosotis orient. perfoliata, lychnidis folio.* DILL. *Elth.* 295. T. 217. f. 284. A.

Cerastium iniquitatum.

DYPLOSTEMONES CARYOPHYLLAEAE.

Calyx monophyllus, tubulosus, ore quinquefido. Petala quinque, vnguibus linearibus, bractea reflexa, latiori. Stamina decem, alterne maiora. Styli duo, tres, quinque. Capsula oblonga, unilocularis. Folia opposita, simplicia. Secundum Hallerum genera minora ex tubis constitui possunt.

VISCAGO. HALL. DILL. Cucubalus.

LINN. Silene. LINN.

Styli tres. Summitates saepe viscidae.

I. Petala simplicia sine corolla auricularum.

Cucubali. LINN.

H. *Viscago (Cucubalus) calycibus campanulatis, petalis distantibus, fructu colorato, ramis divaricatis.* LINN. *Sp.* 1. *Alfine scandens baccifera.* B. *Pin.* 250. *Cucubalus Plinii vulgo.* Circa Jenam. P.

Visca-

Viscago (*Cucubalus*) calycibus subglobosis glabris reticulato-venosis, capsulis trilocularibus, corollis ferratis, bifidis. ^{Achen} LINN. Sp. 2. HALL. H. 2. *Lychnis sylv. quae Beben album vulgo.* B. Pin. 205. In collibus saxosis, siccis. P.

H. Viscago calycibus ovatis glabris venoso-reticulatis, floribus subsessilibus, petalis calyce parum maioribus, denticulatis, radice annua, ^{radice} *Silene* LINN. Sp. 13. Viscago *vesicaria Cretica, flore parvo purpureo*. DILL. Elsb. T. 3 17. f. 409. A.

H. Viscago (*Cucubalus*) petalis bipartitis, floribus secundis decumbentibus, pedunculis solitariis oppositis erectis, caule simplicissimo. *Tataricus*. LINN. Sp. 8. *Lychnis septemtrionalium, foliis hyssopi, floribus uno versu dispositis.* HALL. G. 150. P.

H. Viscago foliis imis petiolatis, ovatis, floribus paniculatis, petalis emarginatis. HALL. Gott. ed. prior. c. ic. in app. & edit. ult. p. 150. *Cucubalus sibiricus* L. 81. 2. p. 492.

H. Viscago foliis imis petiolatis, floribus verticillatim spicatis, petalis indivisis, sexu distincta. HALL. H. n. 5. *Cucubalus Otites*. LINN. Sp. 10. *Lychnis viscosa, flore muscoso.* B. Pin. 206. In *Marchia*. Prope arcem *Regenstein*.

H. Viscago foliis imis petiolatis, floribus verticillatim spicatis, petalis indivisis, androgyna. HALL. G. p. 165.

Viscago (*Cucubalus*) floribus spicatis alternis subsessilibus, corollis obsoletis nudis. ^{pediculis} LINN. Sp. 12. *Lychnis sylv. alba, floribus reflexis.* MAGN. Monsp. II. T. 170.

*Silene*II. Petala denticulata. *Silene* LINN.

H. Viscago calyce duplo petalis longiore, floribus umbellatis. HALL. *H. n.* 3. *Silene floribus fasciculatis fastigiatis, foliis superioribus cordatis glabris.* (*Armeria.*) LINN. *Sp.* 24. In *Valesia, Galp.* A.

Viscago foliis imis petiolatis, ovatis, petalis bifidis se convolventibus. HALL. *Helv. n.* 4. *Silene foliis lanceolatis, caule paniculato, floribus nutantibus, calyce striato, corolla involuta.* LINN. *Sp.* 7. In faxosis circa *Gottingam.*

H. *Silene* petalis bifidis, corolla subcoadunata, floribus secundis, calycibus erectis raris pilosis, pedunculis oppositis trifloris, ramis alternis. *amensna.* LINN. *Sp.* 8. *H. Vps.* 114. In *Sibiria.* A.

H. *Silene* petalis bifidis, caule fruticoso, foliis lato-lanceolatis glabris, panicula trichotoma. *fruticosa.* LINN. *Sp.* 9. *H. Vps.* 111. *n.* 2. *Lychnis sylv. myrtifolia Beben albo similis.* B. *Pin.* 205. *Ocymoides fruticosum.* GAMER. *Hort.* 109. *T.* 33. Fr.

H. Viscago (*Silene*) foliis radicalibus cochleariformibus obtusissimis, caule subverticillato. *gigantea.* LINN. *Sp.* 10. *Lychnis Graeca, Sedi arbore-scentis folio & facie, flore albo.* TOURN. *Cor.* 23. WALTH. *H. p.* 32. *T.* XI. P.

H. Viscago caule foliisque hirsutis ovato lanceolatis, floribus raris, petalis semibifidis. *Lychnis ocymastri facie, flore viridi.* HERM. *Parad.* *T.* 199.

H. *Viscago* foliis inferioribus obverse ovatis, superioribus lineari-lanceolatis, calycibus erectis sulcatis, petalis emarginatis. *Silene cretica*. LINN. *Sp.* 20. *H. Vps.* 112. *Viscago foliis inferioribus bellidis, superioribus tunicae, calyce strictiore & turgidiore*. DILL. *Elth.* 422. *T.* 314. f. 404. 405. In *Creta*.

H. *Viscago* (*Silene*) foliis obverse ovatis, floribus spicatis alternis secundis sessilibus, petalis bifidis inaequalibus. *nocturna* LINN. *Sp.* 4. *Viscago hirta noctiflora, floribus obsolete spicatis*. DILL. *Elth.* *T.* 310. f. 400. In *Pensylvania*. A.

H. *Viscago* (*Silene*) hirsuta, petalis dentatis, floribus erectis, fructibus reflexis subsessilibus alternis. *Lusitanica*: LINN. *Sp.* 2. *Viscago hirsuta Lusitanica, stellato flore*. DILL. *Elth.* *T.* 311. f. 401. In *Lusitania*. A.

H. *Viscago* (*Silene*) petalis integerrimis subrotundis, fructibus erectis alternis. *quinguevulneca*. LINN. *Spec.* 3. *Lychnis hirsuta minor, flore variegato*. HALL. *Gott.* 152. In *Hispania*. A.

H. *Viscago* (*Silene*) calycibus fructiferis pendulis inflatis, angulis decem scabris. *pendula*. LINN. *Sp.* 14. *Viscago hirta Sicula, lychnidis aquaticae facie, supina*. DILL. *Elth.* *T.* 312. f. 404. Folia inferiora spatulata, superiora lanceolata, caulis debilis. In *Creta, Sicilia*. A.

H. *Viscago* (*Silene*) calycibus fructus globosis acuminatis striis triginta foliis glabris. *conoides*.

Et Silene amara — foliis, noctiflora, petalis LINN.
bifidis. In creta non aequa adspici

LINN. Sp. 11. *Lychnis sylvestris latifolia*, calycibus turgidis striatis. B. Pin. 205. A.

Vilcago (*Silene*) erecta laevis, calycibus subglobosis glabris venosis, corollis inapertis. *rubella*. LINN. Sp. 18. H. Vps. 112. Vilcago Lusit. flore rubello vix conspicuo. DILL. Elth. T. 314. f. 406. A.

add. Silene. Deben. p. 189.

LYCHNIS. HALL. 376. *Lychnis* LINN. 517.

Agrostema. LINN. 516.

Styli quinque. ~~*Agrostema*~~ ^{*Agrostemma*} Segmenta calycis coriacei flore longiora; *Lychnis* LINN. Calyx subrotundus inflatus, filamenta staminum alis aucta.

H. *Lychnis* (*Agrostema*) tomentosa, foliis ovato-lanceolatis, petalis coronatis emarginatis. ^{*coronaria*} LINN. Sp. 3. HALL. H. n. 1. *Lychnis coronaria* Dioscoridis sativa. B. Pin. 203. In Helvetia. Italia. A.

Lychnis (*Agrostema*) hirsuta, calycibus corollam aequantibus, petalis integris nudis. ^{*sp. f. Ang.*} LINN. Sp. 1. HALL. H. 2. *Lychnis segetum maior*. B. Pin. 204. Nigellastrum. DODON. Coron. p. 49. Inter segetes. A.

^{*Lychnis*}
Lychnis sexu distincta, foliis ovatis, acutis. HALL. H. 3. ^{*scabra*} LINN. Sp. 6. *Lychnis sylvestris purpurea simplex*. β. eadem alba. n. Pin. 204. γ. Variet. fl. pleno. Ocymastrum rubrum & album. TABERN. 299. Alba circa Rosdorf ad viam.

H. *Lychnis* floribus umbellatis, petalis breviter quadrifidis. HALL. Gott. 153. *Lychnis floribus fasciculatis fastigiatis*. (*Chalcedonica*.) LINN. Sp. 1. Flos Constantinopolitanus. DOD. Pempt. 178. flore coccineo aut incarnato. In Oriente. A.

Lychnis

Lychnis petalis quadrifida, longissimis, fructu subrotundo. LINN. *Sp.* 2. HALL. *H.* 378. Caryophyllus *pratensis*, flore laciniato simplici s. flor. *Cuculi aut flore pleno*. B. *Pin.* 210. In pratis. P.

Lychnis petalis integris, fructu quinqueloculari. *Viscaria*. LINN. *Sp.* 3. *Lychnis sylv. viscosa rubra angustifolia*. B. *Pin.* 205. Circa Elbingero & Blanckenburg.

SAPONARIA. LINN. 499.

Calyx oblongus. Vngues petalorum angusti, capsula cylindrica. Styli duo.

Saponaria calycibus cylindricis, foliis ovato lanceolatis. *officinalis*. LINN. *Sp.* 1. *Saponaria foliis trinerviis, floribus umbellatis*. HALL. *H.* 2. *Saponaria vulgaris*. In vallo vrbis. P.

Saponaria caule brachiato, foliis ovatis acuminatis sessilibus, calyce pentagono turgido. HALL. *H.* 3. *Vaccaria*. LINN. *Sp.* 2. *Lychnis segetum rubra, foliis perfoliatis*. B. *Pin.* 204. Inter segetes. A.

h. *Saponaria caule dichotomo*, foliis lanceolatis, petiolatis, floribus solitariis, calycibus cylindricis villosis. *orientalis*. LINN. *Sp.* 4. *H. Vps.* 106. n. 2. *Lychnis orientalis annua supina, antirrhini folio, flore minimo purpurascens*. DILL. *Elb.* T. 167. f. 204. A.

GYPSOPHILA. LINN. 498.

Calyx campanulatus. Petala ovata, vnguibus minimis. Filamenta staminum subulata.

H. *Gypsophila* foliis lanceolatis scabris, corollis revolutis, (staminibus in flore latentibus.) *paniculata*. LINN. Sp. 3. In *Sibiria*.

H. *Gypsophila* foliis lanceolatis subtrinerviis re-
ctis. *altissima*. LINN. Sp. 4. Gen. nov. 1103. H. Vpf.
107. In *Sibiria*. P.

Gypsophila foliis lanceolato-linearibus enerviis obsolete triquetris, laevibus obtusis secundis. *fasci-
giata*. LINN. Sp. 5. *Saponaria petalis ovatis, fo-
liis glaucis pulposis linearibus*. HALL. Fl. Jen. Rupp.
117. T. 2. f. 1. *Symphytum petraeum*. THAL. Herc.
p. 117. In M. *Altenstollberg*. P.

H. *Gypsophila* foliis lanceolatis semiamplexicau-
libus. *perfoliata*. LINN. Sp. 6. *Spergula multiflora,
foliis inferioribus saponariae, superioribus Beben si-
milibus*. DILL. Elth. T. 276. f. 357. In *Hispania* P.

Gypsophila foliis lineari-lanceolatis planis, caule dichotomo, pedunculis bifloris, petalis emarginatis. *rigida*. LINN. Sp. 6. *Saponaria*. HALL. H. 8. *Lych-
nis annua minima, flore carneo lineis purpureis di-
stincto*. I. R. H. Inter segetes exeunte aestate. A.

H. *Gypsophila* caule dichotomo, foliis subulatis, calycibus quinqueangularibus, squamis ad basin ac-
cessoriis. Caryophyllus muralis minimus. B. Pin.
211. Videtur esse *Gypsophila* ^{minima} LINN. Sp. 7. Est
enim squamae ad basin calycis adsint, summa tamen
cum priori specie analogia, & vita annua impediunt,
quo minus ad *Dianthus saxifragum* LINN. refer-
ri possit. A. *Pro Dianthe saxifrago Haber Willd. Obs. 1. p. 24
est spec. 2 p. 581. 1. Gypsophila saxifraga*
Willd.

DIAN.

DIPLOSTEMONES CALYCIFLORAE.

Calyx monophyllus. Petala patentia & stamina bis tot calyci inserta. Stigmata tot, quot loculi fructus sub calyce positi.

LYTHRUM. LINN. 532. *Salicaria*.

TOURN. 129.

Calyx tubulosus, segmentis sex maioribus, sex minoribus alternis, introsum flexis. Petala sex.

Lythrum (*Salicaria*) foliis oppositis cordato-lanceolatis, floribus spicatis dodecandris. LINN. *Sp.* 1. HALL. *H. n.* 1. *Salicaria vulgaris purpurea foliis oblongis.* 1, R. H., *Lyfimachia spicata purpurea.* B. *Pin.* 246. Ad ripas aquarum. P.

Lythrum (*Salicaria*) foliis alternis linearibus, floribus hexandris. *Hyssipifolia.* LINN. *Sp.* 5. HALL. *Fl. Jen. Rupp. T.* 2. f. 3. B. *Pin.* 246. Auf Aeckern hinter der Graetzels Mühle.

PEPLIS. LINN. 401.

Calyx campanulatus, caeterum priori similis. Petala fugacia, exigua. Stamina sex.

Peplis foliis coniugatis. *Portula.* LINN. *Spec.* 1. *Glaux palustris, flore striato clauso, foliis portulacae.* PETIT. *Gen.* 43. *T.* 43. MICH. *gen.* 21. *T.* 43. VAILL. *Par.* 80. *T.* 15. f. 5. In paludibus M. Hagen. *Ima Giebelhausen.*

H. *Dianthus floribus solitariis squamis calycinis subulatis, tubum aequantibus, corollis crenatis.*
^{Chinensis} LINN. *Sp.* 8. *Caryophyllus Chinensis supinus, Leucocoi folio, flore unico.* TOURN. *Act. Par.* 1705. p. 348. f. 5. A.

H. *Dianthus floribus subsolitariis, squamis calycinis lanceolatis quaternis, corollis crenatis. glaucus.* LINN. *Sp.* 7. *Tunica ramosior, flore candido cum corolla purpurea.* DILL. *Elth.* 400. T. 298. f. 348. In *Anglia.* P.

Dianthus floribus solitariis, squamis calycinis subovatis brevissimis, corollis multifidis faucē pubescentibus. plumarius. LINN. *Sp.* 9. *Tunica.* HALL. n. 5. *Caryophyllus flore tenuissime dissecto.* B. *Pin.* 209. In pascuis nemorosis.

X H. *Dianthus floribus solitariis, foliis subulatis, squamis calycinis muticis, floribus profunde fimbriatis, lacteis.* GMEL. *Hall. Got.* p. 156. P.

Dianthus floribus subsolitariis, squamis calycinis tubo parum minoribus, corollis serratis, punctatis. Caryophyllus simplex supinus latifolius. B. *Pin.* 208. HALL. G. 156. *Caryophyllus minor pulchellus, supinus, maculis aureis argenteisve adspersus.* LOB. ic. 444. Prope *Eichenkrug, Rheinhausen* & alibi in arenosis.

H. *Dianthus caulibus unifloris, squamis calycinis ovatis, corollis multifidis, foliis linearibus. arenarius.* LINN. *Sp.* 10. *Tunica foliis brevibus rigidis, caule unifloro.* HALL. H. n. 8. *Caryophyllus sylv. humilis, flore unico.* B. *Pin.* 209. LOB. ic. 405. *sylv.* L. CLVS. I. p. 282. DILL. *Elth.* 402. In *Marchia.*

~~Dianthus Xanthoxanthus~~ 1794.

DIPLO-

H. *Oenothera* foliis ovato-lanceolatis planis.
^{liennes} LINN. *Spec.* 1. *Onagra latifolia*. 1. R. H. *Lythima-*
chia lutea corniculata. B. *Pin.* 245. In *Marchia*. P.

H. *Oenothera* foliis lanceolatis, capsulis ventri-
 coso-conicis. An *Onagra angustifolia*, caule rubro,
flore minori? TOURN. *Inst.* 302. Flores & fructus
 densius congesti, caulis & calyx rubro tinctus. Fru-
 ctus priori fere aequalis, flos autem triplo
 minor, *Oenothera parviflora* Linn. *2. spec.* 3.

H. *Oenothera* foliis glabris, flore suave purpu-
 reo, *Comm. Soc. R. Sc. Gott.* 1751. p. 224. In *Vir-*
ginia.

H. *Oenothera* foliis lanceolatis undulatis, *mollis-*
sima. LINN. *Sp.* 2. *Onagra bonariensis villosa, flo-*
re mutabili. DILL. *Elth.* 297. T. 219. f. 286.

EPILOBIVM. LINN. 246.

Capsulae cylindricae papposis seminibus foetae
 insitit flos similis priori, sed calyx est quadrifolius,
 tubo carens.

Epilobium spicatum, flore difformi, foliorum
 nervo eminente, ora leviter serrata. HALL. *H. n.* 1.
Epilobium flore difformi, staminibus & pistillis
declinatis. LINN. *Fl. Suec. ed. pr.* 304. *Chamae-*
nerium vulgare latifolium. 1. R. H. B. *Pin.* 245.
 In fylvis.

Epilobium foliis ovato-lanceolatis, serratis, le-
 vissime hirsutis. HALL. *H. n.* 5. *Epilobium foliis*
oppositis ovatis dentatis. (montanum.) LINN. *Sp.* 4.
Chamaenerium glabrum majus. 1. R. H. In umbrosis,

Epilobium foliis ellipticis obtuse lanceolatis, retum leve. HALL. *Gott.* p. 162. *Epilobium foliis oppositis lanceolatis, integris petalis, caule erecto. (palustre.)* LINN. *Sp.* 6. *Chamaenerium glabrum angustifolium.* DILL. *Cat. Giess.* p. 208. In palude inter *Barenfen* & *Gladebeck*.

Epilobium foliis longis, dentatis, villosis, flore minore. HALL. *H. n.* 7. *Epilobium hirsutum.* LINN. *Sp.* 2. β . *Chamaenerium villosum, maius, parvo flore.* I. R. H. In fossis.

Epilobium foliis longis dentatis, villosis, flore maiore. HALL. *H. n.* 8. *Epilobium hirsutum.* LINN. *Sp.* 2. α . *Chamaenerium villosum, flora magno purpureo.* I. R. H. Ad rivos.

DIPLOSTEMONES BICORNES.

Calyx exiguus. Corolla monopetala limbo quinquefido. Stamina aut receptaculo, aut basi corollae tantum affixa. Antherae binas setas exferunt. Tuba simplex. Fructus quadri- aut quinquelocularis.

Vaccinium. LINN. 434.

Corolla campanulata, limbo quadrifido. Germen infra florem. Fr. Bacca umbilicata.

Vaccinium foliis integerrimis revolutis ovatis, caulibus repentibus filiformibus, corolla ad basin fere quadrifida. LINN. *Sp.* 11. *Oxycoccus* HALL. *H.* p. 413. *Vitis Idaea palustris.* B. *Pin.* 471. Corollae segmenta per aetatem ita revolvuntur, ut flos tetrapetalos videatur. In turfæis circa M. *Bruckeri.* Fr.

Vac-

Vaccinium foliis perennantibus obverse ovatis integerrimis, revolutis, subtus punctatis. HALL. *H. n.* 1. *Vaccinium Vitis Idaea*. LINN. *Sp.* 10. *Vitis Idaea foliis subrotundis non crenatis, baccis rubris*. B. *Pin.* 470. 1. R. H. Vbique in sylva hercynica. Fr.

Vaccinium foliis ovalibus integerrimis, basi ciliatis deciduis. *uliginosum*. LINN. *Sp.* 3. *Vaccinium foliis annuis exalbidis*. HALL. *Helv.* 2. *Vitis Idaea* II. CLVS. *Hist.* l. p. 61. T. 62. In sylva Hercynica. Fr.

Vaccinium caule angulato, foliis serratis annuis. LINN. *Cliff.* 148. HALL. *H. n.* 3. *Vaccinium Myrtillus* LINN. *Spec.* 1. BLACKW. *T.* 463. *Vitis Idaea foliis oblongis crenatis, fructu nigro*. B. *Pin.* 470. *Myrtillus* MATH. p. 331. In sylvis. Fr.

ARBUTVS. LINN. 488. TOURN. 368. *Vua vrsi*. T. 370.

Priori similis, sed flos quinquefidus, & fructus intra florem.

H. *Arbutus* caule erecto, foliis glabris serratis, baccis polyspermis, tuberculis seminum exasperatis. *Vnedo*. LINN. *Sp.* 1. *Arbutus folio serrato*. B. *Pin.* 460. *Arbutus*. CLVS. *Hisp.* p. 77. In Europa australi. Fr.

Arbutus caulibus procumbentibus, foliis integerrimis. *Vua vrsi*. LINN. *Sp.* 4. HALL. *H. n.* 2. *Vua vrsi* CLVS. *Hist.* l. p. 63. TABERN. p. 1080. 1. R. H. In *Cellensi* ducatu. Fr.



N 3

ANDRO-

na quinque medio flori sub segmentis inserta. Stylus stigmatibus trifido fimbriato, extra florem pendulo. Folia utrinque ovata, subintegra, versus apicem acute serrata. Fr.

PYROLA, LINN. 490.

Petala quinque, concava. Stamina decem. Capsula pentagona.

Pyrola foliis ovatis acuminatis, serratis, spica unilaterali. HALL. *H. n.* 1. *Pyr. secunda.* LINN. *Sp.* 3. *Pyrola folio mucronato serrato.* B. *Pin.* 181. *Pyrola II, tenerior* CLVS. *Pann.* 506. In sylvis passim.

Pyrola racemosa, staminibus & pistillo rectis. HALL. *Gott.* 165. *Helv. n.* 2. *minor.* LINN. *Sp.* 2. *Pyrola minor.* MILL. *Cat. Giesf.* p. 94. In sylvis *Hercynicis* circa *Ilfeld* & alibi.

Pyrola staminibus adscendentibus, pistillo declinato, rotundifolia. LINN. *Sp.* 1. HALL. *H. n.* 3. *Pyrola rotundifolia maior.* B. *Pin.* 191. In sylvis circa *Eichenkrug, Hardeggen.*

Pyrola scapo uniflora. uniflora. LINN. *Sp.* 6. HALL. *H. n.* 4. *Pyrola flore singulari.* RIV. *Pentap. irreg.* T. 35. *Pyrola IV.* CLVS. *Pann.* p. 508. 509.

TAMARIX, LINN. 337. Tamariscus, HALL.

TOURN.

Petala quinque. Stamina decem, aut quinque. Stigmata tria plumosa. Capsula triquetra, unilocularis.

H. *Tamarix floribus decandris. germanica.* LINN. *Sp.* 2. HALL. *H. p.* 19. *Tamariscus fruticosa, fo-*

lio

lio crassiflore f. Germanica. B. Pin. 485. LOB. ic. 218.
BLACKW. T. 331. Ad Rhenum, Danubium aliosque
fluvios.

OLIGOSTEMONES TRIHILATAE.

LINN.

Calyx quinquepartitus, aut quinquefolius. Petala quinque saepe ex calyce orta. Stamina vario numero quinque, septem, octo. Germen saepe nectarium cingit. Stylus simplex. Fructus bi-aut trilocularis alatus, echinatus, angulatus.

TROPAEOLVM. LINN. 421. Cardamin-
dum. TOURN. 244.

Calyx monophyllus, pone nectario conico instructus. Petala calyci inserta inaequalia. Stamina octo declinata. Stylus stigmatibus trifido. Capsulae tres angulatae congestae.

[†]*Tropaeolum foliis integris, petalis acuminato-se-*
taceis. minus. LINN. *Sp.* 1. *Cardamin-*
dum minus
& vulgare. FEVILL. *Peruv.* 3. p. 14. T. 8. *Nastur-*
tium indicum maius. B. Pin. 306. LOB. ic. 166.
f. 1. 2.

ÆSCVLVS. LINN. 420. Hippocastanum.
TOURN. 382. Pavia. BOERH. 260.

Petala calyci inserta. Capsula echinata trilocularis. Stylus simplex. Flores racemosi.

H. *Æsculus* floribus heptandris. *Hippocastanum.*
LINN. *Vps.* 92. *Cliff.* 142. *Castania equina.* OLIV.
Hist.

Hist. I. p. 7. Exotica quidem, etsi nunc fere spontanea ad vias & in hortis. Folia digitata. Fr.

H. Esculus floribus octandris. *Pavia*. LINN. Sp. 2. *Pavia*. BOERH. *Lugd. bat.* 2. T. 260. *Cliff.* 143. In *Carolina*. Fr.

ACER. LINN. 1023.

Calyx coloratus. Petala quinque & stamina octo ex circulo tumore balsamico, cui germen immersum. Stilus stigmatibus duplici. Capsulae duae alis maximis auctae. Sexus varius.

Acer foliis semitrilobis, lobis obtuse incisis. HALL. H. n. 1. Acer foliis lobatis obtuse emarginatis. (*campestre*.) LINN. Spec. 7. Acer *campestre* & minus. B. Pin. 431. Acer minus *Hist.* I. p. 8. 9. Vbiq. ad sepes. Fr.

Acer foliis quinquangulis, vndique serratis. HALL. H. 2. Acer foliis quinquelobis inaequaliter serratis, floribus racemosis. (*Pseudo-platanus*.) LINN. Spec. 2. Acer montanum candidum. B. Pin. 430. I. R. H. Acer latifolium. GLVS. *Hist.* I. p. 10. In sylvis. Fr.

Acer foliis quinquangulis, acutissime & rariter dentatis. HALL. H. n. 3. Gott. 168. Acer foliis quinquelobis acuminatis acute dentatis glabris, floribus corymbosis. (*Platanoides*.) LINN. Sp. 5. Acer maior. CAMER. Ept. 63. In M. Hengst. et *Cliff.*

CARDIOSPERMVM. LINN. 447.

Calyx trifolius. Petala quatuor. Nectarium quadrifolium circa germen. Stamina octo. Capsula inflata triloba.

OLIGOSTEMONES

H. *Cardiospermum. Halicacabum*. LINN. *Sp.* 1.
Vesicaria. RIV. *Tetrap. irreg.* T. 144. *Corindum*
ampliori folio, fructu maiore. 1. R. H. 431. In
India. A.

STAPHYLAEA. LINN. 336. *Staphylodendron*.
 TOVRN. 386.

Calyx coloratus. Petala quinque. Circulus balsa-
micus circa germen. Capsulae tres inflatae, con-
natae.

Staphylaea foliis pinnatis. pinnata. LINN. *Sp.* 1.
 HALL. p. 423. *Staphylodendron.* TABERN. p. 1022.
 CAMER. *Epit.* 171. 1. R. H. In hortis & ad sepes
 fere spontanea. Fr.

BERBERIS. LINN. 399. TOVRN. 385.

Petala sex, ad basin binis glandulis instructa. Bac-
ca disperma.

Berberis pedunculis racemosis. vulgaris. LINN. *Sp.*
 1. *Berberis spinis triplicibus.* CLIFF. 122. HALL.
 H. p. 242. *Berberis.* BLACKW. T. 165. *dumetorum.*
 B. Pin. 454. 1. R. H. In dumetis. Fr.

TELEPHIUM. LINN. 339. TOVRN. 128.

Petala quinque calyci aequalia. Capsula triquetra.

H. *Telephium*. LINN. *Sp.* p. 271. *Telephium*
Imperati. *Dioscoridis Imperati. Telephium legitimum.* GLVS.
Hist. II. p. 67. In Gallo-Provincia. P.

TRAPA.

TRAPA. LINN. 246. Tribuloides.

TOVRN. 431.

Calyx quadripartitus. Petala quatuor. Capsula ossea spinis quatuor crassis, ex calyce mutato enatis, armata.

Trapa natans. LINN. *Sp. p.* 120. *Trapa perfoliatarum natantium ventricosis*. CLIFF. 483. *Tribuloides* HALL. *Helv.* 468. *Tribulus aquaticus* CAMER. *epit. p.* 715. & aliorum. In stagnis. P.

PTELEAE *similis*.

Frutex *Pteleae* affinis, foliis ternatis, crenatis, fructu trihilato. Ita interim vocare liceat fruticem, qui noui generis nomen forte meretur. Frutex humilis vix bipedalis, ramis patulis. Folia ternata utrinque lanceolata, medio maiori. Flores cymosi. Calycis exigui foliola quinque angusta. Petala quinque ovato lanceolata, cava, patentia, calyce multo maiora, ex albo viridia. Stamina quinque, filamentis basi crassioribus, hirsutis. Germen intra corollam tetraedrum, stylo brevi, stigmate quadrifido, abit in capsulam quadrilocularem, quatuor extantibus latis alis auctam. Saepe calycis folia, petala & stamina quatuor; stigmata tria, capsulae alae & loculi tres. Vides affinitatem & diversitatem a *Ptelea*. *Ptelea trifoliata* L. *f. p.* 173.

ISOSTEMONES CYMOSAE. LINN.

Flos fructui innatus. Calycis segmenta, petala aut corollae segmenta, & stamina quinque. Stilus brevis stigmate tumido. Fructus pulposus, aut bacca. Flores in cymas & racemos congesti.

EVO-

EVONYMVS. LINN. 240.

Petala & stamina ex placenta communi. Stigma obtusum. Capsula pulposa, quadrangularis, quadrilocularis.

Evonymus foliis oblongo-ovatis. HALL. H. 523.
n. 1. Evonymus floribus plerisque quadrifidis. (euro-
paeus) LINN. Sp. 1. EVON. vulgaris, gravis rubentibus.
B. Pin. 428. GAMER. Epit. 102. Ad sepes. Fr.

CELASTRVS. LINN. 239. Evonymoides.

Fabrica priori similis: sed stigma trifidum, & capsula trigona.

H. Celastrus spinis nudis, ramis teretibus, foliis acutis. pyracanthus. LINN. Sp. 5. Lycium aethiopicum, pyracanthae folio. COMM. H. l. p. 163, T. 84. In Africa. Fr.

RHVS. LINN. 331. Toxicodendron.

Cotinus. TOURN.

Calyx erectus, petala erecta. Stamina brevissima. Stigmata tria sessilia. Capsula monosperma.

H. Rhus foliis pinnatis, obtusiuscule serratis ovalibus, subtus villosis. Coriaria. LINN. Sp. 1. Rhus coriariorum. LOB. ic. p. 11. p. 98. DOD. Pempt. 779. Sumach vulgo. In Europa australi. Fr.

H. Rhus foliis ternatis, foliolis petiolatis ovatis nudis integerrimis, caule radicante. radicans. LINN. Sp. 6. Hedera trifolia Canadensis. CORNV. Canad. T. 97. In Virginia. Canada. Fr.

H. Rhus

H. *Rhus* foliis simplicibus obovatis. *Cotinus*. LINN. *Sp.* 12. *Coccygia*. *Theophrasti*. LOB. *ic.* II. p. 99. *Cotinus coriaria*. DOD. *Pempt.* 780. In *Austria*. *Italia*, *Servia*. Fr.

VIBURNVM. LINN. 332. *Tinus*. *Opulus*.

Calyx exiguus. Petalum campanulatum. Stigmata & bacca, vt in priori.

H. *Viburnum* foliis integerrimis ovatis, ramificationibus subtus villoso-glandulosis. *Tinus*. LINN. *Sp.* 1. *Tinus*. II. CLVS. *Hist.* I. p. 49. *Laurus Tinus*, *f. sylvestris*, I. B. *Hist.* III. p. 418. In *Europa australi*. Fr.

H. *Viburnum* foliis cordatis, crenatis venosis, subtus tomentosis. *Lantana*. LINN. *Sp.* 5. HALL. *H.* p. 462. *Viburnum vulgo*. B. *Pin.* 249. I. R. H. *Lantana* DOD. *Pempt.* 701. In *Thuringia*. *Europa australi*. Fr.

Viburnum foliis lobatis, petiolis glandulosis. LINN. *Spec.* 7. *Opulus* I. R. H. LINN. *Cliff.* 109. HALL. *H.* p. 463. *Sambucus aquatica*. B. *Pin.* 456. Flores in radio cymae steriles, petali segmento exteriori maximo.

β. Varietas floribus omnibus amplioribus, sterilibus, in cymas globosas congestis. *Opulus flore globoso*. I. R. H. *Sambucus arbor rosea*.

SAMBVCVS. LINN. 334.

Petalum rotatum, bacca di- & trisperma.

Sambucus caule arboreo, floribus umbellatis.
 HALL. H. n. 1. Sambucus *nigra*, LINN. Spec. 3.
 BLACKW. T. 151. Sambucus *fructu in umbella nigro*, B. Pin. 456. Ad sepes. Fr.

Sambucus caule herbaceo, floribus umbellatis.
 HALL. H. n. 2. Sambucus *Ebulus*, LINN. Sp. 1.
 BLACKW. T. 488. Sambucus *humilis* f. *Ebulus*.
 B. Pin. 456. Rarius in gramineis & hortis. P.

Sambucus caule arboreo, floribus racemosis.
 HALL. H. n. 3. ^{racemosa} LINN. Sp. 4. Sambucus *racemosa*
rubra, BAVH. Pin. 456. In sylvis v. g. Eulenloch &
 versus Besigebausen. Fr.

CASSINE. LINN. 333.

Corolla patens. Bacca trilocularis stigmatibus
 umbilicata.

H. Cassine foliis subovatis integerrimis. *Mauroce-*
nia, LINN. Sp. 2. Frangula *sempervirens*, folio
rigido subrotundo, DILL. Elb. 146. T. 121. f. 147.
Cerasus Capensis, fructu rubro, folio fere obtuso.
 PETIV. Gaz. T. 57. f. 4. In Aethiopia. Fr.

CAPRIFOLIUM. HALL. 464. Lonicera.

LINN. 210.

Petalum ex longo tubo bilabiatum, in aliis fere
 aequaliter quinquefidum. Stilus stigmatibus tumido.
 Bacca trilocularis.

H. Caprifolium foliis supremis coalescentibus,
 floribus bilabiatis. HALL. Gott. 192. Lonicera *flo-*
ribus verticillatis terminalibus sessilibus, foliis *sum-*
mis connato-persolatis, LINN. Sp. 1. Periclyme-
 num

num *perfoliatum*. B. Pin. 302. LOBEL. ic. 652.
Caprifolium Italicum. DOD. Pempt. 411. RIV. T.
 122. In Europa australi. Fr.

H. *Caprifolium* foliis connatis, floribus tubulo-
 sis. HALL. l. c. *Lonicera spicis nudis verticillatis*
terminalibus, foliis summis connato-perfoliatis.
 LINN. Spec. 2. *Periclymenum perfoliatum Virgi-*
nianum sempervirens & florens. HERM. Lugdb.
 484. T. 483. *Virginianum*. RIV. Monop. irreg.
 T. 116. Fr.

Caprifolium floribus racemosis, foliis distinctis.
 HALL. H. n. 5. *Lonicera capitulis ovatis imbricatis*
terminalibus, foliis omnibus distinctis. LINN. Sp. 3.
Periclymenum non perfoliatum Germanicum. B. Pin.
 302. In sepibus & sylvis. v. g. versus Seeburg, Offe-
 rode. Fr.

II. *Caprifolium* fructu gemino rubro, foliis gla-
 bris, cordatis. HALL. Gott. l. c. AMM. Ruth. n. 262.
Lonicera pedunculis bifloris, baccis distinctis, fo-
liis cordatis obtusis. (Tatarica.) LINN. Sp. 5. In
 Sibiria. Fr.

Caprifolium floribus geminis, foliis ovatis, la-
 nuginosis, integerrimis. HALL. Helv. n. 1. *Lonicera*
pedunculis bifloris, baccis distinctis, foliis integer-
rimis pubescentibus. (*Xylosteum*.) LINN. Sp. 6.
Chamaecerasus dumetorum, fructu gemino rubro.
 B. Pin. 451. In sepibus. Fr.

H. *Caprifolium* floribus geminis, foliis ovatis,
 nervosis, ciliatis. HALL. H. n. 3. *Lonicera pedun-*
culis bifloris, baccis coadunatis globosis, stylis indi-
 visis.

^{cacumlea}
vifis. LINN. *Sp.* 9. *Periclymenum rectum, fructu*
coeruleo. CLVS. *Pann.* p. 85. *Fructus sicut fructus Lonicerae.*

DIERVILLA. LINN. ed. pr.

Calyx persistens, segmentis longis, subulatis. Capsula ovata, bilocularis.

Diervilla LINN. *Cliff.* 63. T. 7. Diervilla *Aca-*
diensis fruticosa, flore luteo. TOURN. *Act. Par.*
 1706. T. 7. f. 1. *Lonicera racemis terminalibus,*
foliis serratis. LINN. *Sp.* 14. In *America septen-*
trionali.

VMBELLIFERAE.

Caulis fistulosus. Folia alterna plerumque com-
 posita. Petioli ex eodem puncto enati flores in
 umbellis proferunt. Umbellis foliola subiecta invo-
 lucri nomine veniunt. Semina duo ad mediam axin
 sibi applicata in apice communem gerunt placen-
 tam, ex qua enascuntur, in paucissimis calyx exi-
 guus, in omnibus quinque petala, totidem stami-
 na, styli duo simplices.

I. Umbella obscuriori.

PHYLLIS. LINN. 286.

Calyx bifolius minimus. Fructus turbinato-ob-
 longus, angulatus.

h. *Phyllis stipulis dentatis*. LINN. *Sp.* 1. *Vale-*
rianella Canariensis frutescens simpla nobis dicta.
 DILL. *Eltb.* 405. T. 299. f. 386. In *Canariis.*

HYDROCOTYLE, LINN. 288.

Vmbella simplex. Involucrum quadrifolium. Fructus orbiculatus.

Hydrocotyle foliis peltatis orbiculatis, vndique emarginatis. LINN. *Cliff.* 88. HALL. *H.* 425. *vulgaris.* LINN. *Sp.* 1. 1. R. H. *Cotyledon aquatica.* BOB. *ic.* 387. *Jater Engles et Cila vasa P. Martius ne Martius*
Oct. 76. D. 76. 90.

II. *Vmbella manifesta.*

Seminibus gibbis striatis, involucri nullo.

ANETHVM, LINN. 326. TOURN. 169.

Foeniculum. T. 164. HALL. 425.

Vmbella multiplex; petala aequalia. Foliorum ultimae pinnulae tenuissimae.

H. *Anethum* fructibus compressis. *graveolens.* LINN. *Sp.* 1. *Mat. med.* 146. *Anethum hortense.* B. *Pin.* 147. Inter *Hispaniae* segetes.

H. *Anethum* fructibus ovatis. *Foeniculum.* LINN. *Sp.* 2. *Foeniculum vulgare Germanicum.* B. *Pin.* 147. HALL. *H.* 425. In *Helvetia;* *Gallia Narbonensi.*

APIVM. LINN. 329.

Vmbella vniversalis radiis paucioribus; peculiaris pluribus. Petala aequalia.

Apium foliolis caulinis cuneiformibus semitrifidis, ora loborum circumferrata. *graveolens.* LINN. *Sp.* 2. HALL. *H.* 427. *Apium palustre seu officinarum.* B. *Pin.* 154. *Apium* BLACKW. *T.* 443. β . *Apium*

dulce, *Celleri italorum*. I. R. H. 305. α . in fossis
Europae. β . in hortis culta.

Apium Abrusianum.

SMYRNIVM. LINN. 325.

Umbella vniversalis inaequalis, partialis erecta,
flores subaequales.

H. *Smyrnium* foliis caulinis ternatis petiolatis ser-
ratis. *Olusatrum*. LINN. Sp. 2. *Smyrnium Diosc.*
B. Pin. 154. Ad rupes Angliae & Galliae.

AEGOPODIVM. LINN. 330. Podagraria.

HALL. 427.

Umbella multiplex, petala inaequalia, semina
magna.

Aegopodium foliis caulinis summis ternatis.
LINN. Sp. p. 265. Podagraria. HALL. l. c. RIV. T.
46. *Angelica sylvestris minor*. B. Pin. 115.

PASTINACA. LINN. 324.

Petala aequalia flava. Fructus compresso-planus
margine cinctus.

Pastinaca foliis simpliciter pinnatis. ^{*latius*} LINN. Sp. 1.
HALL. H. 446. *Pastinaca sylvestris latifolia*. B. Pin.
155. In gramineis.

CARVM. LINN. 327. Carvi. T. 160.

Petala subaequalia. Semina ovato-oblonga.

^{*Carui*}
Carum. LINN. Sp. 263. Mat. med. 138. HALL.
H. 428. CAMER. Epit. § 16. *Carum pratense*, *Car-*
vi officinarum. B. Pin. 158. Pinnulae circa costam se
decussant. In pratis.

PIMPINELLA. LINN. 328. Tragofelinum. T.

Vmbella laxa. Petala subaequalia.

Pimpinella foliis pinnatis: foliolis radicalibus subrotundis; summis linearibus. *saxifraga*. LINN. Sp. 1. Tragofelinum pinnis semibilobis, circumscissatis. HALL. H. n. 1. Pimpinella saxifraga maior. B. Pin. 264. I.

β. Pimpinella (*Tragofelinum*) pinnis laciniatis, superioribus tenuissime divisis. HALL. H. n. 3. Pimpinella saxifraga minor. B. Pin. 150. In locis ficcis. Num varietas prioris.

h. Pimpinella foliis radicalibus trifidis incis. *Anisum*. LINN. Sp. 4. Anisum vulgare. CLVS. Hist. 2. p. 202. BLACKW. T. 374. Anisum herbariis. B. Pin. 159. In Aegypto. A.

β. Seminibus gibbis striatis, involucro alterutro aut utroque.

SESELI. HALL. LINN.

h. Sefeli foliis glaucis, foliolis longe linearibus, radice praelonga, seminibus subhirsutis. HALL. Comm. Soc. Gott. T. II. p. 347. vbi vide fusioem descriptionem. GMEL. Fl. Sibir. T. I. p. 207.

OENANTHE. LINN. 314.

Vmbella difformis. Flores radii mares inaequales, maiores, disci androgyni aequales. Fructus calyce coronatus.

Oenanthe stolonifera, foliis caulinis pinnatis fistulosis teretibus. *fistulosa*. LINN. *Sp.* 1. HALL. *H.* 432. n. 1. *Oenanthe aquatica*. B. *Pin.* 162. RIVIN. 66. in fossis & pratis v. g. versus *Maschmüble*. P.

H. *Oenanthe umbellularum* pedunculis marginalibus longioribus ramosis masculis. *prolifera*. LINN. *Sp.* 3. *Oenanthe prolifera apula*. B. *Pin.* 163. MORIS. *Hist.* III. p. 289. S. 9. T. 7. f. 5. A.

H. *Oenanthe* fructibus globosis. *globulosa*. LINN. *Sp.* 4. *Oenanthe Lusitanica*, semine crassiori globoso. I. R. H. 313. P.

H. *Oenanthe* foliolis radicalibus ovatis incis, caulibus integris linearibus longissimis simplicioribus. *pimpinelloides*. LINN. *Sp.* 5. *Oenanthe apii folio*. B. *Pin.* 162.

PHELLANDRIVM. LINN. 315.

Flores inaequales. Fructus ovatus, fere glaber, calyce coronatus.

Phellandrium foliorum ramificationibus divaricatis. LINN. *Sp.* 1. *Phellandrium foliis ad angulos obtusos ramosis*. HALL. *H.* p. 432. RIV. T. 64. In fossis.

CORIANDRVM. LINN. 318.

Flores radii maiores inaequales androgyni, fructu sphaerico calyce coronato. Flores disci mares minores.

H. *Coriandrum* fructibus globosis. *sativum*. LINN. *Sp.* 1. *Mat. med.* 135. *Coriandrum maius*. B. *Pin.* 158. RIV. T. 69. 70. In Italia. A.

H. Coriandrum fructibus didymis. *testiculatum*.
 LINN. Sp. 2. Coriandrum *sylvestre foetidissimum*. B.
 Pin. 158. Prope *Vnam*. A.

AETHVSA. LINN. 317. Cynapium. RIV.

Involucrum parziale trifolium, laterale, reflexum. Petala inaequalia.

Aethusa Cynapium. LINN. Sp. p. 256. *Cicutaria fatua*. LOBEL. ic. 280. Cynapium. RIV. T. 75. In hortis.

AMMI. LINN. 297.

Involucrum vniuersale pinnatifidum.

H. Ammi foliis inferioribus pinnatis lanceolatis serratis, superioribus multifidis linearibus. LINN. Sp. 1. Ammi *maius*. B. Pin. 159. Ammioselinum. RIV. T. 84. In *Europa* australi. A.

CONIVM. LINN. 299. Cicuta.

TOVRN.

Involucri vtriusque foliola lata. Fructus subrotundus striatus.

Conium seminibus striatis. *maculatum*. LINN. Sp. 1. Cicuta. BLACKW. T. 451. HALL. H. p. 434. Cicuta *maior*. B. Pin. 160. *vulgaris*. GLVS. Hist. II. p. 200. In ruderatis, ad sepes.

SIVM. HALL. 435. LINN. 310. *Cicuta*.

LINN. 316.

Involucrum vniuersale latifolium reflexum. Petala aequalia.

Sium foliis pinnatis, serratis, umbella terminali. latifolium. LINN. *Sp.* 1. HALL. *H. n.* 1. C. B. *Pin.* 154. I. R. H. *Sium*. DODON. *Pempt.* 589. RIV. T. 77. In fossis ad loca salsa. v. g. prope *Harste*; & *Salz der Helden*.

Sium foliis pinnatis, umbellis axillaribus sessilibus, nodiflorum. LINN. *Sp.* 2. *Sium foliis pinnatis, inferioribus simplicibus, superioribus trifidis, argente dissectis*. HALL. *H.* 2. *Sium aquaticum ad alas floridum*. F. R. H. In fossis.

H. *Sium* foliis pinnatis, floralibus ternatis. Sisarum. LINN. *Spec.* 3. *Sisarum Germanorum*. B. *Pin.* 155. *Sisarum*. RIV. T. 77. A.

Sium pinnis multifidis, serratis, nervo folioso. HALL. *H.* 6. *Sium foliolis linearibus decurrentibus connatis*. LINN. *Spec.* 6. Falcaria. RIV. T. 47. In agris super *Weende*.

Sium foliis serratis, *Cicuta* pinnulis tripartitis. HALL. *H. n.* 5. *Cicuta* umbellis folio oppositis, petiolis marginatis obtusis. (*virosa*.) LINN. *Sp.* 1. *Cicuta aquatica*. WEPFER in proprio libro T. 1-4. LINN. *Lapp.* 103. In lacu ad *Seeburg* & *Denckershausen*.

SISON.

Sison foliis ternatis. Canadense. LINN. *Sp.* 3. HALL. *Gott.* 175. *Myrrhis Canadensis trilobata*. MORIS. *Hist.* III. p. 301. S. 9. T. II. f. 4. Huic speciei involucrum utrumque deest.

LIGVSTICVM. LINN. 308.

H. Ligusticum foliis multiplicibus, foliolis superne incis. Levisticum. LINN. *Sp.* 1. *Mat. med.*

119. *Ligusticum vulgare*. B. Pin. 157. In montibus *Liguriae*. P.

H. *Ligusticum foliis biternatis*. *Scoticum*. LINN. Sp. 2. *Ligusticum humilius Scoticum*, apii folio. J. R. H. PLVKN. Alm. 217. T. 96. f. 2. *Seseli maritimum Scoticum*. HERM. Parad. T. 227. RIV. T. 68. In *Scotia*.

ATHAMANTA. LINN. 301. Meum

TOURN. 165.

Athamanta foliolis capillaribus, seminibus glabris striatis. *Meum*. LINN. Sp. 8. *Meum*. HALL. p. 426. *Meum vulgare*, foliis *Anethi*. B. Pin. 148. In sylva *Hercynica* prope *Andreasberg* & *Altenau*.

Athamanta Libanotis. p. 226. 227. | *Athamanta Cuneata* p. 221.

BUPLEVRVM. LINN. 291. HALL. 436.

Involucrum univcrsale interdum deest: partiale quinquefolium aut quinquesidum. Semina profunde sulcata. Flores lutei. Folia simplicia.

Bupleurum foliis ovatis perfoliatis. HALL. H. 437. *Bupleurum involucris universalibus nullis*, foliis perfoliatis. LINN. Sp. 1. *Perfoliata vulgarissima arvensis*. B. Pin. 277. BLACKW. T. 95. Ubique inter segetes.

Bupleurum foliis imis petiolatis, superioribus ovatis perfoliatis. HALL. H. n. 2. *Bupleurum longifolium*. LINN. Sp. 5. *Perfoliata montana latifolia*. B. Pin. 277. In sylva prope arcem *Plessi*.

Bupleurum foliis petiolatis falcato-ellipticis acutis. HALL. H. n. 11. *Bupleurum falcatum*. LINN. Sp. 6. *Bupleurum longis & angustis foliis incurvis*

H.

H. Ox. III. p. 300. *Über der Papiermühle & in M. Altenstollberg.* ad res aprem inter stanz burg et 901166

H. Bupleurum foliis gramineis, calyce peculiari quinquesido aristato rigido. *HALL. H. n. 6. Bupleurum odontites. LINN. Sp. 7. A*

CRITHMUM. LINN. 303.

H. Crithmum foliolis lanceolatis carnosiss. *maritimum. LINN. Sp. 1. Crithmum. f. Foeniculum marinum. B. Pin. 288. LOB. ic. 392. Ad littora maris.*

Seminibus, gibbis, striatis, margine alatis.

PEUCEDANUM. LINN. 302.

H. Peucedanum foliis quinquies tripartitis filiformibus linearibus. *officinale. LINN. Sp. 1. Mat. med. 126. Peucedanum germanicum. B. Pin. 149. In Germania australi.*

Peucedanum foliis ^{v. laur.} pinnatifidis, foliolis longitridis oppositis. *LINN. Sp. 3. Sefeli pratense, Silaus forte Plinio. B. Pin. 162. Siler pratense. DOD. Pempt. 310. Super Wernigerode.*

Peucedanum foliis duplicato pinnatis, pinnulis longis linearibus, dissitis. An Carvisfolia. 1. B. III. p. 171. Descripsi hanc plantam in *epistola* ad III. WERLHOFFIUM *Observ. bot. p. 12. Caulis striatus. Petioli foliorum membranacei longe amplexicaules. Umbella utraque densissima. Involucrum utrumque decaphyllum, radiis appressum, iisque brevius. Petala alba, extus flava. Semina convexa striata, alacincta. Circa Berolinum in sylvis vliginosis.*

SELINUM. HALL. 443. LINN. 300.

Selinum sublaetescens, radice vnica, *palustre*. LINN. *Sp.* 2. HALL. *H.* 3. *Thyselinum angustifolium*. RIV. *T.* 20. *Seseli palustre laetescens*. B. *Pin.* 162. *Prodr.* 85. *Im Landolfsbäuser Sumpf*. Circa *Berolinum*.

Selinum pinnis pinnatis, pinnulis appendiculatis, circumferratis, triangularibus. HALL. *Helv.* n. 4. *Oreoselinum apii folio*. In praecipitiis M. *Plesberg*.

Selinum caule alato, foliolis bi- & trifidis. *Angelica foliis duplicato pinnatis, foliolis angustis, saepe trifidis*. BOEHM. *Lips.* 197. *An Peucedanum*. HALL. 3. *Angelica tenuifolia*. RIV. *T.* 13. DILL. *Nov. gen.* p. 226. Caulis inferius alis membranaceis notatus. Foliola apice alba. Involucrum univertale nullum aut unifolium, parziale multifolium lineare, petiolis brevius. Semen tribus sulcis profunde exaratum, margine alatum. *Selinum Carn. folio L.*
Cf. Hall. *Hist. Nat. Helv.* 1. p. 228. Tab. 20. *Prodr. Hist. Nat. Helv.* 1. p. 42.

ANGELICA. LINN. 309.

H. *Angelica foliorum impari lobato*. *Archangelica*. LINN. *Spec.* 1. *Mat. med.* 120. *Angelica sativa*. B. *Pin.* 155. *Archangelica*. CLVS. *Hist.* 2. p. 295. *Vulgaris in hortis*. Sponte in alpi-
bus Lapponicis.

Angelica foliis aequalibus ovato-lanceolatis serratis. *sylvestris*. LINN. *Sp.* 2. HALL. *H.* 445. *Angelica sylvestris maior*. BAVH. *Pin.* 155. *Vbiq. in pratis*.

H. An-

H. *Angelica* foliolis aequalibus, ovatis, inciso-serratis. LINN. *Sp.* 4. HALL. *Gatt.* 179. *Angelica lucida Canadensis*. CORNV. *Canad.* 196. T. 197. In *Canada*.

IMPERATORIA. LINN. 321.

H. *Imperatoria*. *Ostruthium*. LINN. *Spec.* 259. RIV. T. 7. LOB. *ic. p.* 700. BLACKW. T. 279. B. *Pin.* 156. *Magistrantia*. CAMER. *Epit.* 592. In al-
pibus *Helvetiis*.

Seminibus quatuor & pluribus alis armatis.

ASTRANTIA. LINN. 290. HALL. 439.

Umbella simplex, involucro aequalis. Flores
mares & androgyni. Semen calyce coronatum.

Astrantia foliis quinquelobis, lobis trifidis. *ma-
ior*. HALL. H. 439. *Astrantia*. RIV. T. 67. *Astran-
tia maior, corona floris purpurascens*. J. R. H. In
M. *Altenstollberg*. P.

LASERPITIVM. LINN. 306.

HALL. 440.

Laserpitium foliolis cordatis, inciso-serratis. *lati-
folium*. LINN. *Sp.* 1. HALL. H. n. 1. *Libanotis
latifolia maior*. C. B. *Pin.* 159. *Seseli Aethiopicum
herba*. DOD. *Pempt.* 312. In M. *Altenstollberg*, circa
Elbingerode, super *Wernigerode*. P.

H. *Laserpitium* foliolis lanceolatis integerrimis
petiolatis. *Siler*. LINN. *Sp.* 6. *Ligusticum, quod
Seseli officinarum*. B. *Pin.* 162. *Siler montanum*.
DOD. *Pempt.* 310. CLVS. *Hist.* CXCV. *Seseli*
BLACKW. T. 426. In al-
pibus *Helvetiae*. P.

W. H. Laserpitium foliis lanceolatis integerrimis, optimis coactis. *Prutenicum*. LINN. Sp. 1. *Laserpitium foliis radicalibus patentissimis*, BAYEN. 103. *Laserpitium daucoides Prutenicum, viscoso semine*. BREYN. Cent. 167. T. 84. In *Marchia, Borussia*. Circa *Jenam*.

Seminibus planis margine cinctis.

SPHONDYLIVM. HALL. 447.

LINN. 307.

Sphondylium pinhatum, pinnis hirsutis semel quinquefidis. HALL. H. n. 1. *Heracleum Sphondylium*. LINN. Sp. 1. *Sphondylium vulgare hirsutum*. BAYH. Pin. 157. Branca vrsina TABERN. p. 92. V. bique in pratis.

Sphondylium foliis angustioribus. RIV. T. 5. HALLERVS a priori distinguit. Enum. Gott. 248.

TORDYLIVM. LINN. 293.

Semina plana margine crenulato.

H. *Tordylium involucris umbella longioribus*. *Syriacum*. LINN. Sp. 1. *Tordylium Syriacum*. RIV. T. 3. *humilius, semine granulato maiore*. MORIS. Hist. III. p. 317. f. 9. T. 16. f. 7. In *Syria*. A.

H. *Tordylium umbellulis remotis, foliis pinnatis, pinnis subrotundis laciniatis*. *apulum*. LINN. Sp. 3. *Tordylium apulum minimum*. COLVMN. Ecpbr. I. p. 122. T. 124. *Seseli creticum minimum*, B. Pin. 161. In *Italia*.

H. Tor-

H. *Tordylium umbella conferta radiata*, foliolis lanceolatis inciso-ferratis. (*maximum*.) LINN. *Sp.*

5. *Tordylium*. § IV. *Pentap.* I.

~~Tordyl. latifolium~~ *Tordyl. Anthriscus* *Tordyl. nodosum*
Seminibus aculeatis. *Y. aspera*

CAUCALIS. HALL. 448. LINN. 294.

Corolla universalis radiata.

Caucalis involucris latifoliis, petalo extremo maximo. HALL. n. 1. *Caucalis involucris singulis pentaphyllis: foliolo unico duplo maiore. (grandiflora.)* LINN. *Sp.* 1. *Caucalis arvensis echinata*, magno flore. B. *Pin.* 152. *Echinophora pycnocarpus*. COLUMN. *Ecpbr.* l. p. 91. T. 94. Inter segetes circa Webnde. A.

H. *Caucalis* seminibus cylindricis asperis. HALL. G. 181. cum delc. *Anthriscus Creticus*. RIV. T. 38.

Caucalis involucro universalis nullo, laciniis foliorum perangustis. HALL. H. n. 2. *Caucalis seminum aculeis triglochidibus uncinatis; pilis verticillatis hispidis. (leptophylla.)* LINN. *Sp.* 6. *Caucalis arvensis echinata parvo flore & fructu*. B. *Pin.* 152. In agris.

Caucalis *umbella* universalis trifida, partialibus pentaspermis, foliis pinnatis, pinnis ferratis. LINN. *Cliff.* 91. ROY. *Lugdb.* 95. *Tordylium latifolium*. LINN. *Sp.* 5. *Caucalis arvensis echinata latifolia*. B. *Pin.* 152. In agris.

Caucalis seminibus hispidis, foliis hirsutis obtuse lobatis. HALL. *Gott.* 181. *Helv.* 449. *Tordylium Anthriscus*. LINN. *Sp.* 6. *Caucalis semine aspero, floscu-*

flosculis rubentibus. BAVH. Pin. 152. Prodr. 80. In
sylvis & ad sepes.) *Caucalis umbellata* Linn. *Sp. Pl.* 113. *Ep. humilis* et *maritima* *Prodr.* 80.

H. *Caucalis umbellis sessilibus simplicibus.* LINN.

Cliff. 91. HALL. Gott. 182. *Tordylium nodosum.*

LINN. Sp. 7. *Caucalis nodosa, echinato semine.*

B. Pin. 153. Prodr. 80. In Gallia, Italia. A.

Daucus
ARTEDIA, LINN. 295.

Semen in nostra alis profunde ferratis percur-
sum. Involucrum pinnatifidum.

Daucus muricata Linn. *Sp. Pl.* 5.

H. Artedia feminibus aculeatis, *muricata.* LINN.

Sp. 2. *Caucalis maior daucoides Tingitana.* HERM.

Parad. p. 111. cum ic. HALL. Gott. 182. *Echino-*

phora Tingitana. RIV. T. 26. A.

SANICVLA, LINN. 289.

Umbellae minimae globosae. Sexus Caucalidi si-
milis.

Sanicula foliis radicalibus simplicibus, flosculis
omnibus sessilibus. *Europaea.* LINN. Sp. 1. *Sani-*
cula RIV. T. 30. BLACKW. T. 63. *officinarum.*
B. Pin. 319.

Seminibus hirsutis,

DAUCVS, LINN. 296.

Involucrum pinnatifidum, umbella cava.

Daucus feminibus hispidis. *Carota.* LINN. Sp. 1.
HALL. H. 451. *Daucus vulgaris.* CLVS. *Hist.* II. p.
198. 1. R. H.

Chaerophyllum feminibus brevibus hispidis.
HALL. H. n. 1. Scandix ^{Tachylan} LINN. Sp. 4. Myrrhis syl-
vestris aequicolorum. COLUMN. Ecphr. I. p. 110.
T. 112. Caulalis folio Chaerifolii. RIV. T. 34.
Circa Hannoveram.

MYRRHIS. HALL. 453. Scandicis spec.

LINN. 319.

Semina striata.

Myrrhis radice lignosa, perenni, foliis hirsutis,
feminibus flavis, obscure striatis. HALL. Gott. 184.
Myrrhis perennis alba minor, foliis hirsutis, semi-
ne aureo. HALL. Rupp. Jen. 282. T. 5. Myrrhis
minor. LÖBEL. ^{Adrophyllo} Über Wernigerode.

Myrrhis perennis, feminibus strictis longissimis.
HALL. H. n. 4. Chaerophyllum ^{Scutellum} LINN. Sp. 4. Myr-
his Brockenbergenfis. THAL. p. 77. Myrrhis pala-
stris. RIV. T. 50. Cicutaria palustris latifolia. C. B.
Pin. 161. In pratis Hercynicis.

Myrrhis caule inferius hirsuto, superius glabro, ra-
dice verna globosa, ramosa. HALL. H. 3. Chaero-
phyllum ^{multiflorum} LINN. Sp. 2. Myrrhis tuberosa nodosa,
coniophyllum. I. R. H. Myrrhis foetens. RIV. T. 49.
Ad viam versus Edigebausen in dumetis.

Myrrhis pinnulis semipinnatis, obtusis, ordina-
tim decrefcentibus. HALL. H. 1. Chaerophyllum
caule maculato scabro, geniculis tumidis. ^{legumicentrum} LINN.
Sp. 3. Myrrhis annua, semine striato, levi. I. R. H.
315. BLACKW. T. 243.

H. Myrrhis feminibus profunde sulcatis, angula-
tis, maximis. Scandix odorata. LINN. Sp. 1. Odo-
rata

rata RIV. T. 56. Myrrhis maior odorata. B. Pin. 160. In Italia. P.

SCANDIX. HALL. 454.

Semina ventricoso-subulata.

Scandix foliis omnibus tenuissimis divisis. HALL. 1. c. Scandix seminibus laevibus rostro longissimo. LINN. Spec. 2. Scandix semine rostrato, vulgaris. B. Pin. 152. Pecten veneris. CAMER. epit. 302. LOB. ic. 726. In agris vbique.

H. Scandix seminibus subulatis hispidis, floribus radiatis, caulibus laevibus. australis. LINN. Sp. 5. Scandix cretica minor. B. Pin. 152. Anisomarathrum. COLUMN. Erphr. 1. p. 89. T. 90. In Gallia Narbonensi. Creta. A.

Scandix laevifolia; *Sc. Anthriscus*; *Sc. odorata*.^{p. 227.}

Seminibus receptaculo communi paleaceo insertis.

ERYNGIVM. LINN. 287. TOURN. 173.

Eryngium ramosissimum, pinnis foliorum pinnatis, alatis. HALL. H. n. 1. Eryngium foliis radicalibus pinnatis tripartitis. (campestre.) LINN. Sp. 7. Eryngium vulgare. B. Pin. 386. CLVS. Hist. II. p. 157. BLACKW. T. 297. Ad viam versus Bovenenden. P.

H. Eryngium foliis radicalibus ovalibus, planis, crenatis, floribus pedunculatis. planum LINN. Sp. 3. Eryngium Pannonicum latifolium. CLVS. Hist. II. p. 158. Ad Danubium. P.

H. Eryngium foliis radicalibus subrotundis plicatis spinosis, capitulis pedunculatis. LINN. Sp. 5.
Erya-

STELLATAE.

229

Eryngium maritimum. B. Pin. 386. CLVS. Hist. II. p. 169.

Dipetalae.

CIRCAEA. LINN. 24.

Petala & stamina duo fructui biloculari innata.

Circaea caule erecto, racemis pluribus. Lutesiana. LINN. Sp. 1. *Circaea foliis longioribus, obiter crenatis*. HALL. H. n. 1. *Solanifolia Circaea dicta maior*. B. Pin. 168. In umbrosis circa Plessi. Gleichen. Um die Quelle bey der Webender Papiermühle.

Circaea caule adscendente, racemo unico. alpina. LINN. Sp. 2. *Circaea foliis brevioribus dentatis*. HALL. H. 2. *Circaea minima*. COLUMN. Ephr. II. p. 80. In sylvis Hercynicis.

ISOSTEMONES STELLATAE.

Flos, vt in umbelliferis, duobus seminibus conatis impositus, sed corolla monopetala, limbo quadrifido, stamina quatuor, stylus stigmatibus bifido. Caulis quadrangularis, folia verticillata, vires adstringentes.

ASPERVLA. LINN. 113. *Sherardia*. LINN. 112. Rubeola. T. 50.

Flos nudus rubo cylindraceo.

(*Asperula* ^{*Sherardia*} *feminibus coronatis*. HALL. Gott. 186. *Sherardia* ^{*argentea*} *foliis omnibus verticillatis, floribus terminalibus*. LINN. Sp. 2. HALL. H. 457. Rubeola ar-

P 3

venfis

venfis repens coerulea. B. Pin. 334. Prodr. 145.

In agris vbique. A. *Asperula*.

H. *Asperula umbellis pilosis, foliis senis, longis & obtusis.* HALL. H. 457. *Asperula foliis senis, floribus sessilibus aggregatis terminalibus. (arvensis.)*

LINN. Sp. 2. *Asperula coerulea arvensis.* B. Pin. 334. In Anglia. Belgio. Helvetia. A.

Asperula foliis octonis, feminibus hispidis, ex summo caule paniculata. LINN. Sp. 1. *Asperula foliis octonis lanceolatis, florum fasciculis pedunculatis. (odorata.)* LINN. Sp. 1. *Asperula montana odorata.* B. Pin. l. c. In sylvis. P.

H. *Asperula foliis quaternis, ovato-lanceolatis, ramis alternis.* LINN. Cliff. 33. HALL. Helv. l. c. *floribus fasciculatis terminalibus. (Taurina.)* LINN. Sp. 3. *Rubia levis Taurinensium.* LOB. ic. p. 800. In Helvetia.

Asperula foliis quaternis linearibus, floribus paniculatis. HALL. G. l. c. H. 458. *Asperula cynanchica.* LINN. Sp. 6. *Rubia quadrifolia laevis, floribus purpurascens.* I. R. H. *Gallium mont. latifolium cruciatum.* COLUMN. Eschbr. l. p. 297.

H. *Asperula foliis plerisque binis, inferioribus ellipticis, superioribus linearibus, segmentis floris in apice glandula notatis. Plantula procumbens, diffusula. Flores extus rubent. An Asperula foliis quaternis lanceolato-linearibus, caule erecto, floribus saepe trifidis. Pyrenaica.* LINN. Sp. 5. *Rubia cynanchica saxatilis.* B. Pin. 133. Prodr. 146.? Descriptionem dedi in Comm. Gott. T. V.

Aspe-

STELLATAE.

231

Asperula foliis linearibus, inferioribus senis, intermediis quaternis, caule flaccido, floribus trifidis. HALL. G. 186. HALL. Rupp. Fen. p. 8. *tinctoria*. LINN. Sp. 4. *Galium album*. III. TABERN. T. 733. f. 1. In M. *Attenßollberg*.

CRUCIANELLA. LINN. 118.

Flos tubulosus, calyce trifolio cinctus. Segmenta apice inflexa.

H. *Crucianella* erecta, foliis senis linearibus. *angustifolia*. LINN. Sp. 1. *Asperula spicata, glumis floralibus maximis*. HALL. G. 187. *Rubeola angustifolia spicata*. B. Prodr. 145. Pin. 334. *Monspeli*.

H. *Crucianella* procumbens, foliis quaternis lanceolatis, floribus spicatis. *latifolia*. LINN. Sp. 2. *Rubia latifolia spicata*. B. Pin. 334. *Monspeli*.

GALIVM. HALL. 458. LINN. 117. Vailantia. LINN. 1019.

Corolla rotata sine tubo. Semina subrotunda.

Galium caule tereti, fulcrato, foliis octonis, glaucis & obtusis. HALL. H. v. 8. *Mollugo montana latifolia ramosa*. B. Pin. 334. *Mollugo* CLVS. Hist. p. CXLV. In sylvis. *Galium glutinosum* Lin p. 2 p. 156.

Galium foliis octonis, glabris, acuminatis. *Mollugo vulgarior*. LÖB. ic. 862. *Galium album vulgare*. I. R. H. Ad vias.

Galium caule recto, foliis senis, inferne canaliculatis. *Galium mont. altissimum, foliis angustis, albi*

albicantibus. RVPP. *Jen.* p. 5. In M. *Altenstollberg.*

Galium caule subaspero, foliis quaternis & senis, spinula terminatis. HALL. *H. n.* 1. *vliginsum.* LINN. *Sp.* 2. *Aparine palustris minor Parisien- sum, flore albo.* 1. R. H. 114. In vliginosis.

Galium foliis quaternis obovatis inaequalibus, caulibus diffusis. *palustre.* LINN. *Sp.* 2. HALL. *H. n.* 6. *Galium palustre album.* B. *Pin.* 335. In aquo- sis pratis.

Galium foliis senis & novenis, subasperis, spinu- la terminatis. HALL. *H. n.* 5. *Galium album minus.* VAILL. *T.* 78. Ad vias: locis lapidosis.

Galium caule ramosissimo procumbente, foliis quinis obverse ovatis. *saxatile.* LINN. *Sp.* 8. HALL. *H.* 460. *Galium saxatile supinum molliore folio.* JVSSIEV. *Atl. Par.* 1714. p. 490. Super *Hardegfen* & in sylvis *bercynicis* vbique.

Galium foliis octonis confertis, gracilissimis, spi- ca flava caulem terminante. HALL. *H. n.* 9. ^{verum} LINN. *Sp.* 9. *Galium luteum.* B. *Pin.* 335. BLACKW. *T.* 435. In arenosis. v. g. *beym Bovender Galgen.*

H. *Galium* procumbens, foliis senis, gracilissi- mis, confertis feminibus subhirsutis. HALL. *G.* 188. An *Gal. minutum* LINN. *Sp.* 9. In *Sibiria.*

Galium foliis quaternis lanceolatis trinerviis, cau- le erecto, feminibus hispidis. HALL. *H.* 2. ^{verum} LINN. *Sp.* 16. *Rubia prat. laevis, acuto folio.* B. *Pin.* 333. In M. *Altenstollberg.*

Galium caulis alis, foliorum ora nervoque serratis, fructu levi. HALL. *Aparine semine minore*. VAILL. *Par.* T. 4. f. 4.

Galium caulis alis, foliorumque ora nervoque serrato, fructu subaspero. HALL. G. 188. *Vaillantia* ^{Aparine} LINN. *Sp.* 2. *Aparine semine laevi*. VAILL. *Par.* T. 4. f. 3. In agris.

Galium caulis alis, foliorumque ora & nervo serratis, fructu hispido. HALL. H. 3. ^{Aparine} LINN. 19. *Aparine vulgaris*. B. *Pin.* 334. Ad sepes.

Galium foliis quaternis, flosculis in alis confertis. HALL H. 10. *Vaillantia floribus masculis quadrifidis, pedunculis diphyllis*. ^{Cruciata} LINN. *Sp.* 4. *Cruciata hirsuta*. B. *Pin.* 335. Flores mares & androgyni misti. Ad vias.

H. *Galium* feminibus exasperatis, flosculis lateralibus masculis. HALL. G. 188. *Aparine semine Coriandri saccharati* PARKINS. *Monspeliis*. ^{Halleri (H. p. 191) videtur eundem fuisse Vaillantia aparine}

H. *Galium* flosculis inter folia ovata, amplissima reflexa positis. HALL. G. 187. *Cruciata Orient. glabra humifusa*. I. R. H. *Cor.* ^{Vaillantia umbellata L. 2. p. 1491}

H. *Galium* floribus masculis trifidis, omnibus plantae partibus hispidis. *Vaillantia erecta hispida*. LINN. *Cliff.* 460. ^{Asperula} LINN. 2. *Sp.* 1. ~~Asperula~~

H. *Galium* floribus masculis trifidis, pedunculo communi cristato, foliis ciliatis. *Vaillantia procumbens glabra*. LINN. *Cliff.* 468. *Cruciata muralis minima Romana*. COLUMN. *Ecphr.* I. T. 297. Vtriusque diversitatem proxime fusius exponam. *V. muralis* Lin. 2. p. 2.

RVBIA. LINN. 19.

Calyx minimus. Flos quinquefidus. Baccae duae glabrae.

H. *Rubia* foliis senis. *tinctorum*. LINN. *Sp.* 1.
HALL. *H.* 462. *Rubia sylvestris aspera*. B. *Pin.* 33.
BLACKW. *T.* 326. In *Europa* australi.

SPERMACOCE. LINN. III. DILL.

Elth. 227.

Calyx minimus. Capsulae bicornes connatae.

H. *Spermacoce* glabra, staminibus inclusis. ^{tenuior} LINN.
Sp. 1. *Spermacoce verticillis tenuioribus*. DILL.
Elth. 370. *T.* 227. f. 359. In *Carolina*. A.

ANTHOSPERMUM. LINN. 1029.

Calyx revolutus sine petalo. Stamina receptaculo inferta. In polygamia dioicia.

H. *Anthospermum*. ^{Aethiopicum} LINN. *Cliff. T.* 27. *Sp. p.*
1058. *Frutex Africanus ambram spirans*. WALTH.
H. 24. *T.* 9. In horto planta mascula tantum adest.
In *Aethiopia*. Fr. ;

CORNVS. LINN. 139.

Flores umbellati. Bacca officulo biloculari.

Cornus arborea, umbellis involucrium aequantibus. *mas.* LINN. *Sp.* 2. *Cornus sylvestris mas.*
B. *Pin.* 447. CAMER. *Epit.* 158. In M. *Hengst*
prope *Lützelengen*.

Cornus arborea cymis nudis. *sanguinea*. LINN.
Sp. 3. *Cornus umbellis involucri multoties longioribus*.

vibus. HALL. H. 463. *Cornus femina.* B. Pin. 447.
Ad sepes.

ISOSTEMONES CONGLOMERATAE.

Stellatis affines sunt flore quadrifido, quatuor staminibus, & fructu saepe biloculari, qui in aliis sub flore, in aliis intra florem est. Flores semper aut in spicas aut alas congesti.

SANGVISORBA. LINN. 136. *Pimpinella.* HALL.

469. Poterium. LINN. 948.

Corolla persistens. Fructus bilocularis. Stigmata duo plumosa.

Sanguisorba tetrastemon, staminibus tubo floris aequalibus, spicis ovalibus. *Pimpinella.* HALL. l. c. n. 1. *Sanguisorba officinalis.* LINN. Sp. 1. *Pimpin. sanguisorba maior.* B. Pin. 160. Prope *Elbingeroode.* *Jen in Ned. Flor. Pan. Tab. 97.*

H. *Sanguisorba tetrastemon*, staminibus tubo floris, longioribus, spicis cylindricis. An *Pimpinella tetrastemon*, *spica ovata, foliis ovatis serratis.* HALL. G. 197. *Pimpinella minore di Canada.* ZANNON. T. 138. Flores rubri, spica cylindrica, flores superiores prius florere incipiunt, ut in priori specie. Diversissima a *sanguisorba Canad. spicis longissimis.* *Sanguisorba media dñ. 2 spec. 2.*

Poterium
Sanguisorba polytemon, staminibus longissimis pendulis. *Pimpinella.* HALL. n. 2. BLACKW. T. 413. *Poterium sanguisorba.* LINN. Sp. 1. *Pimpinella sanguisorba minor laevis.* B. Pin. 160. In siccis. P.

H. Sanguisorba polystemon, spinis ramosis, Pimpinella *ramosa*, *spinosa*. HALL. G. 197. MORIS. III. S. VIII. T. 18. f. 5. Poterium *spinosum*. LINN. Sp. 3. In Creta.

PLANTAGO LINN. 133. Psyllium
TOVRN.

Flores spicati, fructus intra florem.

Plantago foliis ovatis glabris, nudo scapo tereti, spica flosculis imbricatis. *maior*. LINN. Sp. 1. HALL. H. n. 5. Pl. *latifolia sinuata*. C. B. I. R. H. *septinervia*. BLACKW. T. 35. In cultis.

β. *Plantago latifolia rosea*, foliis inter flores longioribus enatis.

H. *Plantago* foliis latissimis, ovatis dentatis, rigidioribus, spica brevi, densa. AMMAN. n. 23. HALL. G. 199.

Plantago foliis ovato-lanceolatis pubescentibus, spica cylindrica, scapo tereti. *media*. LINN. Sp. 3. HALL. *Helv.* n. 4. Pl. *latifolia incana*. B. Pin. 189. Ad vias.

H. *Plantago* foliis lanceolato-ovatis pubescentibus subdenticulatis, spicis laxis pubescentibus, scapo angulato. *Virginica*. LINN. Sp. 4. Pl. *media incana*, *Virginiana*, *serratis foliis*. MORIS. III. S. 8. T. 15. f. 8. In Virginia. A.

'*Plantago* foliis lanceolatis, spica subovata nuda scapo angulato. *lanceolata*. LINN. Sp. 5. Pl. *angustifolia maior*. B. Pin. 189. BLACKW. T. 14. In gramineis.

H. Plan-

CONGLOMERATAE

237

H. *Plantago foliis lanceolatis, spica ovata hirsuta, scapo angulato.* *Lagopus*. LINN. *Sp.* 6. *Pl. angustifolia paniculis lagopi.* B. *Prodr.* 98. MORIS. III. S. 8. T. 16. f. 13. In *Europa australi.*

H. *Plantago foliis semicylindraceis integerrimis basi lanatis, scapo tereti.* *maritima*. LINN. *Sp.* 10. *Pl. marit. minima, folio gramineo, rigido.* I. R. H. In *falsis. v. g. am g. salzenen See.*

H. *Plantago foliis linearibus pinnato-dentatis, scapo tereti.* LINN. *Sp.* 10. *Coronopus sylvestris hirsutior.* B. *Pin.* 190. Eodem loco cum priori.

H. *Plantago caule ramoso, foliis integerrimis, spicis foliosis.* *Psyllium*. LINN. *Sp.* 15. HALL. *H. n.* 1. *Psyllium matius erectum.* B. *Pin.* 191. In *Marchia.*

Plantago (Psyllium) perennis, angustifolia & supina. I. R. H. *Psyllium.* CAMER. *Epi.* 811,

H. *Plantago caule ramoso, foliis dentatis, spicis aphyllis.* *Cynopr.* LINN. *Sp.* 16. *Psyllium Dioscoridis vel Indicum, foliis crenatis.* B. *Pin.* 191.

BASELLA. LINN. 343.

Corolla sexfida succulenta facta semen fovet.

H. *Basella foliis planis. rubra.* LINN. *Sp.* 1. *Fl. Zeyl.* 119. In *India.*

CVSCVTA. LINN. 156.

Planta aphylla, parasitica. Flores in glomeris congesti.

Cuscuta floribus sessilibus, repens filiformis. LINN. *Sp.* 1. *Cuscuta maior.* I. R. H.

β. Epi-

β. *Epithymum*. f. *Cuscuta minor*. C. B.

ILEX. LINN. 158. *Aquifolium*. T.

Flores quadrifidi ad alas foliorum congesti. Stylis quatuor.

Ilex foliis ovatis acutis spinosis. LINN. *Sp.* 1.

Aquifolium. HALL. *Helv.* 467. BLACKW. *T.* 205.

In arenosis *Cellenfis*.

ISOSTEMONES CAMPANACEAE.

Calyx quinquefidus. Corolla plerumque campanulata. Stamina tot, quot segmenta floris. Stylus simplex. Fructus in aliis sub flore, in aliis intra florem, quae autem ob summam analogiam a se invicem divelli nequeunt.

CONVOLVULVS. LINN. 198.

Petalum plicatum, limbo leviter quinquefido. Stigmata duo.

Convolvulus foliis sagittatis utrinque acutis, pedunculis unifloris biflorisque. *arvensis*. LINN. *Sp.* 1. HALL. *H.* 489. n. 2. B. *Pin.* 294. Volabilis *arvensis* flore roseo. TABERN. 877. In arvis ubique.

Convolvulus foliis sagittatis, postice truncatis, pedunculis unifloris. *sepium*. LINN. *Sp.* 2. *Conv. maior*, *albus*. I. R. H. BLACKW. *T.* 38. In sepibus.

Convolvulus foliis sagittatis, postice truncatis, pedunculis ramosis bifloris. LINN. *Sp.* 3. *Mat. med.*

82. *Convolvulus Syriacus*. MORIS, II, S. I. *T.* 3.

f. 5. *Scammonia Syriaca*. B. Pin. 294. DOD. Purg. 206. In *Syria*.

H. *Convolvulus* foliis ovato-lanceolatis, duobus foliis flori substratis. HALL. G. 205. ROYEN. 428. ^{Scand.} LINN. Sp. 17. *Convolv. Siculus minor, flore parvo articulato*. BOCC. Sic. 89. T. 48. In *Sicilia*.

H. *Convolvulus* foliis lanceolato-ovatis glabris, caule declinato, floribus solitariis. *tricolor*. LINN. Sp. 25. *Conv. peregrinus coerulescens*. B. Pin. 295. In *Europa australi*.

H. *Convolvulus* foliis lanceolatis obtusis sericeis, pedunculis multifloris. ^{Grecum} LINN. Sp. 21. *Conv. maritimus rectus, Creticus argenteus*. MORIS. II. S. I. T. 3. f. 1. *Conv. serpens maritimus spicae folio*. LOB. Ec. 622. P.

H. *Convolvulus* calycibus tuberculatis pilosis, foliis cordatis. ^{Aedonensis} LINN. Sp. 6. *a. coeruleus folio subrotundo*. DILL. Elsb. 97. T. 82. f. 94. Stigma tridum.

H. *Convolvulus* calycibus tuberculatis pilosis, folio anguloso. ^{vi?} LINN. Sp. 8. *ζ-η* DILL. Elsb. 96. T. 89. f. 90. 91. Stigma obtusum.

IPOMOEAE. LINN. 199. Quamoclit. T.

Petalum infundibuliforme. Stigma capitatum.

H. *Ipomoea* foliis cordatis acuminatis, basi angulatis, pedunculis multifloris. *coccinea*. LINN. Sp. 4. Quamoclit *Americ. folio bederae, flore coccineo*. COMMEL. rar. 21. T. 21. A.

H. Ipomoea foliis pinnatifidis linearibus, floribus solitariis. ^{Quamoclit} LINN. Sp. 1. Quamoclit. f. *Jasminum americanum*. CAMER. Hort. f. 40. CLVS. *Posth.* 9. PON. Bald. p. 71. In India.

H. Ipomoea foliis trilobis cordatis, pedunculis unifloris. An ^{Triloba} LINN. Sp. 12.? Folia accurate conveniunt, sed pedunculus triflorus dicit.

STRAMONIVM. TOVRN. 43. Datura.

LINN. 218.

Calyx circa basin rotunde deciduus. Petalum pliatum. Fructus hispidus.

Stramonium (*Datura*) pericarpus spinosis erectis ovatis. ^{Stramonium} LINN. Sp. 1. HALL. H. 489. BLACKW. T. 313. Solanum *manicum Dioscoridis*. COLUMN. *Phytobasf.* p. 47. Solanum *foetidum, pomo spinoso, oblongo*. c. B. Pin. 168. Sponte nunc in hortis & fimetis. β. flore & caule violaceo.

H. Stramonium calyce pentagono, fructu pendulo spinoso globofo. HALL. Gott. 206. *Datura* ^{varior} LINN. Spec. 2. Stramonium *ferox*. BOCCON. Sic. p. 5.

H. Stramonium fructu pendulo, spinoso, calyce campaniformi. HALL. Gott. l. c. Nux metella. CAMER. epit. 173. *Datura Metel Lini*.

H. Stramonium calyce campaniformi, fructu pendulo papillato. HALL. l. c. c. descr. *foetida* 2 Spec. 5.

H. Stramonium Aegyptiacum, flore pleno, intus albo, extus violaceo. 1. R. H. *Datura Contarena*. P. ALP. exot. 167. *Datura Aegyptiaca*. VESL. p. 58.

Varietas praecedentis.

MIRA-

MIRABILIS. LINN. 215. Jalappa.

TOVRN.

Petalum infundibuliforme ex nectario globoso ortum.

H. Mirabilis foliis corollisque glabris, corollis folio brevioribus. *Mirabilis Jalappa*. LINN. Sp. 1. Admirabilis *Peruviana*. CLVS. Hist. II. p. 87.

H. Mirabilis foliis villosis viscidis, tubo floris cylindrico, villoso, foliis longiore. COMM. Soc. Gott. T. V. ubi novam hanc speciem, icone addita, descripsi. Eodem tempore eiusdem descriptionem & iconem edidit Ill. LINNAEVS in Act. Soc. Holm. 1755. p. 176. T. VI. eandemque habet in Cent. I. plant. n. 18. *Mirabilis longiflora* Linn.

PLUMBAGO. LINN. 196.

Calyx hispidus. Petalum infundibuliforme. Valvulae circa germen staminiferae.

H. Plumbago foliis petiolatis. *Plumbago indica spicata*, *ocymastri foliis*, *fructibus sapaceis oblongis*. COMMEL. H. II. p. 169. T. 5. In India.

POLEMONIUM. LINN. 196.

Campanulae similis, sed fructus intra florem.

H. *Polemonium* calycibus corollae tubo longioribus. *coeruleum*. LINN. Sp. 1. HALL. H. 490. *vulgare coeruleum & album*. L. R. H. In Hassia, Helvetia.

CAMPANULA. LINN. 201.

Valvae latiores flaminiferae fundum floris claudunt. Stigma trifidum.

Campanula caule angulato, foliis petiolatis, calycibus ciliatis, pedunculis trifloris. *Trachelium*. LINN. *Sp.* 18. HALL. *H.* n. 1. Camp. *vulgatior vrticæ folio maior vel asperior*. B. *Pin.* 94. In sylvis. Folia subcordata. P.

Campanula foliis cordato-lanceolatis, alis unifloris, floribus secundis sparsis. LINN. *Sp.* 14. HALL. *H.* 2. Camp. *hort. rapunculi radice*. B. *Pin.* 94. Ad vias sylvarum.

H. Campanula foliis ovatis glabris subserratis, caule erecto paniculato, ramulis brevibus. LINN. *Sp.* 8. Camp. *pyramidalis altissima*. T. R. H. BAVH. *Pin.* 93.

H. Campanula capsulis quinquelocularibus, tectis calycis sinibus reflexis. LINN. *Spec.* 22. Campan. *hortensis, folio & flore oblongo*. B. *Pin.* 94. *Viola Mariana*. DOD. *Pempt.* 163. β. Varietas flore albo & γ. pleno.

H. Campanula capsulis trilocularibus, tectis calycis sinibus reflexis, floribus maculosis pendulis. *Sibirica*. LINN. *Sp.* 23. HALL. *G.* 193. In *Sibiria*. P.

Campanula floribus sessilibus latioribus foliis insidentibus, capitulo terminali. HALL. *H.* 4. *glomerata*. LINN. *Sp.* 19. Camp. *pratensis flore conglomerato*. B. *Pin.* 94. *Rapunculus umbellatus* &c.

THAL. *ic.* VIII. n. 2. In locis aridis circa *Harperhausen*.

Campanula hispida, racemo ovato-oblongo terminali, caule simplicissimo, foliis lanceolato-linearibus. ^{*perfoliatis*} LINN. *Sp.* 21. *Camp. foliis Echii*. B. *Pin.* 94. *Prodr.* 36. In sylvis montosis versus *Dudersbatt. Northheim: circa Rheinhausen*. ^{*Salp. Fr. L. hinc Drichl*}

Campanula foliis undulatis, radicalibus lanceolato-ovalibus, panicula coarctata. ^{*Campanuloides*} LINN. *Sp.* 5. HALL. *H. n.* 11. *Rapunculus esculentus*. B. *Pin.* 92. In colle arenoso prope *Boenden*.

Campanula foliis radicalibus obovatis, caulinis lanceolato-linearibus subserratis, floribus raris. perfoliata. LINN. *Sp.* 7. HALL. *H. n.* 12. *Rapunculus perfoliatus magno flore*. B. *Pin.* 93. flore coeruleo & albo. In dumetis. P. *c. currens* ^{*perfoliata*} *Sp.* 6. *ed.* 2.

Campanula foliis radicalibus reniformibus, serratis, caulinis linearibus integris. ^{*perfoliata*} LINN. *Sp.* 3. HALL. *Helv.* 14. *Camp. minor rotundifolia vulgaris*. B. *Pin.* 93. In gramineis. P.

Campanula fructu prismatico longissimo, flore rotato. HALL. *H. n.* 17. *Camp. speculum Veneris*. LINN. *Sp.* 27. *Camp. arvensis procumbens, aut erecta*. I. R. H. *Viola pentagonia*. TABERN. 316. In segetibus M. *Heynberg*. A.

H. *Campanula foliis perfoliatis, floribus inferioribus apetalis*. HALL. *G.* 195. LINN. *Sp.* 31. *Camp. pentagonia perfoliata*. MORIS. *S. V. T.* 2. f. 23. In *Virginia*. A.

PHYTEVMA, LINN. 203. Rapunculus
TOVRN.

Corolla rotata in lacinias lineares profunde fissa, reliqua prioris.

Phyteuma spica oblonga, foliis radicalibus cordatis, superioribus angustis. LINN. *Sp.* 4. Rapunculus. HALL. *H.* 5. Rapunc. *spicatus*, flore flavescente. 1. R. H. In sylvis. β . Varietas spica ovali obesa in M. Bruelero.

Phyteuma capitulo subrotundo, foliis ferratis, radicalibus cordatis. LINN. *Sp.* 3. Rapunculus. HALL. *H.* 3. Rapuntium *mont. corniculatum*. COLV. *Epbr.* I. T. 224. Rapunc. *spica orbiculari*. B. *Pin.* 92. In pratis prope Elbingerode.

JASIONE, LINN. 896.

Flores in capitulum collecti, rapunculi similes. Antherae connatae.

Jasione ^{montana}. LINN. *Sp.* p. 928. HALL. *H.* 496. Rapunculus *scabiosae capitulo coeruleo*. B. *Pin.* 92. In ficcioribus prope Bovenden. Rheinhausen. Seeburg.

LOBELIA, LINN. 897.

Floris tubus ad basin fissus, limbo fere ringente. Antherae connatae.

H. Lobelia caule erecto, foliis lanceolatis ferratis, spica terminali. *Cardinalis*. LINN. *Sp.* 10. Rapuntium *coccineum*. MORIS. *Hist.* II. S. V. T. 5. f. 54. RIVIN. *Monop. irreg.* T. 107. Flos fere tripetalos, calycis segmenta tubo aequalia. In *Virginia*. P.

H. Lobelia caule erecto, foliis ovato-lanceolatis erenatis, floribus lateralibus, calycum sinibus reflexis. ^{inpartem} LINN. Sp. 11. Rapuntium Amer. flore diluto coeruleo. I. R. H. 163. In Virginia.

H. Lobelia caule erecto, foliis ovatis subserratis pedunculo longioribus, capsulis inflatis. *inflata*. LINN. Sp. 12. Hort. Vps. 276. A.

H. Lobelia caulibus procumbentibus, foliis lanceolatis serratis, pedunculis lateralibus. ^{inpartem} LINN. Sp. 17. Campanula Afr. Erini facie, caulibus procumbentibus. HERM. Lugdb. p. 109. T. 52. In Africa. A.

TRACHELIVM. LINN. 204.

Corolla infundibuliformis. Stylus corolla multo longior. Flores corymbosi.

H. Trachelium (foliis ovatis serratis, corymbis compositis.) ^{coeruleum} LINN. Sp. p. 171. ROY. Lugdb. 248. Trachelium azureum umbelliferum. I. R. H. PON. Bald. p. 44. In Italia.

SAMOLVS. LINN. 205.

Corolla exigua. Squamae stamina munientes.

H. Samolus Valerandi. LINN. Sp. | p. 171. H. Vps. 42. Samolus Afr. folio rotundiore. WALTH. H. 162. T. 23. In maritimis. A.

ISOSTEMONES CVCVRBITACEAE.

Omnes fere in Monoecia. Mas. Calyx quinquepartitus; petalum campanulatum quinquepartitum

ex calyce ortum. Stamina filamenta tria, antherae quinque flexuosae. *Femina*. Flos insidet fructui carnosio; calyx & petalum, vt in mare; stylus simplex stigmate trifido. In omnibus caulis scandens.

MOMORDICA. LINN. 967. Elaterium.

H. Momordica pomis angulatis tuberculatis, foliis glabris patentipalmatis. *Balsamina*. LINN. *Sp.* 1. Mom. *vulgaris*. I. R. H. *Balsamina rotundifolia repens*. B. Pin. 306. Conf. charact. floris in *Comm. Soc. Gott. T. V.*

H. Momordica pomis angulatis tuberculatis, foliis villosis longitudinaliter palmatis. *Charantia*. LINN. *Sp.* 2. *Ceylanica* VOLCAM. *Hesp.* 244. *Balsamina cucumerina indica, fructu maiore flavescente*. COMM. H. 1. p. 103. T. 54.

H. Momordica pomis oblongis sulcatis catenulatis, foliis incis. *Luffa*. LINN. *Sp.* 3. Cucumis *Aegyptius reticulatus f. Luffa Arabum*. VESLING. *Aegypt.* 199. T. 58. 59. MORIS. I. p. 35. S. 7. T. 7. f. 1. 2. In *Ceylona*.

H. Momordica pomis hispidis, cirrhis nullis. ^{Elaterium} LINN. *Sp.* 6. Elaterium *Offic.*, Cucumis *sylvestris affinis*. B. Pin. 314. In *Europa australi*.

CUCURBITA. LINN. 968.

Cucurbita foliis subangulatis tomentosis, basi subtus biglandulosis, pomis lignosis. ^{lagenaria} LINN. *Sp.* 1. Cucurbita *lagenaria, flore albo*. MORIS. *Hist.* 2. S. 1. T. 5. f. 3.

H. *Cucurbita* foliis lobatis, pomis laevibus, seminum margine tumido. *Pepo*. LINN. *Sp.* 2. *Cucurbita maior Pepo oblongus*. B. *Pin.* 311.

H. *Cucurbita* foliis lobatis, pomis nodoso-verrucosis. LINN. *Sp.* 3. *Melo pepo verrucosus*. I. R. H. 106.

H. *Cucurbita* foliis lobatis, caule erecto, pomis depresso-nodosis. LINN. *Sp.* 4. *Cucurbita clypeiformis*. I. B. II. p. 224.

CUCUMIS. LINN. 969. Melo.

H. *Cucumis* foliis multifidis, pomis globosis glabris. LINN. *Sp.* 1. *Colocynthis* B. *Pin.* 313. LOB. *ic.* 644.

H. *Cucumis* foliorum angulis rectis, pomis oblongis scabris. LINN. *Sp.* 6. *Cucumis sativus vulgaris*. B. *Pin.* 310.

H. *Cucumis* foliis palmato-sinuatis, pomis subovatis echinatis. LINN. *Sp.* 2. *Cucumis sylv. Amer. anguriae folio, fructu ovi figura, spinosis tuberculis momordicae instar muricato*. PLUK. *Phyt.* 170. f. 3.

H. *Cucumis* foliorum angulis rotundatis, pomis torulosis. LINN. *Sp.* 4. *Melo vulgaris*. B. *Pin.* 310. BLACKW. *T.* 329.

H. β. *Melo minimus*, pomo non maior, sphaericus. HALL. *Gott.* 208. An *Cucumis hirsutus pomis subglobosis*. LINN. *Sp.* 5. *Melo variegatus, aurantii figura, odoratissimus*. DILL. *Elth.* T. 177. f. 218.

ET *Cucumis prophetarum*

BRYONIA. LINN. 970. *Melothria*.

LINN. 48.

Bacca ovalis, laevis. Sexus varius.

^{alba} Bryonia foliis palmatis, vtrinque calloso-scabris.
LINN. Sp. 1. Bryonia *foliis quinquangulis scabris*.
HALL. H. 506. Br. *alba*, *baccis nigris*. B. Pin. 297.
BLACKW. T. 37. Flores in dioicia. Ad sepes prope
portam Webndanam.

H. Bryonia foliis palmatis quinquepartitis vtrin-
que laevibus: laciniis pinnatifidis. ^{atrica} LINN. Sp. 5.
Bryonia *Afr. laciniata, tuberosa radice, floribus*
verbaceis. HERM. *Parad.* 107. T. 108.

H. Bryonia foliis palmatis scabris, laciniis lanceola-
tis serratis, lateralibus minimis. ^{acutula} LINN. Sp. 4. Fl.
Zeyl. n. 355. Bryonia Zeylonica fructu rubro, striis
albis variegato. Picta est in HERM. *Lugab.* T. 95.
Hort. Malab. T. VIII. T. 19.

^{Melothria} H. Bryonia floribus androgynis. HALL. G. 208.
^{acutula} Melothria. LINN. Sp. p. 35. Omnes flores andro-
gyni. In *America*.

SYCIOS. LINN. 971.

Bacca spinis obsita, unilocularis.

H. ^{angulata} Sycios foliis angulatis. LINN. Sp. 1. Bryo-
nioides *flore & fructu minore*. DILL. *Elth.* T. 51.
f. 59. In *Canada*.

PASSIFLORA. LINN. 910. Granadilla.

Calycis folia & petala quinque. Triplex corona
setarum circa stylum, qui stamina quinque & tres
stylos profert.

H. *Passiflora* foliis palmatis integerrimis, involu-
cris cordatis tryphyllis. *coerulea*. LINN. *Sp.* 22. H.
Cliff. 432. HALL. *G.* 210.

H. *Passiflora* foliis crenatis tripartito divisis.
HALL. *l. c.* An *Passiflora foliis trilobis serratis*.
(*incarnata*.) LINN. *Sp.* 21.

H. *Passiflora* foliis trilobis cordatis aequalibus
obtusis glabris integerrimis. *lutea*. LINN. *Sp.* 15.
Passiflora folio tricuspide, flore parvo flavescente.
I. R. H. HALL. *l. c.*

H. *Passiflora* foliis trilobis cordatis pilosis, involu-
cris multifido-capillaribus. *foetida*. LINN. *Sp.* 19.
HALL. *l. c.* *Clematis Indica hirsuta, foetida*. PLUM.
Amer. 71. T. 86.

ISOSTEMONES ROTACEAE.

Campanaceis valde similes calyce, staminibus & tu-
bis. Flos in plurimis infundibuliformis. Fructus po-
lypermus intra florem.

GENTIANA. HALL. 473. LINN. 285. *Swertia*
LINN. 284.

Corolla varia. Fructus oblongus, bivalvis, disty-
lus. Folia alterna, simplicia.

Gentiana corollis brevissime quadrifidis, imber-
bibus verticillatis, sessilibus. LINN. *Sp.* 20. HALL.
Sp. 3. *Gentiana Cruciata*. I. R. H. T. 40. C. B. *Pin.*
188. *Im Göttinger-Wald. In pratis prope Herper-*
hausen, in valle. Anthera vido

Gentiana corollis quadrifidis, corollis ciliatis. ciliata. LINN. *Sp.* 19. HALL. *H.* n. 1. *Gent. coerulea oris pilosis.* B. *Pin.* l. c. *Gent. coerulea, simbricata, autumnalis, angustifolia.* COLUMN. *Ecpbr. Tab.* 221. In sylvis. v. g. *Gottinger Wald.*

Gentiana foliis amplexicaulibus, corollis quinquefidis hypocrateriformibus, fauce barbata. ^{*amarula*} LINN. *Sp.* 17. HALL. *H.* 2. *Gent. autumnalis ramosa.* B. *Pin.* 188. In sylvis M. *Heymberg.*

H. *Gentiana uniflora, corolla quinquefida campanulata, caulem excedente.* ^{*acutius*} LINN. *Sp.* 8. HALL. *H.* 9. *Gent. alp. magno flore.* C. B. *Gentianella Helvetica.* LOBEL. *ic.* 310. In montibus *Helvetiae.*

Gentiana foliis linearibus, floribus in alis sessilibus. *Pneumonanthe.* LINN. *Sp.* 5. HALL. *H.* 10. LOBEL. *ic.* 309. *Gent. angustifolia autumnalis.* I. R. H. In paludosis ducatus *Cellensis.*

Gentiana foliis trinerviis, ovatis, floribus umbellatis, tubo strictissimo. ^{*Centaurium*} LINN. *Sp.* 14. HALL. *H.* 4. *Centaurium minus.* B. *Pin.* 178. BLACKW. *T.* 452. In sylvis.

β. *Centaurium minus palustre, ramosissimum.* VAILL. *T.* 6. f. 1. In paludosis. *et palus v. d. g. (v. g. 183)*

Gentiana caule brachiato, floribus flavis quadrifidis. HALL. *Gott.* 201. *filiformis.* LINN. *Sp.* 2. *Centaureum palustre, luteum, minimum.* VAILL. *T.* 6. f. 3. Circa ^{*Engelsen, v. g. 183*} *Engelsen* verius *Cellam.*

LYSIMACHIA. LINN. 188.

Flos rotatus. Capsula unilocularis decemvalvis.

Lyse

ANAGALLIS. LINN. 189.

Priori similis. Capsula horizontaliter dehiscens.

Anagallis foliis indivisis, flore phoeniceo. HALL.
H. 1. LINN. ^{crucifera} Sp. 1. BAVH. Pin. 252. Anag. terre-
stris mar. BLACKW. T. 43.

Anagallis flore coeruleo. HALL. H. 2. LINN. & c. ter-
restris femina. T. 274. Utraque super Papiermühle.

(β.) Anagallis foliis cordatis amplexicaulibus, cau-
libus compressis, LINN. ^{basifolia} Sp. 3. præter magnitudi-
nem omnium partium mihi quidem diversa non
videtur.

TRIENTALIS. LINN. 419.

Omnium partium numerus septenarius.

Trientalis. LINN. ^{europa} Sp. 1. Fl. Lapp. 139. Alsinan-
themum. HALL. Iter. Hercyn. §. 22. In rupibus prope
Waken & sylvis densioribus. Planta septentrionalis.

GLAVX. LINN. 257.

Calyx nullus. Petalum campanulatum.

Glaux maritima. LINN. Sp. p. 207. B. Pin. 215.
LOBEL. ic. p. 415. In prato sale turgido prope
Harste.

MENYANTHES. LINN. 185.

Segmenta floris fimbriata.

Menyanthes foliis ternatis, ^{trifoliata} LINN. Sp. 3. HALL.
Helv. 487. Trifolium palustre. BLACKW. T. 476.
fibrinum. TABERN. 521. & offic. In paludibus. v. g.
circa Diemern, Seeburg, Harste.

HOT-

HOTTONIA. LINN. 186.

Corollae infundibuliformis tubo stamina imposita.

Hottonia palustris. LINN. *Sp.* 145. HALL. *H.* 487. *Millefolium aquaticum*. B. *Pini* 141. *Stratiotes vulgaris*. VAILL. *Aff. Par.* 1719. p. 27. In vdis *Cellenfis*bus.

CYCLAMEN. LINN. 184.

Segmenta floris sursum flexa. Stamina ut in priori.

H. *Cyclamen corolla retroflexa*. LINN. *Sp.* 1. HALL. *H. n.* 1. *Artamita cyclamen* BLACKW. *T.* 147. LDB. *ic.* 604.

PHLOX. LINN. 197. *Lychnidea* DILL. Elth. 166.

Corolla infundibuliformis. Stamina inaequalia.

H. *Phlox foliis lato-lanceolatis*, inferioribus alternis, caule ramoso. *divaricata*. LINN. *Sp.* 5. *Lychnidea Virginiana*, *alsines aquaticae foliis*. PLUK. *Mant.* 121. In *Virginia*.

PRIMVLA. LINN. 180.

Flores plures involucro excepti. Calyx quinqueangularis.

Primula foliis rugosis, hirsutis, scapo unifloro. HALL. *H.* 1. *Primula veris*. BLACKW. *T.* 52. singulari flore. I. R. H.

β. calyce colorato in segmenta petaloidea fissus.

Primula foliis rugosis hirsutis, spica inclinata, flore pallido inodoro. HALL. *Helv.* n. 2. *Primula veris pallido flore elatior*. CLVS. *Hist.* I. p. 301. 1. R. H. LINN. *Sp.* I. β . In pratis & sylvis.

Primula foliis rugosis hirsutis, spica inclinata, flore flavo odoro. HALL. *U.* 1. *Primula odorata, flore luteo simplici*. 1. R. H. Paralysis. BLACKW. *T.* 226. In sylvis. Omnino a priori diversa.

Primula foliis serratis carnosis glabris. LINN. *Sp.* 5. HALL. *H.* 5. *Auricula vrsi*. CLVS. *Hist.* 301.

β . Varietates hortenses variorum colorum.

ANDROSACE, LINN. 179.

Flores ex involucro. Petalum tubo ventricoso.

H. Androsace perianthiis maximis. LINN. *Sp.* I. *Androsace vulgaris latifolia*. 1. R. H. Andros. *altera*. CLVS. *Hist.* II. p. 134.

Androsace foliis lanceolatis dentatis glabris, perianthiis angulatis corolla brevioribus. LINN. *Sp.* 2. Andros. *Coronopi folio glabra, flosculis ex verticilli foliosi umbilico plurimis albis*. AMMANN. *Roth.* n. 22. In *Sibiria*.

COFFEA, LINN. 209.

Corolla infundibuliformis. Stigmata duo. In bacca femina duo.

H. *Coffea*. LINN. *Sp.* 172. JUSSIEV. *Art. Par.* 1713. p. 388. T. 7. TREW. *Commer. Nor.* 1745. hebdomada 52. BLACKW. *T.* 337. BON, ALPIN, *Aegypt.* T. 36. In *Arabia*.

LYCIUM. LINN. 1232.

Superiori similis, sed bacca bilocularis, polysperma.

H. *Lycium* foliis cuneiformibus. *Europaeum*. LINN. Sp. 3. *Jasminoides aculeatum*, *falidis folio*, *flore parvo ex albo purpurascens*. MICHEL. gen. 224. T. 105. f. 1. In Europa australi.

H. *Lycium* foliis linearibus. *Afrum*. LINN. Sp. 1. *Jasminoides aculeatum humile*, *balimi minoris folio*, *flore violaceo*. MICH. gen. 224. T. 105. f. 2. In Africa.

ISOSTEMONES ASPERIFOLIAE.

Calyx quinquefidus. Corolla limbo quinquefido. Stamina plerumque in tubo floris latentia, minima; tubum saepe claudunt pali aut glandulae aut plicae. In fundo tubi semina quatuor nuda, in quorum medio exsurgit tuba unica simplex. Folia alterna, aspera aut hirsuta.

ECHIUM. LINN. 175. HALL. H. 513.

Corolla campaniformis limbo inaequali.

Echium caule simplici erecto, foliis caulinis lanceolatis hispidis, floribus spicatis lateralibus. *vulgarre*. LINN. Sp. 3. HALL. H. n. 1. BLACKW. T. 299. CLVS. CLXIII. Vbique ad vias.

H. Echium calycibus fructescens distantibus, caule procumbente. *Creticum*. LINN. Sp. 4. Ech. *Cret. rubrum*. B. Pin. 254. CLVS. Hist. 2. CLXV.

CERIN-

CERINTHE. LINN. 171. HALL. H. 515.

Corolla regularis tubulato ventricosa, fauce nuda pervia. Capsulae seminales binae, dispermae.

H. Cerinthe foliis amplexicaulibus, dentibus floris revolutis brevissimis. HALL. H. 1. *maior*. LINN. Sp. 1. CLVS. CLXVII. α . flore rubro purpurascente,

H. β . flavo flore asperior. B. Pin. 258. In *Helvetia, Italia*. &c. A.

H. Cerinthe corollis fere semiquinquefidis acutis, clausis. HALL. H. 2. *minor*. LINN. Sp. 3. CLVS. CLXVIII. B. Pin. 268. Glastum montanum. DALECH. Prope *Jenam*.

SYMPHYTUM. LINN. 170. HALL. H. 514.

Corolla priori similis, sed faux palis clausa.

Symphytum foliis ovato-lanceolatis, decurrentibus. LINN. Sp. 1. HALL. H. 1. *Symph. Consolida* *maior*. BLACKW. T. 252. flore pallido, *femina*. V-bique in pratis.

β . flore purpureo *mas*. In *Helvetia*.

PULMONARIA. LINN. 169.

Calyx pentagonus. Limbus floris erecto-patens. In fauce vngues breves pilosi.

Pulmonaria foliis radicalibus ovato-cordatis scabris. *officinalis*. LINN. Sp. 2. HALL. H. 1. *maculosa*. BLACKW. T. 276.

β . *non maculosa*. In sylvis.

LITHOSPERMUM. LINN. 166. HALL.

H. 517.

Calycis segmenta subulata. Faux plicis munita.
Semina nitentia.

H. Lithospermum seminibus laevibus, floribus in
alis sessilibus parvis, foliis nervosis. HALL. H. 2.
LINN. Sp. 1. *maius erectum*. B. Pin. 158. I. R. H.
Miliun folis. 1. B. In *Helvetia*.

Lithospermum seminibus rugosis, corollis vix caly-
cem superantibus. LINN. Sp. 2. HALL. H. 3.
BLACKW. 436. *arvense radice rubra*. C. B. Vbique
ad vias.

Lithospermum corollis calycem multoties super-
antibus, caulibus procumbentibus radicatis. LINN.
Sp. 4. HALL. H. 1. Lithosp. *repens maius*. CLVS.
CLXIII. In praecipitio montis super *Lutielengen*.
In sylva prope *Grossen-Schneen*, P.

HELIOTROPIUM. LINN. 164. HALL.

Limbus planus. Tubus plicis concoloribus
clausus.

H. Heliotropium foliis ovatis acutis, spicis soli-
tariis, fructibus bifidis. LINN. Sp. 1. *americanum*
coeruleum, *hormini folio*. FLVK. Phyt. 245. f. 4.
HERM. Lugdb. 307. In *India*, P.

H. Heliotropium foliis ovatis integerrimis rugo-
sis, spicis coniugatis. LINN. Sp. 2. HALL. H. 520.
Hel. *maius Dioscoridis*. B. Pin. 253. CLVS. XLVI.
In *Europa australi*, A.

H. *Helitropium* foliis lanceolato-linearibus glabris, spicis coniugatis. ^{*crucianicum*} LINN. Sp. 4. *americanum procumbens, facie lini umblicati*. HERM. Parad. T. 143. In America.

MYOSOTIS. LINN. 165. *Scorpiurus* HALL. 519.

Limbus planus. Faux clausa glandulis coloratis.

Myosotis feminibus nudis, foliis hirsutis, radice annua, flore minori. ^{*scorpioides*} LINN. Sp. 1. α . *Scorpiurus*. HALL. 1. *Echium scorpioides arvense*. c. B. flore coeruleo in humidis, albo in macilentis.

Myosotis foliis subhirsutis, radice perenni, flore maiori. ^{*palustre*} LINN. Sp. 1. β . *Scorpiurus* HALL. 2. *Echium scorpioides palustre*. c. B. in palustribus. β . Varietas hirsutior in ficcis.

H. *Myosotis* feminibus aculeatis glochidibus, foliis ovato-oblongis, ramis divaricatis. *Virginiana*. LINN. Sp. 2. Semina angulata. Fructus nutans. In Virginia.

Myosotis feminibus aculeatis glochidibus, foliis lanceolatis pilosis. *Lappula*. LINN. Sp. 3. *Cynoglossum foliis lingulatis, floris minimi tubo longiore*. HALL. H. 1. *Cynogl. minus*. B. Pin. 257. *Cynoglossa montana, minor, altera serotina*. COLUMN. Ecphr. p. 180. Circa Plesi. In muris.

CYNOGLOSSVM. LINN. 168.

Pali villosi tubum claudunt. Semina compressa, scabra.

Cynoglossum staminibus corolla brevioribus, foliis lato-lanceolatis tomentosis sessilibus. *officinale*. LINN. Sp. 1. HALL. H. 2. BLACKW. T. 249. Cyn. vulg. B. Pin. 257. Vbique ad vias.

Cynoglossum staminibus corollam aequantibus, foliis asperis. LINN. Sp. 4. Cynogl. mont. flore minore, folio virenti. 1. R. H. Cyn. mont. maximum frigidarum regionum. COLUMN. Ecphr. I. T. 170. In rupibus prope Herzberg; super Ilfeld.

H. Cynoglossum corollis calyce duplo longioribus, foliis lanceolatis sericeis. LINN. Sp. 3. Cyn. Creticum argenteo angusto folio. B. Pin. 257. Cyn. Creticum L. CLVS. CLXII. Corollae albae venis sanguineis. In Creta.

H. Cynoglossum foliis lineari-lanceolatis glabris. *linifolium*. LINN. Sp. 5. Cynogl. minus album, lini foliis glaucis, semine umbilicato. MOR. III. S. XI. T. 30. f. 11. Omphalodes Lusitanica elatior, cynoglossi folio. 1. R. H. 140. In Lusitania.

H. Cynoglossum repens, foliis radicalibus cordatis. *omphalodes*. LINN. Sp. 6. Omphalodes verna symphyti folio. Hort. G. 223. Symphytum repens, Borriginis facie, B. Pin. 209. In Lusitania.

ANCHVSA. LINN. 167. Buglossum.

HALL. & TOURN.

Calyx teres tubo cylindrico floris aequalis.

Anchusa foliis lanceolatis, spicis imbricatis secundis *officinalis*. LINN. Sp. 1. Buglossum floribus longe spicatis. HALL. H. 1. Bugl. angustifolium ma-

ius. B. Pin. BLACKW. T. 500. Circa Hannoveram.

LYCOPSIS. LINN. 174. HALL. H. 523.

Tubus floris incurvus ventricosus. Semina sublonga.

Lycopsis foliis lingulatis, sinuatis, hispidis, calycibus florecentibus erectis, *arvensis*. LINN. Sp. 1. HALL. H. 523. Echioides RIV. T. 7. Buglossum sylv. minus. C. B. In agris inter segetes. P.

H. Lycopsis foliis integerrimis, calycibus frutescentibus inflatis pendulis, ^{vesicularia} LINN. Sp. 1. Lyc. flore pullo calycibus amplis. HALL. G. 224. Echioides flore pullo. RIV. T. XI. In Thuringia. A.

H. Lycopsis foliis repandis dentatis callosis, caule decumbente, corollis cernuis. *variegata*. LINN. Sp. 2. Lyc. foliis hispidis verrucosis, flore variegato. HALL. G. l. c. Buglossoides Cretica. RIV. T. 9. In Creta. A.

ASPERVGO. LINN. 173.

Calyx quindecimfidus, laciniis quinque maioribus, figura bivalvi semina comprehendit.

Asperugo. LINN. p. 138. HALL. H. 522. Buglossum ^{procumbens} sylvestre, caulibus procumbentibus. B. Pin. C. B. In vallis vrbis ad portam Weendanam. A.

BORRAGO. LINN. 172.

Calyx & flos sine tubo stellatus, expansus.

H. Borrago foliis omnibus alternis, calycibus patentibus, *officinalis*. LINN. Sp. 1. HALL. H. 524.

BORRA

Borrago. BLACKW. T. 36. floribus coeruleis & β . albis. Sponte in hortis. Praeter palos elastra, stamina virgentia. A.

H. Borrago radice perenni, flore reflexo. Differt a vulgari defectu unguium & longiori tubo floris.

H. Borrago foliis ramificationum oppositis, calycinis foliis sagittatis. LINN. ^{Indica} Sp. 2. Cynoglossoides folio caulem amplexante. Act. Par. 1725. f. 10. A.

H. Borrago calycibus tubo corollae brevioribus, foliis cordatis. LINN. ^{orientalis} Spec. 4. Borrago Constantinopolitana, flore reflexo coeruleo, calyce vesicario. TOURN. It. p. 523. Circa Constantinopolin. P.

Nolana prostrata Desf. et al. in *Ann. Bot. Tab. 3.*

ISOSTEMONES SOLANACEAE.

Calyx quinquefidus. Corolla limbo quinquepartito, regulari aut irregulari. Stylus simplex. Fructus ut plurimum bilocularis siccus aut succulentus, seminibus receptaculo communi affixis. Odor gravis. Vires anodynae, narcoticae.

SOLANVM. LINN. 224.

Calyx & corolla rotata. Antherae conniventes. Fr. bacca.

Solanum caule inermi herbaceo, foliis ovatis dentato-angulatis; umbellis nutantibus. *nigrum*. LINN. ^{in solanum turgidum} Sp. 12. HALL. H. 2. *hortense*. BLACKW. T. 107. *officinarum*. B. Pin. 166. Ubique in cultis & hortis. A.

β. *Solanum Guineense*, fructu magno instar *ce-
rasi*. DILL. *Elth.* T. 274. f. 354.

h. *Solanum* ramis teretibus, foliis & floribus
villosis, baccis luteis. ^{in lasium} LINN. *Sp.* 12. γ. DILL. *Elth.*
274. f. 353. a vulgari, constanti partium villositate
diversum videtur.

h. *Solanum* caule inermi herbaceo, foliis pinna-
tis integerrimis. *tuberosum*. LINN. *Sp.* 8. *Solan. tu-
berosum esculentum*. B. *Prodr.* T. 89. *Papas Peru-
vianorum*. Tartüßeln, Gottingae copiosissime co-
litur. A.

h. *Solanum* caule inermi herbaceo, foliis pinna-
tis incis, racemis simplicibus. *Lycopersicum*. LINN.
Spec. 9. *Poma amoris fructu luteo*. CAMER. *Epit.*
821. In *America*. A.

Solanum caule inermi frutescente flexuoso,
foliis superioribus hastatis, racemis cymosis. LINN.
Sp. 5. *Solan. scandens lignosum, f. Dulcanara*. I. R.
H. BLACKW. T. 34. In paludosis ad sepes. P.

h. *Solanum* caule frutescente subinermi, foliis
cuneiformibus sinuato-repandis. ^{Bonariense} LINN. *Sp.* 7. *Solan.*
Bonariense arborescens, papas floribus. DILL. *Elth.*
T. 272. f. 351. Folia & flores maximi. In agro
Bonariensi. Fr.

h. *Solanum* caule inermi fruticoso, foliis lanceo-
latis repandis, umbellis sessilibus. *Pseudo-capsicum*.
LINN. *Sp.* 3. *Solanum fruticosum bacciferum, Ama-
mum Plinii*. I. R. H. Fr.

h. *Solanum* caule inermi fruticoso, foliis ovatis
integerrimis, floribus ad basin ramulorum confertis.

LINN.

LINN. *Sp.* 2. *Solan. Afr. lignosum*, folio atroviridi angusto oblongo obtuso. BOERH. II. p. 68. In Guinea. Fr.

H. *Solanum* foliis ovato-lanceolatis, fructu calloso, striato. HALL. G. 213. c. descr. An *Solanum* LINN. *Sp.* 4.?

H. *Solanum* caule inermi herbaceo, foliis ovatis tomentosis integris, calycibus spinosis. LINN. *Sp.* 13. *Melongoena fructu oblongo luteo*. Mala insana. DOD. *Pempt.* 458. β. fructu rubro longo.

H. *Solanum* caule aculeis recurvis, foliis sinuatis subtus tomentosis utrinque aculeatis, pedunculis (& calycibus) aculeatis. LINN. *Sp.* 19. *Fl. Zeyl.* 95. * *Solanum pomiferum frutescens, Africanum, spinosum, nigricans, borraginis flore, foliis profunde laciniatis*. HERM. *Lugdb.* 573. T. 575. MOR. III. S. XIII. T. I. f. 15. PLUK. T. 226. f. 5.

CAPSICVM. LINN. 225.

Calyx erectus. Flos rotatus plicatus. Baccā magna exsucca.

H. *Capsicum* caule herbaceo, *annuum*. LINN. *Sp.* 1. *Piper Indicum vulgatissimum*. B. *Pin.* 132. Varietates ex fructu plurimae.

α. *Capsicum siliquis longis propendentibus, purpureis*.

β. *Caps. fructu flavo pyramidalis, oblongo*. I. R. H.

γ. *Caps. fructu latiore & rotundiore*. I. R. H.

δ. *Caps. fructu parvo, rotundo, acerrimo*. I. R. H.

- e. Caps. Americ. fructu luteo, minimo.
 g. Caps. Americ. fructu parva, conica, erecto.
 θ. Caps. fructu cordiformi erecto, & flore, ac in
 reliquis, minori.

PHYSALIS. LINN. 223. Alkekengi. T.

HALL. 506.

Calyx inflatus, coloratus, clausus fructum exsuscum comprehendit. Stamina libera.

H. *Physalis* caule fruticoso, foliis ovatis integerrimis, floribus confertis. LINN. *Cliff.* 62. *somnifera*. LINN. *Sp. L.* Alkekengi fruticosum, floribus verticillatis. HALL. G. *Solanum somniferum*. CLVS. II. p. 85. In Syria, Creta.

Physalis foliis integerrimis cordatis geminis, floribus solitariis. LINN. *Sp.* 5. *Mat. med.* 93. Alkekengi. f. *Halicacabum*. BLACKW. T. 161. *officinarium*. I. R. H. *Solanum vesicarium*. C. B. Pin. 166. In M. Pless.

H. *Physalis* ramosissima, ramis angulosis glabris, foliis dentato-serratis, *angulata*. LINN. *Sp.* 6. Alkekengi *Indicum glabrum chenopodii folio*. DILL. *Elib.* T. 12. f. 12. *Solanum vesicarium Indicum*. COMM. H. I. p. 70. T. 17. A.

H. *Physalis* ramosissima, foliis villosis-viscosis, ramis pubescentibus, pedunculis nutantibus. *pubescens*. LINN. *Sp.* 7. Alkekengi *Virginianum, fructu luteo*. FEVILL. *Peruv.* III. p. 5. T. 1. *Solanum vesicarium Virgin. procumbens annuum, folio lanuginoso*. MORIS. III. p. 27. S. 13. T. 3. f. 24. Corollae maculae fuscae. In India. A.

ATRO-

ATROPA. LINN. 222. Belladonna. T.

HALL. 506.

Calyx & corolla tubuloso-campaniformis.

Atropa caule herbaceo, foliis ovatis integris.

LINN. Sp. 1. Belladonna. HALL. H. 507. Solanum melano-cerasus. B. Pin. 166. lethale CLVS.

II. LXXXVI. In sylvis copiose. A.

H. Atropa foliis sinuato-angulatis, calycibus

clausis acutangulis. *physalodes*. LINN. Sp. 2. cum

descr. Alkekengi amplo flore violaceo. FEVILL.

Peruv. 724. T. 16. A.

NICOTIANA. LINN. 220.

Corolla infundibuliformis calyce longior, capsula ovata.

II. Nicotiana foliis lanceolatis. *Tabacum*. LINN.

Sp. 1. Nicot. maior. latifolia. B. Pin. 169. *Tabacum*. Petum. copiose colitur. A.

H. Nicotiana foliis ovatis. *rustica*. LINN. Sp. 2.

Nicotiana minor. B. Pin. 170. In America. A.

H. Nicotiana foliis cordatis, floribus paniculatis.

tubis clavatis. *paniculata*. LINN. Sp. 3. Nicotiana minor, folio cordiformi, tubo floris praelongo. FEVILL. Peruv.

I. p. 717. T. 10. HALL. Gott. 219. In Peru.

H. *Nicotiana glauca*. LINN. Sp. 3. Nicotiana glauca.

VERBASCUM. LINN. 217. *celsea*.

HALL. H. 509.

Corolla rotata, limbo irregulari. In plerisque

stamina inclinata & villosa. Fr. succus.

Verbascum foliis decurrentibus vtrinque tomentosis, floribus densissime spicatis sessilibus. HALL. H. 1. *Thapsus*. LINN. Sp. 1. Verbasc. *mar. latifolium, luteum*. B. Pin. 239. Vbique ad vias. A.

β. Verbascum *flore maiore, stigmate bicolli*.

H. Verbascum foliis sessilibus vtrinque viridibus, non decurrentibus, fasciculis florum remotis. Verbasc. *foliis viridibus annuum, floribus luteis*. HERM. H. Lugdb. 618. HALL. H. 216. simile *Verbasco mari*, foliis autem & spica rariori diversa. A.

Verbascum foliis serratis cuneiformibus oblongis inferne villosis, pedunculis ramosis. HALL. H. 2. *Lychnitis*. LINN. Sp. 2. Verbasc. *flore albo parvo*. C. B. 240. Verb. *candidum femina*. Rarius in ruderatis. A.

Verbascum foliis cordato-oblongis petiolatis, pedunculis ramosis, staminum barba purpurascente. HALL. H. 3. LINN. Sp. 3. Verb. *nigrum, flore ex luteo purpurascente*. I. R. H. Ad viam versus *Papiermühle*. In quatuor speciebus modo dictis praeter reliquas partes calyx & fructus lana obducti sunt.

H. Verbascum foliis sessilibus longe acuminatis, serratis, floribus in raram spicam dispositis, solitariis, staminum barba purpurascente. Planta vix satis nota. Summitates plantae vndique pilis glanduliferis obsitae. Flores fere sessiles, rari, tribus bracteis succincti. Fructus subrotundus calyci aequalis. Folia hirsuta non tomentosa.

A. N. C. Tern. A. p. III.

H. *Verbascum* foliis amplexicaulibus oblongis glabris, floribus solitariis longius pedunculatis. *Blattaria*. LINN. *Sp.* 4. *Blattaria lutea, folio oblongo laciniato*. C. B. 240. α . flore albo extus purpureo; β . flore luteo. Folia radicalia sinuata, superiora serrata. Summitates pilis glanduliferis obsitae. In Germania australi. A.

H. *Verbascum* foliis ovatis nudis crenatis, caule subnudo racemoso. *phoeniceum*. LINN. *Sp.* 7. *Verb. foliis glabris indivisis*. HALL, G. *Blattaria purpurea*. B. *Pin.* 241. In Hungaria, Bohemia, P.

H. *Verbascum* tetrastremon. HALL, G. 217. *Celsia foliis duplicato-pinnatis*. LINN. *Sp.* 621. *Verbascum orientale, sophiae folio*. TOURN. *Cor.* 8. In Oriente. Habitus totus *Verbasci*, sola stamina recedunt. A.

HYOSCYAMVS. LINN. 219.

Corolla limbo irregulari. Fructus operculo tectus.

Hyoscyamus foliis amplexicaulibus. *niger*. LINN. *Sp.* 1. *Mat. med.* 36. HALL, H. 512. *Hyosc. vulgaris vel niger*. B. *Pin.* 169. Ubique ad vias & in ruderalis. A.

H. *Hyoscyamus* foliis crispis circumferratis petiolatis, floribus pedunculatis. HALL, G. 218. LINN. *Sp.* 3. *Hyosc. Cretic. luteus minor*. I. R. H. C. B. *Pin.* *Hyosc. aureus*. ALPIN. *Exot. p.* 98. In Creta. *Monspeliis*. A.

H. *Hyoscyamus* foliis petiolatis oblongis ampliter dentatis, floribus pedunculatis. *Hyosc. minor albo similis, umbilico floris atropurpureo*. TOURN. *Cor.*

MANETT. *Virid.* 518. Habitus totus albi, nisi quod flores pedunculati sint, in fundo violacei. A.

H. *Hyoscyamus* foliis petiolatis, floribus sessilibus, ^{albus} LINN. *Sp.* 2. *Hyosc. albus vulgaris.* CLVS. CXVIII. In *Europa* australi. A.

H. *Hyoscyamus* foliis lanceolatis subdentatis, calycibus spinosis. *pufillus*, LINN. *Sp.* 4. HALL. G. 218. In *Persia*. A.

ISOSTEMONES DICARPAE.

S. CONTORTAE.

- Calyx quinquepartitus. Corolla quinquesida. Segmenta floris oblique dextrorsum flexa. In plurimis circa stamina nectaria cava, antherae conniventes aut connatae, & stigmata duo corpusculo cavo recta. Folliculi duo polyspermi, seminibus pappo involutis.

NERIVM, LINN. 262.

Corollae tubus terminatus corona lacera.

H. *Nerium* foliis lineari-lanceolatis ternis. *Oleander*. LINN. *Sp.* 1. *Rhododendrum*. DOD. *Pempt.* 851. In *Palæstina*.

β. *Nerium indicum*, flore pleno. HERM. *H. Lugdb.* p. 549. Fr.

VINCA. LINN. 261. *Pervinca*

HALL. H. 526.

Corolla hypocrateriformis. Stylus stigmatibus duplici, inferiori orbiculato, superiori capitato.

Vinca

Vinca caulibus procumbentibus, foliis lanceolato-ovatis. LINN. *Sp.* 1. *Pervinca vulgaris angustifolia*. I. R. H. Clematis *Daphnoides minor*. C. B. Pin. In sylvis. v. g. super *Herperhausen*. Fr.

H. Vinca caulibus erectis, foliis ovatis. *maior*. LINN. *Spec.* 2. *Pervinca latifolia flore coerulesco*. GARID. *T.* 81. In *Gallia Narbonensi*. Fr.

II *Vinca vicia*

PERIPLOCA. LINN. 267.

Corolla rotata. Nectarium monophyllum ligulas emittens. Stigmata tecta.

H. *Periploca* foliis oblongis, floribus interne hirsutis. LINN. *Sp.* 1. HALL. *G.* 225. c. descr. Apocynum II. *angustifolium*. CLVS. *Hist.* I. p. 125. In *Syria*. P.

ASCLEPIAS. LINN. 270.

Flores umbellati. Nectaria cava cum corniculis circa machinam stamineam operculatam.

H. *Asclepias* foliis ovalibus subtus tomentosis, caule simplicissimo, umbellis nutantibus. *Syriaca*. LINN. *Sp.* 4. Apocynum *maius Syriacum rectum*. CORNVT. *Canad.* T. 93. Apocyn. *Syriacum*. CLVS. *Hist.* II. LXXXVII. P.

H. *Asclepias* foliis lanceolatis, caule superne diviso, umbellis terminalibus congestis. *incarnata*. LINN. *Sp.* 8. Apocynum *minus rectum Canadense*. CORNVT. *Canad.* 9. T. 93. In *Canada*. P.

H. *Asclepias* foliis lanceolatis petiolatis glabris, caule simplici, umbellis erectis solitariis. *Curassavica*.

vica. LINN. *Sp.* 9. *Apocynum radice fibrosa, petalis coccineis, corniculis croceis*. DILL. *Elth.* T. 30. f. 33. *Apocynum Curassavicum, fibrosa radice, floribus aurantiis, chamaenerii foliis latioribus*. HERM. *Par.* T. 36. HALL. G. 226. P.

H. *Asclepias caule erecto fruticoso, foliis revolutis lanceolatis, umbellis nutantibus, conceptaculis inflatis pilosis. fruticosa*. LINN. *Sp.* 14. *Apocynum erectum Afr. villoso fructu, salicis folio*. HERM. *Par.* 23. T. 24. COMMEL. H. II. f. 25. In *Syria*. Fr.

Asclepias foliis ovato-lanceolatis, glabris, floribus racemosis, albis. HALL. H. 525. ^{*Vincetoxicum*} LINN. *Sp.* 12. *Asclepias albo flore* B. *Pin.* 303. *Vincetoxicum*. DODON. *Pempt.* 407. In collibus lapidosis. v. g. M. *Pless. Hengst.* P.

APOCYNVM. LINN. 269.

Nectaria simplicia. Antherae conniventes.

H. *Apocynum caule rectiusculo herbaceo, foliis ovatis vtrinque glabris, cymis terminalibus*. LINN. ^{*apocyni folium*} *Spec.* 1. *Apocynum Canad. foliis androsaemi maioris*. BOCC. *Sic.* 35. T. 16. f. 3. MORIS. III. p. 609. S. 15. T. 3. f. 16. In *Virginia*. Fr.

STAPELIA. LINN. 271. HALL. G. 227.

Corolla rotata. Nectarium stellatum, lacerum. Machinam stamineam & stigmata stellula tegit.

H. *Stapelia caulibus aculeatis, flore levi*. HALL. G. 227. c. descr. *Stapelia denticulis ramorum patent-*

tentibus. (variegata.) LINN. *Sp.* 1. *Apocynum humile aizoides, siliquis erectis, africanum.* HERM. *Lugdb.* T. 53. Fr.

H. *Stapelia* caulibus quadrangulis, flore pubescente. HALL. *G.* 228. *Stapelia denticulis ramorum erectis. (hirsuta.)* LINN. *Sp.* 2. *Asclepias Afr. aizoides, flore pulchre fimbriato.* COMMEL. *rar.* T. 19. Fr.

ISOSTEMONES FLORE DIFFORMI.

Petala quatuor aut quinque irregularia; Epimedio excepto, cui flos regularis. In omnibus nectaria. Stamina antheris conniventes. Fructus est siliqua modo longior, modo brevior.

MELIANTHVS. LINN. 712.

Flos papilionaceo similis, in cuius fundo fovea nectarifera. Stamina quatuor.

H. *Melanthus* stipulis solitariis petiolo adnatis. *maior.* LINN. *Sp.* 1. *Melanthus Africanus.* HERM. *Lugdb.* 414. T. 415. In *Aethiopia.* Fr.

H. *Melanthus* stipulis geminis distinctis. *minor.* LINN. *Sp.* 2. *Melanthus Africanus foetidus minor.* COMMEL. *rar.* 4. T. 4. Fr.

HYPECOVM. LINN. 157.

Petala quatuor inaequalia trifida. Siliqua articulata.

H. *Hypocoum* siliquis erectis torulosis. LINN. *Sp.* 3. *Hypocoum tenuifolium, siliquis erectis teretibus,*

bus. AMMANN. *Ruth.* 58. T. 9. HALL. *Gott.* 230.
In *Sibiria*. A.

VIOLA. LINN. 898.

Petala quinque, infimum calcare nectarifero. Siliqua ovata.

Viola acaulis, foliis cordatis, stolonibus reptantibus. *odorata*. LINN. *Sp.* 5. HALL. *H.* 1. *Viola martia*. BLACKW. *T.* 55. *Viola martia purpurea, flore simplici odoro*. B. *Pin.* 119. I. R. H.

Viola caulis adscendentibus floriferis, foliis oblongo cordatis. *canina*. LINN. *Sp.* 9. *Viola martia inodora sylvestris*. B. *Pin.* 199. In sylvis.

H. *Viola* foliis reniformibus, caulis foliosis floriferis. HALL. *H.* 5. *Viola caule bifloro, foliis reniformibus serratis. (biflora.)* LINN. *Sp.* 16. *γ. alp. rotundifolia minor lutea*. FLVKN. *T.* 233. f. 7. In rupibus *Helvetiae*.

H. *Viola* caule triquetro, foliis reniformi-cordatis, floribus caulinis apetalis. *mirabilis*. LINN. *Sp.* 15. *Viola montana latifolia, flores e radice, semina in cacumine ferens*. DILL. *Elth.* *T.* 303. f. 390. In *Marchia*.

H. *Viola* stipulis crenatis, foliis ovato-lanceolatis. HALL. *G.* 231. c. descr. *Viola palustris, foliis Trachelii glabris*. RVPP. p. 289.

Viola caule erecto, stipulis laciniatis, reliquis longe ellipticis. HALL. *H.* 10. *montana*. LINN. *Sp.* 10. *Viola martia arborescens purpurea*. B. *Pin.* 199. In sylvis *hercynicis*.

Viola

MEIOSTEMONES

His stamina duo, quotquot segmenta floris sint.
Corolla inundibnaiformis. Fr. bacca intra florem.

NYCTANTHES. LINN. 16.

Corollae limbus octopartitus.

H. *Nyctanthes* foliis inferioribus cordatis obtusis,
superioribus ovatis acutis. LINN. Sp. 2. *Sambac*
Jasmin Arabum. ALP. *Aegypt*. p. 73. CLVS. cur.
post. III. T. 3. Fr.

JASMINVM LINN. 17.

Corollae limbus quinquefidus.

H. *Jasminum* foliis oppositis pinnatis. *officinale*,
LINN. Sp. 1. *Jasmin. vulgare*, flore albo. I. R. H.
B. Pin. Fr.

H. *Jasminum* foliis alternis ternatis simplicibus-
que, ramis angulatis. *fruticans*, LINN. Sp. 3.
Jasmin. luteum vulgo dictum bacciferum. B. Pin.
398. Fr.

H. *Jasminum* foliis alternis ternatis pinnatisque,
ramis angulatis. *humile*, LINN. Sp. 4. *Jasmin. humile*
luteum. B. Pin. 397. Fr.

H. *Jasminum* foliis alternis ternatis pinnatisque
ramis teretibus. *odoratissima*, LINN. Sp. 5. *Jasmin.*
Ind. flavum odoratissimum. FERRAR.

H. *Jasminum grandiflorum*

SYRINGA. LINN. 22. Lilac. T.

Corolla limbus quinquefidus. Capsula bilocu-
laris.

Syria-

MEIOSTEMONES

271

Syringa foliis ovato-cordatis, *vulgaris*. LINN.
Sp. 1. *Syringa coerulea flore*. CLVS. *Hist.* p. 56. β.
flore albo. Ad sepes quasi spontanea. Fr.

H. *Syringa* foliis lanceolatis integris dissectisque.
Perfica. LINN. Sp. 2. *Agemlilac Persarum*. COR-
NVT. 189. In *Asia* minori.

OLEA. LINN. 10.

Corollae limbus quadrifidus. In drupa ovata
nux.

H. *Olea* foliis lanceolatis. ^{*europaea*} LINN. Sp. 1. *Olea*.
CLVS. *Hist.* p. 56. BLACKW. T. 213. *syvestris*.
folio duro subtus incano. B. Pin. 472. In *Europa*
australi. Fr.

LIGVSTRUM. LINN. 18.

Corollae limbus quadrifidus. Bacca bilocularis.

^{*in Gaeae*}
Ligustrum. LINN. Sp. 1. HALL. H. 527. BLACKW.
T. 142. *germanicum*. B. Pin. 475. Ad sepes. Fr.

Variae meiostemones ob analogiam relatae sunt
inter didynamias.

DIDYNAMIAE.

Omibus his communia sunt calyx monophyllus
apice incisus, corolla monopetala irregularis bila-
biata. Stamina quatuor duo maiora, duo minora,
rarius alterum par deficit. Stylus simplex. Fructus
in priori classe capsularis, in altera semina quatuor
nuda.

DIDYNAMIAE ANGIOSPERMAE.

Calyx quadri- aut quinquepartitus. Corolla vario modo difformis & bilabiata. Capsula unilocularis aut bilocularis.

MONTIA. LINN. 96.

Corolla quinquefida, stamina tria.

Montia. ^{*fontana*} LINN. *Sp.* p. 87. HALL. *H.* 608. *Montia aquatica minor*. MICHEL. *gen.* 18. *T.* 13. *f.* 2. *Alsinoides annua verna*. VAILL. *Paris.* *T.* 3. *f.* 4. In pratis vdis super *Hardeggen*.

β. Varietas *minor* MICHEL. *T.* 13. *f.* 2. Circa *Herzberg*. *Nordheim* esse *Helv.*

LIMOSELLA. LINN. 694.

Limosella. ^{*aquatica*} LINN. *Sp.* 631. HALL. *Helv.* 609. LIND. *Alf.* 156. *T.* 5. *Plantaginella* HALL. *Jen.* 23. *T.* 6. *f.* 2. *palustris*. B. *Pin.* 190. In inundatis limosis. v. g. vor dem Wehder Thor gleich zwischen den Garten; an dem Lichtenmeer; bey der alten Leinenbrücke. Corollae limbus subaequalis. Plantula per flagella repens cespitosa, foliis longe petiolatis ovalibus.

OROBANCHE. LINN. 697.

Corollae ringens. Habitus aphyllus.

Orobanche caule simplici pubescente, bracteis longioribus, staminibus inclusis. ^{*major*} LINN. *Sp.* 2. HALL. *H.* 611. *n.* 3. *Orob.* I. CLVS. *L.* p. 270. RIV. *Irreg.* *T.* 88. In M. *Altenstollberg*.

SQUAMARIA. HALL. 611. Lathraea.

LINN.

Squamaria, HALL. l. c. RIV. irreg. T. 88. Lathraea caule simplicissimo, corollis pendulis, labio inferiore trifido. ^{Squamaria} LINN. Sp. 4. Orobanche radice dentata maior. C. B. Calyx & corolla quadrifida, Pl. aphylla radice squamata.

LANTANA. LINN. 683.

H. Lantana foliis oppositis, caule inermi ramofo, floribus capitato-vimbellatis. Camara. LINN. Sp. 3. Viburnum Amer. verticæ foliis, lamii odore, floribus miniatis. COMMEL. H. l. p. 151. T. 78. In America calidiore. P.

H. Lantana ~~avulgate~~

VITEX. LINN. 708.

H. Vitex foliis digitatis, spicis verticillatis. Agnus castus. LINN. Sp. 1. RIV. T. 117. Vitex foliis angustioribus cannabis modo dispositis. B. Pin. 475. In Europa australi. Fr.

VERONICA. LINN. 25.

Corolla rotata quadrifida, stamina duo.

H. Veronica spicis terminalibus, foliis quaternis quinisve. ^{Veronica} LINN. Sp. 1. Veron. Virgin. procerior, foliis ternis quaternis & quinis caulem amplexantibus, spicis florum candidissimis. PLUK. T. 70. f. 2. P.

H. Veronica foliis ternis, ovato-lanceolatis, serratis, floribus spicatis. HALL. G. 234. An Veron ^{additionem h. h. de lat. sent. in p. 1. n. 1. a} foliis ternis aequaliter serratis, spicis terminalibus. LINN. Sp. 2. Veron. spicata recta, profunde serrata. J. B. III. p. 284. In Europa australi. P.

H. Veronica foliis ternis, rariter dentatis, spica rariori. HALL. G. I. c. Veron. *spicis terminalibus, foliis ternis inaequaliter seriatis. (maritima.)* LINN. Sp. 3. In maritimis Europae. P.

H. Veronica spicis terminalibus, foliis oppositis crenatis obtusis, caule erecto tomentoso. ^{incana} LINN. Sp.

5. Veronica *spicata lanuginosa & incana, floribus coeruleis.* AMMANN. Ruth. 30. In Vkrainia. P.

Veronica caule simplici, nudo, spicato, foliis oppositis scabris, ovatis, dentatis. HALL. G. I. c. LINN. Sp. 6. Veronica *spicata minor.* B. Pin. 247. Ad arcem Regenshein.

Veronica spicis lateralibus pedunculatis, foliis oppositis scabris ovatis, caule procumbente. HALL. H. 7. *officinalis.* LINN. Sp. 8. Veron. *mas supina & vulgarissima.* B. Pin. 246. BLACKW. T. 143. In sylvis. P.

myrtilloides
ad Bremen
in A. 12
in V. 1. 1. 1.
in A. 12.
Veronica foliis imis cordatis, superioribus angustioribus ellipticis, ex alis racemosa. HALL. H. n. 6. Veron. *supina, facie Teucrii pratensis.* I. R. H. Ad vias. v. g. versus Bovenen. P. *prostrata* L. Sp. 2. 11. *in comitat. Regenshein*

Veronica foliis cordatis, superioribus maioribus, ex alis racemosa. HALL. H. 5. Veron. *racemis lateralibus, foliis ovatis rugosis dentatis, caule debili. (Chamaedrye.)* LINN. Sp. 18. *γ. minor, foliis imis rotundioribus.* I. R. H. In pratis.

H. Veronica foliis cordatis dentatis longe mucronatis, ex alis racemosa. HALL. H. 4. *latifolia.* LINN. Sp. 19. Veron. *prat. omnium maxima.* BUXB. L. T. 34. In Helvetia.

Vero.

Veronica caule debili, foliis ovatis rugosis crenatis petiolatis, racemis lateralibus paucifloris, calyce hirsuto. LINN. *Cent. I. nov. spec. n. 1.* Veron. *procumbens*. RIV. T. 92. HALL. *Gott.* 235. Ad rivum supra Geismar in sylv. *montana* B. & C. *spec.* 21.

Veronica racemis lateralibus, foliis ovatis planis, caule repente. *Beccabunga*. LINN. *Sp.* 14. HALL. *H.* 3. Anagallis *aquatica maior, folio subrotundo*. C. B. BLACKW. T. 48. In fossis. P.

Veronica racemis lateralibus, foliis lanceolatis serratis, caule erecto. LINN. *Sp.* 15. HALL. *H.* 2. Veron. *aquat. maior, folio oblongo*. B. Pin. 252. In fossis. P.

Veronica racemis lateralibus, pedicellis pendulis, foliis linearibus integerrimis. LINN. *Sp.* 16. HALL. *H.* 1. Anagallis *aquatica angustifolia scutellata*. Ad lacum prope Seeburg & alibi.

Veronica foliis subrotundis, crenatis, hirsutis, pedunculis solitariis calyce brevioribus. HALL. *H.* 15. *arvensis*. LINN. *Sp.* 21. Veron. *flosculis cauleculis adhaerentibus*. B. Pin. 250. In pratis & hortis.

Veronica foliis glabris, ovatis, floribus rarius spicatis, sessilibus. HALL. *H.* 14. Veron. *racemo terminali subspicato, foliis ovatis glabris, crenatis*. LINN. *Sp.* 13. Veron. *pratensis Serpyllifolia*. C. B. In gramineis.

Veronica floribus solitariis, foliis cordatis incis-
fis pedunculo brevioribus. *agrestis*. LINN. *Sp.* 20.
HALL. *H.* 18. Veron. *chamaedryos folio*. RIV.
monop. irreg. T. 96. In cultis & agris.

Veronica foliis cordatis quinquelobis, calycis segmentis sagittatis. HALL. H. 17. LINN. Sp. 22. Veron. *Hederae folio*. RIV. T. 96. In agris & hortis.

β. *Veronica Chia verna*, *Cymbalariae folio*, flore albo. HALL. G. 237. BVXB. Cent. I. T. 40.

Veronica foliis trifidis & quinquefidis. HALL. H. 18. Veron. floribus solitariis, foliis digitato-partitis pedunculo brevioribus. *triphyllor*. LINN. Sp. 23. Ver. *rutae folio*. RIV. T. 95. In agris.

* *Veronica australis*

Cymbalaria vulgaris
et *Cymbalaria alba*

CAPRARIA. LINN. 686.

H. *Capraria biflora*. LINN. Sp. 628. foliis alternis, corollis quinquefidis. CLIFF. 320. *Curassavica*. HERM. *Parad.* T. 110. *Gratiolae affinis frutescens Americana*, foliis *Agerati* s. *Veronicae erectae maioris* COMMEL. H. I. p. 79. T. 40. In America meridionali, Fr.

DODARTIA. LINN. 698.

H. *Dodartia* foliis linearibus integerrimis glabris, *orientalis*. LINN. Sp. 1. *Dodartia orient. flore purpurascente*. TOURN. itin. III. T. 208. In Oriente.

MIMVLVS. LINN. 701.

H. *Mimulus*. LINN. Sp. 634. H. Vpf. 176. T. 2. *Lyfimachia galericulata*. GRONOV. Virg. 69. *Digitalis perfoliata glabra, flore violaceo minori*. MORIS. II. S. 5. T. 8. f. 6. In Canada.

BROWALLIA. LINN. 691.

H. Browallia ^{Peruvia} LINN. Sp. 631. H. Cliff. 318. T. 17. H. Vps. 179. In America australi.

RVELLIA. LINN. 702.

H. Ruellia foliis petiolatis, floribus verticillatis subsessilibus. ^{Strepens} LINN. Sp. 1. Vps. 178. Ruellia strepens, capitulis comosis. DILL. Elth. 300. T. 249. f. 321. HALL. G. 304. In Virginia.

H. Ruellia foliis petiolatis, pedunculis longis subdivisis nudis. ^{Flan. de Spina} LINN. Sp. 2. Vps. 179. * Ruellia capsulis teretibus. DILL. Elth. 328. T. 248. f. 320. HALL G. l. c. In Barbados.

BIGNONIA. LINN. 677.

H. Bignonia foliis simplicibus cordatis, caule erecto, floribus diandris. *Catalpa*. LINN. Sp. 1. Bignonia flore sordide albo, intus maculis purpureis & luteis adperso, &c. CATESB. Car. I. T. 49. In Carolina. Fr.

H. *Bignonia* ^{ad. l. 179}

DIGITALIS. LINN. 676.

Digitalis calycinis foliolis ovatis acutis, corollis obtusis. ^{fl. purp.} LINN. Sp. 1. HALL. H. 1. BLACKW. T. 16. Digit. purpurea. DOD. Pempt. 168. LOBEL. ic. 572. In sylvis hercynicis & sylva Solingen. β . flore albo.

Digitalis foliolis calycinis linearibus acutis, flore obtuso venoso, maculoso. HALL. H. 2. Digit. lutea magno flore. TABERN. p. 567. C. B. Pin. I. R. H. In sylva hercynica circa Elbingerode & Herzberg, ^{in sylva hercynica}

H. *Digitalis* foliis calycinis lanceolatis, corollia acutis, labio superiore bifido. ^{lutea} LINN. *Sp.* 2. *Digit. lutea minore flore.* MORIS. II. p. 479. S. 5. T. 8. f. 5. RIV. T. 104. *parva.* LOBEL. *ic. p.* 573. In Gallia. B.

H. *Digitalis* calycinis foliolis ovatis obtusis, corollae labio inferiore longitudine floris. ^{ferruginea} LINN. *Sp.* 3. *Digit. angustifolia, flore ferrugineo.* B. Pin. 244. HALL. G. T. 309. B.

CHELONE. LINN. 666.

H. *Chelone* foliis lanceolatis serratis, summis oppositis. *glabra.* LINN. *Sp.* 1. *Chelone Acadensis flore albo.* TOURN. *Art. Par.* 1706. In Canada. Stamina secunda quatuor, quintum abortivum.

GRATIOLA. LINN. 27.

H. *Gratiola* floribus pedunculatis, foliis lanceolatis serratis. *officinalis.* LINN. *Sp.* 1. p. 17. *Gratiola.* DOD. *Pempt.* 361. 362. BLACKW. T. 411. HALL. H. 618. 619. In paludosis *Europae* australioris. Filamenta duo antherifera, tria abortiva.

ANTIRRHINVM. LINN. 668.

H. *Antirrhinum* foliis oppositis cordatis crenatis. *Asarina.* LINN. *Sp.* 1. *Asarina f. Hedera saxatilis.* LOB. *ic.* 601. P.

Antirrhinum foliis cordatis quinquelobis alternis. ^{Gymnarrhena} LINN. *Sp.* 2. HALL. H. 1. *Cymbalaria.* C. B. CAMER. *Epit.* 860. RIV. T. 85. In rupibus prope *Münden.*

Antirrhinum foliis hastatis imis coniugatis, superioribus alternis, caule flaccido. ^{calcare} LINN. *Sp.* 3.

HALL.

HALL. H. 3. *Elatine folio acuminato*. C. B. 253.
Elatine f. Veronica femina. BLACKW. T. 170. In
 sylva Lith bey Lenglern. A.

Antirrhinum foliis ovatis, imis conjugatis, su-
 perioribus alternis, caule flaccido. LINN. Sp. 4.

HALL. H. 3. *Elatine folio subrotundo*. C. B. *Bey der
 Papiermühle*. A.

H. Antirrhinum foliis ternis ovatis. *triphyllum*.
 LINN. Sp. 6. *Linaria Valentina Hisp.* CLVS. I. p.
 320. *Linaria coerulea apula*. COLUMN. Ecptr. p.
 78. c. ic. Variat coeruleo & luteo flore. In Hi-
 spania. A.

H. Antirrhinum foliis quaternis lanceolatis, caule
 erecto ramoso, floribus pedunculatis. LINN. Sp. 7.
Linaria Americana maxima, purpureo flore. HERM.
 Lugdb. p. 377. RIV. T. 81. In America.

H. Antirrhinum foliis quaternis lineari-lanceola-
 tis, nectariis subulatis recurvis. LINN. Sp. 9. *Linaria
 purpurea maior odorata*. C. B. 1. R. H. *Linaria
 flore purpurea*. RIV. T. 84. In Gallia.

H. Antirrhinum foliis sublinearibus, inferne qua-
 ternis, calycibus capsulae aequalibus. LINN. Sp. 11.
Linaria angustifolia, flore cinereo striato. DILL. Elth.
 T. 163. f. 197. In Anglia.

Antirrhinum foliis sublinearibus, inferioribus
 verticillatis, calycibus piloso-vilcidis, floribus spica-
 tis, caule erecto. *arvense*. LINN. Sp. 12. *Linaria
 arvensis coerulea*. B. Pin. 213, DILL. Elth. T. 163.
 f. 198. In arenosis.

G. *Linaria arvensis*, foliolis carnosiss, flosculis minimis flavis. C. B. l. c. HALL. G. 307.

H. Antirrhinum foliis quaternis subulatis, caulis erectis. *molle*. LINN. Cent. 1. n. 44. *Linaria foliis confertis linearibus carnosiss*. HALL. G. 306. *Linaria maritima foliis succulentis*. BUXB. Cent. 4. p. 23. T. 37. A.

H. Antirrhinum foliis quinis linearibus. *multicaule*. LINN. Sp. 15. *Linaria Sicula multicaulis, molluginis folio*. ROCC. Sic. 38. T. 19. f. 1. HALL. G. l. c. In Sicilia. A.

H. Antirrhinum foliis lanceolatis alternis, caule fruticoso. *Dalmaticum*. LINN. Sp. 18. *Linaria maxima folio Lauri*. I. B. III. p. 458. *Linaria fruticosa di Ragusa, odorata*. ZANON. T. 106. HALL. G. 307.

Antirrhinum foliis lanceolato-linearibus confertis adscendentibus, spicis terminalibus sessilibus, floribus imbricatis. *Linaria*. LINN. Sp. 21. HALL. H. n. 6. *Linaria lutea vulgaris*. B. Pin. 212. BLACKW. T. 115. In gramineis & arvis.

H. Antirrhinum foliis lineari-lanceolatis alternis, floribus racemosis, calycibus corolla longioribus, caule erecto. *Chalepense*. LINN. Sp. 23. *Linaria annua angustifolia, flosculis albis longius caudatis*. TRIVMF. Obs. T. 87. f. 2. In Italia. A.

Antirrhinum foliis lanceolatis obtusis viscidis alternis, caule ramosissimo diffuso. LINN. Sp. 24. HALL. Helv. 7. Antirrhinum *arvense minus*. B. Pin. 212. *Linaria arvensis minima*. RIV. T. 84. In siccis. A.

H. An-

H. *Antirrhinum* corollis ecaudatis spicatis, calycis foliolis brevibus ovatis. *matus*. LINN. *Sp.* 25. HALL. *H.* 9. *Antirrhinum* DOD. *Pempt.* 182. LOB. *ic.* 404. RIV. *T.* 81. A.

Antirrhinum corollis ecaudatis, sparsis, sessilibus, calyce brevioribus. *Orontium*. LINN. *Sp.* 26. HALL. *Hetv.* 8. p. 615. Gott. 308. *Antirrhinum sylvestre*, *Phyteuma*, DOD. *Pempt.* p. 182. In agris.

ACANTHVS. LINN. 711.

H. *Acanthus* foliis sinuatis inermibus ^{*mollis*}. LINN. *Sp.* 1. p. 939. *Mat. med.* 318. *Acanthus mollis Virgilii*. B. *Pin.* 383. RIV. *T.* 86.

JUSTICIA. LINN. 26. Adhatoda. T.

H. *Justicia* arborea, foliis lanceolato-ovatis, bracteis ovatis persistens, corollarum galea concava. *Adhatoda*. LINN. *Sp.* 1. *Adhatoda Zeylanensis*. HERM. *Lugdb.* T. 643. *Ecbolium*. RIV. *T.* 87. Fr.

H. *Justicia* fruticosa, foliis lanceolatis integermis, pedunculis trifloris ancipitibus, bracteis calyce brevioribus. *hyssopifolia*. LINN. *Sp.* 4. In insulis fortunatis. Fr.

SCROPHVLARIA. LINN. 674.

HALL. 618.

Scrophularia caule obtusangulo, foliis cordatis, racemo terminali, calyce tubo floris aequali, galea emarginata. *Scrophularia nodosa foetida*. B. *Pin.* 235. LINN. *Sp.* 2. HALL. *H.* 2. RIV. *T.* 106. CAMER. *Epit.* 866. In fossis. P.

Stro-

H. Scrophularia foliis cordato-oblongis
pendiculatis, racemis terminalibus. *auriculata*
Spec. 4. Betonica aquatica septentrionalium
ic. 133. P.

H. Scrophularia foliis cordatis duplicato-
racemo composito. ^{*Scroph. rodonia*} LINN. *Sp. 5. Scrophula*
rodoniae folio. MORIS. S. V. T. 35. PL
59 f. 5. Scroph. melissae folio. I. R. H. 11
Lusitania. P.

H. Scrophularia foliis cordatis superior
ternis, pedunculis axillaribus bifloris. *pe*
LINN. *Sp. 10. Scrophul. peregrina. CAMEL*
157. T. 43. An Scroph. folio rubro. I.
p. 422. ? A.

H. Scrophularia foliis cordatis acuminat
ribus globosis, squama sub galea destitutis.
H. p 619. HALL. Gott. 311. vernalis. LINN
Scroph. flore luteo. B. Prodr. T. 112. RIV.
Lamium Pannonicum II. CLVS. Hist. II. X
In Italia. Helvetia. P.

G. 311. *Sideritis Sambuci folio*. P. ALP. Exot. p. 202. Flores in hoc genere maximi. In *Hispania*, P.

H. *Scrophularia foliis pinnatis*, calycis segmentis ovatis, flore ventricoso, squamula transversim latiori. *Scrophularia Cretica* L. f. *foliis filicis modo laciniatis*. CLVS. CCIX. An *Scroph. glauco folio*; in *umplas laciniis diviso*. TOURN. Voy. Levant. 1784. Fule hanc speciem descripsi, & a recentioribus neglectam restitui. in *Comment. Gott. Tom. V.* Tuba recta, stigmate hamato, recurvo. *Lucida* *Lin. 2. f. 11.*

H. *Scrophularia foliis pinnatis*, calycis segmentis rotundatis ala alba cinctis, tubo floris recto, squamula lineari. *Scrophularia Ruta canina dicta*. B. Pin. 136. HALL. H. n. 4. LINN. Sp. 9. *Ruta canina*. CLVS. CCIX. Tuba propendens reflexa, stigmate obliquo. In *Helvetia*.

PEDICULARIS. HALL. 620. LINN. 664.

Pedicularis caule ramoso erecto, calycibus bifidis crenatis, corollis labio obliquis. HALL. H. 621. *pratensis*. LINN. Sp. 1. *Pedicularis pratensis rubra latior*. RAIL. Hist. 700. RIV. T. 91. Ad lacum prope *Seeburg*.

Pedicularis caule ramoso procumbente, calycibus quinquefidis crenatis, floribus solitariis labio cordato. HALL. H. 622. *sylvatica*. LINN. Sp. 1. *pratensis purpurea*. B. Pin. 163. In sylvis palustribus.

ALECTO-

ALECTOROLOPHVS. HALL. H. 624.

Rhinanthus LINN. 658.

Alectorolophus corollarum labio superiore compresso, calycibus glabris. HALL. H. 624. I. Rhinanthus. LINN. *Sp.* 3. Pedicularis *pratensis lutea* L. *crista Galli*. B. Pin. 163. In pratis.

Alectorolophus corollarum labio superiore compresso, calycibus hirsutis. HALL. H. 624. n. 2. Pedicularis lutea. TABERN. 791.

MELAMPYRVM. LINN. 660.

Melampyrum spica densa longiori, foliis inter flores laciniatis, adscendentibus. HALL. H. 2. Melamp. *spicis conicis laxis laceris*. (*arvense*.) LINN. *Sp.* 2. Melamp. *purpurascente coma*. B. Pin. 234. In segetibus.

Melampyrum spica densa brevi, foliis inter flores cristatis, reflexis. HALL. H. 625. 1. Melamp. *spicis quadrangularibus compactis, obtusis, imbricatis*. (*cristatum*.) LINN. *Sp.* 1. RIV. T. 80. *luteum angustifolium*. C. B. In M. *Altenstollberg*.

Melampyrum spica laxissima, foliis inter flores laciniatis, patentibus. HALL. H. 3. Melamp. *bracteis cordato lanceolatis, summis coloratis sterilibus, calycibus lanatis*. (*nemorosum*.) LINN. *Sp.* 3. Melamp. *coma coerulea*. C. B. Ad oras nemorum.

Melampyrum floribus sparsis sessilibus, binatis, foliis superioribus incis. HALL. H. 4. Melamp. *florum coniugationibus remotis, crrollis clausis*. (*pratense*.) LINN. *Sp.* 4. Mel. *luteum latifolium*. C. B. In sylvis circa *grossen Schneen* & *Hardeggen*.

Melam-

Melampyrum floribus sparsis, sessilibus, binatis, hiantibus, foliis omnibus indivisis. HALL. H. 5. Melamp. *corollis hiantibus sylvaticum*. In sylvis *Hercynicis*, ~~sylvaticum~~ L. & spec. 6.

ODONTITES. HALL. 627. Euphrasiae.
LINN. Sp.

Odontites foliis inter flores serratis. HALL. H. 1. Euphrasia foliis linearibus, omnibus serratis. ^{Odontites} LINN. Sp. 4. Euphr. *pratensis rubra*. C. B. In pratis & agris humentibus.

EUPHRASIA. HALL. 628. LINN.
659. Sp.

Euphrasia foliis ovatis, acute dentatis. HALL. H. 2. 1. officinalis. LINN. Sp. 2. officinarum. C. B. In pratis.

β. flore minori. In sylvis.

DIDYNAMIAE GYMNASPERMAE. S. VERTICILLATAE.

Caulis quadrangularis. Folia opposita. Flores in verticillos dispositi. Calyx monophyllus, quinquefidus. Corolla irregularis, ut plurimum bilabiata, ringens, labio superiori bifido, inferiori tripartito. Stamina quatuor duo maiora, duo minora. Stylus simplex, stigmate bifido. In fundo calycis semina quatuor nuda. Omnibus vires aromaticae, robustantes,

I. Flore galea destituto.

BVGULA, HALL. 633. *Aiuga*,
LINN. 624.

Labium inferius tripartitum.

Bugula foliis ovatis, inter flores stipatis, flagellis reptans. HALL. H. p. 634. *Aiuga stolonibus reptatricibus*. LINN. Sp. 4. Bugula, DOD. Pempt. 135. In sylvis.

(Bugula foliis angulose dentatis, caule simplici. HALL. H. 635. *Aiuga tetragono-pyramidalis*. LINN. Sp. 3. Bugula montana. RIV. T. 75. In sylvis M. Heymberg & alibi.) ~~Et *Aiuga geminata* L.~~

II. Bugula (*Aiuga*) floribus resupinatis. LINN. Sp. 1. Bugula Orient. villosa, flore inverso coeruleo, alba macula notato. DILL. Elth. T. 43. f. 61. HALL. Goit. 316.

TEVCRIVM, LINN. 625. Chamaedrys.
HALL. 630.

Labium inferius quinquepartitum.

Teucrium foliis multifidis, floribus verticillatis utrinque ternis. LINN. Sp. 3. Chamaedrys foliis pinnatis, pinnis obtusis, trifidis. HALL. H. 633. Iva moschata, folio multifido. RIV. T. 14. In M. arcis Pless.

H. Teucrium foliis trifidis linearibus integerrimis, floribus sessilibus axillaribus. Chamaepitys. LINN. Sp. 5. Bugula foliis trifidis, floribus per alas sparsis. HALL. H. Chamaepitys lutea vulgaris. B. Pin. 249. Sponte in Thuringia. A.

H. Teu

h. *Teucrium* foliis integerrimis oblongo-ovatis petiolatis supra villosis, subtus tomentosis incanis. *Teucrium latifolium*. LINN. *Sp.* 10. *H. Vpf.* 159.

n. 1. *Teucrium fruticans baeticum-ampliore folio*. DILL. *Elth.* 397. *T.* 284. *f.* 367. Folia superne ex atroviridia, lana telae araneae instar obducta, fine obtusiori; inferne cum ramis tomento niveo tecta. Fr.

h. *Teucrium* foliis integerrimis ovatis utrinque acutis, racemis secundis villosis. *Marum*. LINN. *Sp.* 12. *Maum Syriacum*. BLACKW. *T.* 47. Fr.

h. *Teucrium* foliis ovatis acutis subdenticulatis, floribus solitariis. *Marum Hispan. nigrum Piperella dicta*. BOCC. *Mus.* II. *T.* 117. BARREL. *rar.* *T.* 694. Prioris similis, sed folio differunt, & odor ille penetrans & tortis desideratur. Petioli foliis breviores. an Teucrium baeticum?

Teucrium foliis cordatis serratis petiolatis, racemis lateralibus secundis, caule erecto. LINN. *Sp.* 17. *Scorodonia*. RIV. *T.* 12. In sylvis prope *Münden*; in nemore *Solingen*. P.

h. *Teucrium* foliis cordatis crenatis petiolatis, spicis longissimis, densissimis. *Veronica* *HALL. Comm. Gott.* 1752. cum *ic.* *H. Gott.* 315. In *Hyrkania*. P. *Teucrium baeticum* L. *Sp.* 10. *T.* 4.

h. *Teucrium* foliis cordatis undulatis obtuse serratis, floralibus integerrimis concavis, caule fruticoso. *flavum*. LINN. *Sp.* 20. *Teucrium vulgare fruticans* L. CLVS. *Hist.* I. p. 348. *Teucrium* C. B. RIV. *T.* 10. In *Europa australi*.

Teucrium foliis oblongis sessilibus dentato-serratis, ^{foliis} geminis axillaribus, caule diffuso. LINN. *Sp.* 18. *Chamaedrys foliis ellipticis, serratis, molibus, verticillis inter folia paucifloris.* HALL. *H. p.* 631. *Scordium* C. B. BLACKW. *T.* 475. In paludosis. P.

H. *Teucrium* foliis ovatis inciso-crenatis petiolatis, floribus subverticillatis. ^{CHAMAEDRYS} LINN. *Sp.* 19. *Chamaedrys procumbens, foliis solidis, ovatis, rotunda crenatis.* HALL. *H. p.* 630. *Chamaedrys repens.* C. B. RIV. *T.* 10. In *Thuringia*. P.

H. *Teucrium* corymbo terminali, foliis lanceolatis integerrimis. ^{CHAMAEDRYS} LINN. *Sp.* 21. *Chamaedrys foliis angustis integris, ellipticis subtus incanis.* HALL. *H. p.* 632. *Polium matius.* Cordi *Hist.* 124. b. *Ainiga folio integro.* RIV. *T.* 15. In *Thuringia*. P.

H. *Teucrium* *fruticosum.*

Galea trifida.

SCVTELLARIA. LINN. 653. *Cassida.*

Calyx cristatus. Tubus floris incurvus.

Scutellaria floribus cordato-lanceolatis crenatis, floribus axillaribus. *galericulata.* LINN. *Sp.* 4. *Cassida.* HALL. *p.* 635. *Cassida palustris vulgatiores, flore coeruleo.* I. R. H. In aquosis & ad fossas.

H. *Scutellaria* caule ramossissimo, foliis ovato-lanceolatis, racemis axillaribus multifloris. *Cassida latifolia, flore minore.* RIV. *T.* 77. HALL. *G.* 317. *Racemi florum inter folia latent. Folia floralia reliquis similia, floribus majora. Flosculi parvi ex minimis huius generis coerulei.* An *Scutellaria palustris.*

lusa. repens Virginiana maior, flore minore. MORIS.
III. p. 416. ^{latente} LINN. Sp. 3.?

H. Scutellaria foliis subcordatis ferratis, spicis interruptis. *peregrina.* LINN. Spec. 8. H. Vps. 172.
n. I. Cassida COLVMNAE Ecpbr. I. p. 187. T. 189.
HALL. G. I. c. *Lamium astragaloides.* CORNVT.
Canad. T. 129. *Bracteae calycibus minores, ovatae, obtuse serratae, glabrae.*

H. Scutellaria foliis cordatis, crenatis, floribus spicatis secundis, bracteis flore multo minoribus. *Bracteae calycis longitudine, ovatae, integerrimae, pilosae. Calyx fuscus, insigniter cristatus. Flores omnes erecti, secundi, purpurei, labio inferiori albo.* An Cassida Orientalis, altissima, verticillae folio.
I. R. H. Coroll. HALL. G. 318. An Scutellaria foliis subovatis crenatis, florum racemis nudiusculis.
^{Cassida} LINN. Sp. 12. ^{et} *Scut. albigina, L.*

H. Scutellaria foliis cordatis ferratis, verticillis diffitis, bracteis ^{longioribus} superantibus. *Caules diffusi ramosi. Bracteae ovatae, integerrimae, floribus duplo maiores. Calyx parvus, fuscus, pilosus. Floris exiguae galea carnei coloris, hirsuta, barba albida.*

H. Scutellaria foliis subovatis, ferratis, villosis, spicis imbricatis, bracteis flores aequantibus. *Bracteae inferiores foliis similes, superiores floris longitudine ellipticae integerrimae pilosae. Calyx pilosus, insigniter cristatus. Corollae villosae ochroleucae.* An Scutellaria foliis cordatis obtusis, obtuseque serratis, spicis foliosis. ^{Cassida} LINN. Sp. 12. Cassida Cret. fruticosa, folio Catariae, flore albo.
TOURN. Cor. II.

H. *Scutellaria* foliis ovatis crenatis, floribus capitato-spicatis bracteas superantibus. *Spicae breves densae. Bracteae ovato-acutae ciliatae imbricatae; corollae uncialis patulae, tubo ochroleuco, limbo violaceo.* ^{*Scut. lupulina L. var. 3.*} An *Scutellaria* LINN. Sp. 9. *Teucrium alpinum inodorum, magno flore.* B. Prodr. 116.?

H. *Scutellaria* (*Cassida*) *spicis foliosis.* GMELIN. HALL. Gott. 318. c. descr.

H. *Scutellaria* foliis incisus subtus tomentosis. *Orientalis.* LINN. Sp. 1. *Cassida Orient. folio Chamaedryos, flore luteo.* TOURN. Itin. III. T. 306. COMMEL. rar. T. 30. RIV. T. 77.

III. Galea erecta, integra, aut emarginata.

SALVIA. LINN.

Calyx bilabiatus $\frac{1}{2}$. Filamenta duo brevissima, quibus in medio fere, articulatione affixa sunt duo alia longa, quorum extremitati superiori anthera, inferiori glandula aut callosa aut parum farinifera insidet. In omnibus rudimenta duorum staminum sterilium.

I. *Salviae.* TOURN. Calycis & corollae fauces hiantes. Galea recta. Filamenta antherifera fere semicircularia, inferius glandula pulverulenta aucta.

H. *Salvia* foliis lanceolato-ovatis integris crenulatis, floribus secundis subdidynamis, tubo calycem superante. *Salvia officinalis.* LINN. Sp. 4. *Salvia maior. minor. α. non auriculata.* C. B. BLACKW. T. 71. *β. auriculata, sive uno alterove foliolo ad basin folii accessorio.* C. B. I. B. III. p. 105. In Europa australi. Fr.

H. *Salvia* foliis pinnatis, pinnis ovatis serratis, calycibus amplis inflatis, tubum floris aequantibus. *Salvia pinnata*. LINN. *Sp.* 26. *Salvia Orientalis latifolia hirsutissima, viscosa, pinnata, fl. re & calyce purpureis, inodora*. BOERH. *Lugdb.* II. T. 167. In Oriente. Fr.

II. *Sclareae*. Galea magna falcata. Filamenta staminum falcata, glandula callosa.

H. *Salvia* foliis rugosis oblongo-cordatis serratis, bracteis concavis acuminatis coloratis, calyce longioribus. *Sclarea*. LINN. *Sp.* 23. H. *Vpf.* 10. n. 4. Horminum *Sclarea dictum*. C. B. Pin. 235. In Syria.

H. *Salvia* foliis cordato-sagittatis serratis acutis, *glutinosa*. LINN. *Sp.* 19. H. *Vpf.* 11. HALL. H. 639. Horminum *luteum glutinosum*. B. Pin. 238. Horminum *sylvestre* II. CLVS. XXIX. In Helvetia.

H. *Salvia* foliis oblongis dentato-angulatis hirsutis, verticillis lanatis, corollarum labio falcato. *Aethiopiae foliis sinuosis*. C. B. 241. A. LINN. *Sp.* 25.

H. *Salvia* foliis rugosis pinnatifidis lanatis verrucosis. *ceratophylla*. LINN. *Sp.* 24. Horminum *rugoso verrucosoque folio, cornu cervi exprimente*. PLVKN. *Phyt.* T. 194. f. 5. HERM. *Parad.* T. 186. HALL. *Gott.* 323. A.

H. *Salvia* foliis hastato-triangularibus oblongis crenatis obtusis. *Canariensis*. LINN. *Sp.* 20. Horminum *Canariense, hastato folio, caule tomentoso*. PLVKN. *Phyt.* T. 301. f. 2. MORIS. III. 394. S. XL T. 13. f. 17. HALL. G. 321. In Canariis. Fr.

Salvia foliis cordato-oblongis crenatis, verticillis subnudis diffusis, calyce tubo floris brevior. *Salvia pratensis*. LINN. Sp. 11. *Sclarea prat. foliis serratis, flore coeruleo*. I. R. H. *Horminum sylvestre*. OAMER. Epit. 629. Prope Rosdorf. P.
Salvia agrestis var. *horminum* sp. pl. 172 p. 35.

H. *Salvia foliis cordatis acute crenatis, verticillis subnudis remotissimis, calyce aristato*. ^{Indica} LINN. Sp. 17. *Sclarea Indica, floribus variegatis*. I. R. H. HALL. G. 320.

H. *Salvia foliis ovatis vtrinque acuminatis serratis*. *Mexicana*. LINN. Sp. 15. *Sclarea Mexic. altissima, facie Heliotropii*. DILL. Elth. T. 254. f. 330. HALL. G. 320. *Jungia*. HEISTERI Praefat. ad Epist. Burkhardianam. T. I. p. 92.

H. *Salvia foliis cordatis inaequaliter basi excisis, caule nudo, spica florente pendula*. HALL. Comm. Gott. T. I. T. XI. LINN. Sp. 27. *Var. flore albo*. In *Russia*.

III. *Hormina*. Galea brevis, parum erecta. Stamina recta.

H. *Salvia foliis ovatis, crenatis, bracteis cordato-acuminatis sensim decrescentibus, viridibus, in spicam tetragonam digestis*. *Salvia foliis oblongis crenatis, corollarum galea semiorbiculata, calycibus fructiferis reflexis*. LINN. Sp. 7. H. Vps. 11. n. 8. *Horminum coma viridi*. I. R. H. *Calyces fructiferi reflexi, labii superioris dentibus divergentibus*. A.

H. *Salvia foliis obtusis, crenatis, bracteis summis sterilibus maioribus coloratis*. ^{*Horminum*} LINN. Sp. 8. *Horminum*

minum *coma violacea, rubra*. HALL. G. 320. I. R. H. *Calyx priori similis, vix autem reflectitur*. A.

H. *Salvia foliis cordato-sagittatis dentatis, verticillis multifloris, stylo labio inferiori incumbente*. *Salvia verticillata*. LINN. Sp. 18. H. Vps. n. 7. *Horminum sylvestre latifolium verticillatum*. C. B. HALL. Gott. 322. In *Austria, Helvetia*. A.

H. *Salvia foliis cordato-lanceolatis, duplicato-crenatis, undulatis, galea punctata, staminibus galea multo brevioribus*. *Hormini sylvestris quinti altera species*. CLVS. XXXI. *Horminum sylvestre salvisolium maius vel maculatum*. C. B. *Sclarea folio salviae maior vel maculata*. I. R. H. BOERH. Lugdb. II. 165. Hanc cum sequente iungit LINNAEVS Sp. 9. nomine *Salviae foliis cordato-lanceolatis, bracteis coloratis flore brevioribus*. P.

H. *Salvia foliis cordato-lanceolatis, simpliciter ferratis, staminibus galeam aequantibus*. *Hormini sylv. quinti species prior*. CLVS. l. c. *Horminum sylv. f. salvisolium minus*. C. B. RVPP. Fl. Jen. 221. MORISON. III. p. 394. *Sclarea folio Salviae minor f. glabra*. I. R. H. BOERH. l. c. *salvisolium*. p. 12. l. p. 35.

H. *Salvia foliis sinuato-ferratis, hirsutis galea obtusa, stylo galeae longitudine*. *Salvia verbenaca*. LINN. Spec. 13. *Hormini sylv. quarti quinta species*. CLVS. XXXIII. *Horm. sylv. lavendulae flore*. C. B. 329. I. R. H. *Horm. sylv. inciso folio, parvo flore, italicum*. BARREL. 220. *Horm. Verbenae laciniis angustifolium*. TRIVMF. Obs. T. 66. HALL. G. 322.

G. *Horminum minus, folio sinuato, flore minori coeruleo.* DILL. HALL. G. l. c. A.

H. *Salvia foliis pinnato-dentatis, asperis, hirsutis, verticillis pilosis, floribus apetalis. Planta a priori omnino diversa, foliis exsuccis, birtis, rugosioribus, terrae appressis, caule simplicissimo piloso, bracteis & calycibus undique hirsutissimis. In calyce iuniori quidem corolla minima emarcida reperitur, quae vix unquam autem adolescit, semina tamen fecunda sequuntur.* A.

H. *Salvia foliis sinuato-serratis glabris, calyce piloso viscido, corolla vix calycem superante. Horminum Virginianum Blattariae folio.* HALL. G. 322. *Calyx latus post florescentiam amplo rictu bians violaceus. Corolla violacea, galea recta brevissima. An Salvia foliis radicalibus lyratis dentatis, corollarum galea brevissima. (lyrata.)* LINN. Sp. 3? *Salvia corollarum labio superiore brevior, fauce patente.* GRONOV. Virg. 8? A.

H. *Salvia foliis ovato-oblongis simpliciter serratis, calyce hirsuto longius aristato, corolla brevissima. An Horminum folio betonicae, flore coeruleo.* BOERH. L. p. 165. *Calyx angustior viridis, neque viscidus. Corolla coerulea, tenuior.*

H. *Salvia foliis ovatis serratis, spicis tetragonis imbricatis, calycibus tridentatis, corollarum galea brevissima. Salvia foliis ovatis basibus mucronato-tuberosis, spicis imbricatis, calycibus trifidis. Hispanica.* LINN. Sp. 16. c. desc.

ROSMARINVS. LINN. 35.

H. *Rosmarinus officinalis*. LINN. *Sp.* p. 23. *Mat. med.* 16. *Rosmarinus hort. angustiore folio*, C. B. In *Europa* australi. Stamina duo, sed simplicia, neque bifurcata. Fr.

MONARDA. LINN. 34.

Galea linearis, longissima. Stamina duo.

H. *Monarda capitulis terminalibus*, caule obtusangulo. *fistulosa*. LINN. *Sp.* 1. *Origanum fistulosum Canadense*. CORNVT. *Canad. T.* 14. RIV. *T.* 61. In *Canada*. P.

H. *Monarda floribus verticillatis*, corollis punctatis. *lanceolata*. LINN. *Sp.* 4. *Clinopodium Virgin. angustifolium*, quovis verticillo duodecim foliolis rubentibus cincto. PLVKN. *Alm.* III. *T.* 24. f. In *Virginia*. P.

ZIZIPHORA. LINN. 33.

H. *Ziziphora capitulis terminalibus*, foliis ovatis. *capitata*. LINN. *Sp.* 1. *Clinopodium fistulosum punilum Indiae occidentalis*, summo caule floridum. PLVKN. *Alm.* *T.* 164. f. 4. In *Syria*. Huic generi calyx tubulosus, corollae limbus parvus.

H. *Ziziphora floribus lateralibus*, foliis lanceolatis. *tenuior*. LINN. *Sp.* 2. HALL. *G.* 324. *Acinos Syriaca*, folio mucronato, capsulis hirsutis. MORIS. III. p. 304. S. XI. *T.* 19. f. 3. 4.

OCIMUM. LINN. 651.

Calyx superne latus, planus. Flores inversi.

H. *Oci.*

H. *Ocimum* foliis ovatis, glabris, calycibus ciliatis. *Basilicum*. LINN. *Sp.* 2. *Ocimum styracis* f. *melissae* odore. BOERH. *Ocimum caryophyllatum*, maximum & minus. C. B.

β. *latifolium maculatum vel crispum*. C. B.

γ. *viride foliis bullatis*. C. B. A.

H. *Ocimum* foliis ovatis integerrimis. *minimum*. LINN. *Sp.* 3. *Ocimum minimum*. C. B. MORIS. III. p. 407. S. XI. T. 10. f. 17. A.

BRVNELLA. LINN. 654.

H. *Brunella* foliis imis integris, superioribus laciniatis. HALL. H. 637. *Brunella foliis ovato-oblongis petiolatis*. LINN. *Sp.* 7. γ. *laciniata*. *Brunella fol. laciniato*. C. B. 261. RIV. T. 30. In *Helvetia*. P.

Brunella foliis dentatis, spica foliis minore. HALL. H. 636. *Brunella* LINN. *Sp.* 1. *Var.* α. *Brunella maior, foliis non dissectis*. C. B. In pratis. P.

Brunella foliis vix dentatis, spica foliis maiore. HALL. l. c. LINN. *Sp.* pr. *Var.* β. *Brunella flore maiore*. RIV. T. 29. Circa *Gescheroder Feld*. P.

LEONVRVS. LINN. 641. *Cardiaca*. TOURN.

Calyx pentagonus, tubulosus. Galea concava, integra. Antherae punctatae.

Leonurus foliis caulinis lanceolatis trilobis. *Cardiaca*. LINN. *Sp.* 1. HALL. H. p. 639. *Cardiaca* 1. R. H. & auct. BLACKW. T. 171. In rudertis.

β. *Car-*

β. *Cardiaca Orient. canescens, folio ramunculi pratensis, flore purpureo, calyce molli.* I. R. H. Hirsutiae calycis & foliorum differre videtur.

H. *Leonurus foliis tripartitis laciniatis, calycibus villosis. Tartaricus.* LINN. *Sp.* 3. *Cardiaca foliis tenuis & profundius incisus, glabra.* AMMANN. *Ruth.* 49. 50.

H. *Leonurus foliis tripartitis multifidis linearibus obtusiusculis. Sibiricus.* LINN. *Sp.* 4. *Phlomis foliis multifidis.* HALL. *G.* 329. *Ballote inodora, foliis Coronopi.* AMM. *Ruth.* 43. *T.* 8. In *Sibiria*.

H. *Leonurus foliis palmatis dentatis, calycibus & caule lanatis. Cardiaca Chinenfis, albis calycibus, aculeatis, semine nigro.* Comm. Soc. Sc. Gott. *T. III. Ballote foliis palmatis dentatis, caule lanato.* LINN. *Sp.* 2.

(*Leonurus (Cardiaca) foliis cordiformibus.* HALL. *H.* 2. *Galeopsis* LINN. *Sp.* 4. *Lamium oblongo folio, flore luteo.* C. B.)

H. *Leonurus hispidus* p. 305.

PHLOMIS. LINN. 642.

Involucrum setaceum verticillis densis subiectum. Priori affinis.

H. *Phlomis foliis subrotundis tomentosis crenatis, involucris lanceolatis. fruticosa.* LINN. *Sp.* 1. *Phlomis fruticosa, Salviae folio longiore.* I. R. H. *Phlomis latifolia capitata lutea grandiflora.* DILL. *Elth.* *T.* 237. f. 306. In *Hispania*. Fr.

H. *Phlomis involucris hispidis subulatis, foliis cordatis scabris caule herbaceo. tuberosa.* LINN. *Sp.*

Sp. 8. *Phlomis foliis cordiformibus galea lacera.* HALL. *Comm. Gott. T. I. T. 12. Enum. Gott. 328.* *Phlomis floribus parvis subrubentibus, villosissimis, bormini foliis.* AMMANN. *Ruth. n. 28. P.*

H. *Phlomis foliis lanceolatis serratis, involucris linearibus nudis, calycibus decagonis decemdentatis.* ^{Leonurus} LINN. *Sp. 12. Leonurus capitis bonae Spei.* BREYN. *Cent. T. 86. Leonurus Afr. frutescens angustifolia, flore longissimo phoeniceo, Leonurus dicta.* MORIS, III. S. XI. T. 19. f. 17.

LAMIVM. LINN. 636.

Lamium foliis dissitis cordatis acutis. HALL. *H. n. 1. Lamium foliis cordatis acuminatis serratis petiolatis.* LINN. *Sp. 1. Lamium album. β. purpureum non foetens, folio oblongo.* C. B. *Archangelica alba.* BLACKW. *T. 33. Ad sepes. Huic generi calyx aristatus, fauces floris spinula terminantur.*

Lamium foliis petiolatis, cordatis, obtusis, in summo caule congestis. HALL. *H. 2. ^{purpureum} LINN. Sp. 2. Lamium purpureum foetidum, folio subrotundo.* C. B. *Vertica iners altera.* DODON. *Pempt. 153. In hortis.*

Lamium foliis superioribus amplexicaulibus obtusis incis. HALL. H. 2. ^{anagallis} LINN. Sp. 3. Lamium minus, folio caulem ambiente. C. B. 231. In hortis.

GALEOPSIS. LINN. 637.

Galeopsis foliis & caule fulcrato, hispidis, flore calyce duplo longiore. HALL. *H. n. 1. Galeopsis internodiis superne incrassatis, verticillis summis sub-*

^{Tetradact}
 2 *subcontiguus*. LINN. *Sp.* 2. var. *α*. *Cannabis spuria*. RIV. T. 31. In sylvis caeduis & tuderibus.

β. flore albo.

γ. flore magnovario colore. *Cannabis spuria flore maiore*. RIV. T. 32. In dumetis um den Landolshäuser Teich.

Galeopsis foliis ovato-lanceolatis, mollissimis & tomentosis. HALL. H. 3. *Galeopsis pr. spec. var. β*.

1. var. flore ochroleuco, barba violacea.

2. flore purpureo.

Galeopsis foliis longe ellipticis subhirsutis. HALL. H. 4. *Galeopsis internodiis caulinis aequalibus, verticillis omnibus remotis*. LINN. *Sp.* 1. *Ladanum segetum*. RIV. T. 24. In segetibus.

Galeopsis Galeobdolon p. 301.

STACHYS. LINN. 638.

Stachys verticillis sexfloris, foliis cordatis petiolatis acuminatis. *sylvatica*. LINN. *Sp.* 1. HALL. H. 1.

Stachys sylvatica RIV. T. 26. *Galeopsis*. BLACKW. T. 84. In sylvis & ad sepes.

... *Stachys verticillis sexfloris, foliis lineari-lanceolatis semiamplexicaulibus*. *palustris*. LINN. *Sp.* 2.

Stachys verticillis spicatis, foliis subhirsutis ellipticis, ad basin emarginatis. HALL. H. 4. *Stachys palustris foetida*. C. B. In agris humectis.

α. *Stachys foliis hirsutis spicam distinguentibus, imis cordiformibus*. HALL. H. 2. *Stachys verticillis multifloris, tomentosis, foliis lanceolato-ovatis basi emarginatis*. LINN. *Cliff.* 301. *alpina*. *Sp.* 3. *Stachys*

chys folio obscure virente, flore ferrugineo. MORIS II. p. 382. T. X. f. 11. In montosis Helvetiae. ^{mont. Plur.}

H. Stachys caule lanato foliis tomentosis albis, oblongo-cordatis, verticillis multifloris, densissimis, HALL. H. 3. *germanica*. LINN. Sp. 4. Stachys *maior Germanica*. C. B. I. R. H. *montana*. RIV. T. 27. ^{Stachys annua} ^{Stachys amensis p. 206.}

BETONICA. LINN. 631.

Betonica foliis, inter verticillos in vertice congestos, brevissimis & integris. HALL. H. 3. *officinalis*. LINN. Sp. 1. Betonica BLACKW. T. 46. *vulgaris purpurea*. C. B. I. R. H. In sylvis. Ch. gen. Galea plana erecta.

Betonica foliis petiolatis, crenatis, subglabris, bracteis inter verticillos diffitos longioribus, ellipticis, ferratis. HALL. H. 1. Betonica ^{caum.} *verticillata*, ^{Stachys annua 2 spec. 9.} *calycibus spinosis*. LINN. Sp. 4. Var. α . *Sideritis arvensis*, *latifolia*, *glabra*. C. B. Inter fegetes.)

Betonica foliis sessilibus, crenatis hirsutis, bracteis inter verticillos diffitos brevibus, aristatis, integris. HALL. H. 2. ^{hirsuta} LINN. Sp. pr. var. β . *Sideritis vulgaris hirsuta*. C. B. *vulgaris* I. CLVS. H. XXXIX. Vtriusque diversitatem specificam fusius dixi in COMM. Gott. T. V. *Stachys recta* L. Mant. p. 82.

SIDERITIS. LINN. 632. Cunila

HALL. 334.

Corolla subaequalis, calyce parum longior, virgineis, stamina intra tubum floris.

Sideritis (*Stachys*) foliis oblongis acutis amplexicaulibus ferratis: floralibus cordatis acutis integerrimis.

gerrimis. ROY. *Lugdb.* 318. HALL. G. 327. *Sideritis perfoliata.* LINN. *Sp.* 4. *Sideritis orient. phlo-
midis folio.* 1. R. H. *Corollae albae venis rufis.*

H. *Sideritis fruticosa tomentosa, foliis cordatis,
pedunculis ante florescentiam nutantibus.* *Candrien-
sis.* LINN. *Sp.* 1. *Stachys Canariensis frutescens.*
Verbasci folio. COMMEL. H. II. p. 197. T. 99.
HALL. G. 327. In *Canariis.* Fr.

H. *Sideritis fruticosa, tomentoso-lanata, foliis
lineari-lanceolatis.* *Syriaca.* LINN. *Sp.* 3. *Cunila
foliis ovatis tomentosis.* HALL. G. 335. *Sideritis
Cretica tomentosa, candidissima, flore luteo.* 1. R. H.
In *Creta.* Fr.

H. *Sideritis foliis lanceolatis subdentatis, bracteis
cordatis dentato-spinosis, calycibus aequalibus.*
birsuta. LINN. *Sp.* 8. *Sideritis foliis ellipticis, cre-
natis, disiunctis verticillis.* HALL. *Helv.* p. 647.
Sideritis Monspelica scordioides. LOBEL. *Adv.* 424.
Vide characterem in *Obs. botan. epist. ad Ill. WERL-
HOFIVM.* In *Gallia Narbonensi.* P.

H. *Sideritis herbacea decumbens, calycibus spi-
nosis, labio superiori trifido.* *montana.* LINN. *Sp.* 5.
Cunila calycibus paucis aristatis. HALL. G. 335.
*Sideritis montana, parvo flore nigro purpureo, ca-
pite medio croceo.* COLUMN. *Ephr.* I. T. 196. In
Italia. A.

H. *Sideritis (Cunila) foliis ovatis dentatis, ver-
ticillis multifloris, calycibus sessilibus spinosis.*
HALL. G. 334. *Leonurus.* LINN. *Sp.* 2. *Marrubia-
strum Cardiacae folio viridi.* BOCC. *Mus.* II. T
98. *Creta Wistebergam.* A.

— Sideritis [foliis cordatis oblongis crenatis, verticillis sexfloris, staminibus tubo florispaulo maioribus. Cunila *foliis ovatis, dentatis, hirsutis*, HALL. G. 334. (*Glechoma arvensis*. LINN. Sp. 2.) *Stachys minima*. RIV. T. 27. In collibus inter segetes. *Stachys arvensis* Linn. 2^a spec. p. 814.

IV. Galea profunde fissa.

BALLOTE. LINN. 639.

Ballote foliis cordatis indivisis serratis. *nigra*. LINN. Sp. 1. *Ballote* HALL. H. 648. BLACKW. T. 136. *Marrubiastrum*. TABERN. 506. RIV. T. 65. Ad sepes.

ⁿ *Ballote lanata* p. 301.

MARRUBIUM. LINN. 640.

I. Calycibus quinquedentatis.

H. *Marrubium* foliis cuneiformibus, verticillis involucro destitutis. ^{Alysson} LINN. Sp. 1. *Marrubium foliis cordatis, retusis, serratis*. HALL. G. 336. *Alysson Galeni*. CLVS. II. XXXV. In Hispania.

H. *Marrubium* foliis ovato-lanceolatis serratis, calycum denticulis fetaceis. *peregrinum*. LINN. Sp. 2. *Marrub. foliis ellipticis serratis venosis, calycibus quinquedentatis*. HALL. G. 336. *Marrubium Creticum*. TABERN. 539. In pago Erdeborn.

H. *Marrubium* foliis oblongis obtusis, mollibus, crenatis, calycum denticulis subulatis. *Marrubium Hisp. supinum, foliis sericeis, argenteis*. DILL. *Elth. F.* 215. nostram optime exprimit. A priori differt,

fert, foliis & caule lanatis, verticillis densioribus.

Marrubium dentibus calycinis setaceis vncinatis.
^{ru. gae} LINN. Sp. 5. Marrubium *album vulgare*. CLVS. II.
 XXXIV. C. B. 1. R. H. In ruderatis v. g. ad arcem
Pless.

β. Marrubium *album villosum*. C. B. Prodr. 110.
 Pin. 230. 1. R. H. 192. egregia varietas, caule &
 inferna parte foliorum denso tomento obductis,
 odore suavi, aromatico.

h. Marrubium foliis cordatis, serratis, tomentosis. HALL. G. 336. Marr. *calycum limbis patentibus, denticulis acutis*. LINN. Sp. 7. Marr. *album rotundifolium Hispanicum*. HERM. Par. T. 201. In Hispania. P.

h. Marrubium foliis ovatis tomentosis integerrimis. HALL. G. 1. c. Marrub. *calycum limbis planis villosis, foliis cordatis, caule fruticosa*. LINN. Sp. 8. *Pseudodictamnus*. DOD. Pempt. 281. In Creta. Fr.

NEPETA. LINN. 629. Cataria HALL.
 649. TOVRN.

Nepeta floribus interrupte spicatis, verticillis subpedicellatis, foliis petiolatis cordatis dentato serratis. ^{Cataria} LINN. Sp. 1. Cataria HALL. H. 1. Herba Cataria MATH. 719. Nepeta f. *Menba cataria*. BLACKW. T. 455. In ruderosis.

h. Nepeta floribus paniculatis, foliis cordatis petiolatis obsolete crenatis. *Pannonica*. LINN. Sp. 2.

Cataria angustifolia maior. 1. R. H. *Calamintha montana praealta*, *Pulegii* odore, BOCC. Mus. II. ic. 38. In *Pannonia*.

Nepeta virginica

H. *Nepeta* (*Cataria*) foliis ovatis, ferratis, hirsutis, HALL. G. 337. *Cataria peregrina latifolia*. LOBEL. 511.

H. *Nepeta* foliis petiolatis cordato-oblongis dentatis, verticillis pedunculatis dilute spicatis. LINN. Sp. 3. HALL. H. 2. *Cataria Hispanica Betonicae folio angustiori*, flore coeruleo. 1. R. H. In *Helvetia*. P.

H. *Nepeta* floribus spicatis sessilibus, bracteis ovatis coloratis verticillos excipientibus, foliis sessilibus. LINN. Sp. 8. *Nepeta tuberosa*, *spicata*, *purpurea*, *Hispanica*, BOCC. Mus. T. 36. In *Hispania*. P.

H. *Nepeta* caule acuto quadrangulo, spicis densissimis nudis. HALL. G. 338. *Hyssopus* LINN. Sp. 3. *Betonica maxima*, *Scrophulariae folio*, floribus incarnatis. HERM. Par. p. 106. c. ic. Flores iuniores ochroleuci, aetate rubescunt. In *Virginia*.

H. *Nepeta* (*Cataria*) floribus inversis, HALL. G. 338. Comment. Gott. 1752. *Hyssopus corollis transversalibus*. LINN. Sp. 2. In *Siberia* & *China*.

H. *Nepeta latifolia*

HYSSOPVS. LINN. 628. HALL.

H. *Hyssopus* spicis secundis. *officinalis*. LINN. Sp. 1. *Hyssopus officinarum coerulea* f. *spicata*. C. V. 217. BLACKW. T. 296. β. flore albo. In rupibus arcis *Salz der Helden* quasi sponte.

H. *Hyssopus nepetoides*

H. *Hyssopus lephantus*

DRACO-

DRACOCEPHALVM. LINN. 648.

Moldavica HALL.

H. Dracocephalum floribus spicatis, foliis lanceolatis ferratis. ^{Virginiana} LINN. *Sp.* 1. BREYN. *ic.* 33. *T.* 27. RIV. *T.* 25. Pseudodigitalis *Persicae foliis*. BOGG. *Sic.* 12. *T.* 6. *f.* 3. In *Virginia*.

H. Dracocephalum floribus spicatis, foliis compositis. ^{Canariensis} LINN. *Sp.* 2. Moldavica *foliis ternatis*. HALL. *G.* 331. Cedronella *Canariensis, viscosa, foliis plerumque ex eodem pedunculo ternis*. COM-MEL. *Amst.* II. *T.* 41. Dracocephalo *affinis, Americana trifoliata, Terebinthi odore*. VOLKAM. *Norrib.* *T.* 145. In *Canariis*.

H. Dracocephalum foliis ex lanceolato-linearibus rarius dentatis spinulosis. *Comm. Gott.* III. *p.* 436. Dracoceph. *floribus oppositis, bracteis lanceolatis integerrimis, foliis lanceolatis, mucronatis, dentatis*. ^{Ungarica} LINN. *Cent.* II. *Sp.* nov. n. 160. In *Sibiria*.

H. Dracocephalum floribus spicatis, foliis bracteisque lanceolatis indivisis muticis. ^{Russica} LINN. *Sp.* 5. *H. Vps.* *p.* 156. Ruyschiana *glabra foliis integris*. AMMANN. *Ruth.* 50. In *Sibiria*. P.

H. Dracocephalum floribus verticillatis, bracteis lanceolatis, ferraturis capillaceis. ^{Moldavica} LINN. *Sp.* 7. Moldavica *foliis ellipticis, profunde crenatis*. HALL. *G.* 331. Melissa *Moldavica*. CAMER. *Epit.* 576. Melissophyllum *Turcicum*. LOB. *adv.* 220. Melissa *Turcica offic.* In *Moldavia*. A.

H. Dracocephalum floribus verticillatis, bracteis oblongis, ferraturis spinosis, foliis subtomentosis.

canescens. LINN. Sp. 8. H. Vpf. 66. *. Moldavica orientalis betonicae folio, flore magno violaceo, COMMEſ. rar. T. 28. TOVRN. Cor. p. 11.

β. Eadem flore magno albido. TOVRN. l.c. p. 11. Sideritis annua flore luteo, utriculis & foliis longioribus. MORIS. III. p. 389. S. XI. T. 8. f. 18. In Oriente.

h. Dracocephalum floribus verticillatis, bracteis orbiculatis, serrato-ciliatis. *peltatum*. LINN. Sp. 9. H. Vpf. 166. *. Moldavica foliis ovatis, crenatis, floralibus orbiculatis creberrime ciliatis, peramplis. HALL. G. 332. Moldavica orient. salicis folio, flore coeruleo. I. R. H. Cor. A.

h. Dracocephalum floribus verticillatis, bracteis oblongis ovatis integerrimis, corollis ^{mutatis} maiusculis purantibus. LINN. Sp. 10. H. Vpf. 167. n. 6. Moldavica foliis glabris ovatis, floralibus lanceolatis, floribus calyce duplo longioribus. HALL. G. 333. Moldavica Betonicae folio, floribus minoribus coeruleis pendulis. AMM. Ruth. 44. In Sibiria.

h. Dracocephalum floribus verticillatis, bracteis oblongis ^{longioribus} integerrimis, corollis vix calyce maioribus. LINN. Sp. 11. H. Vpf. 167. n. 7. Moldavica foliis ovatis, floribus calycem vix superantibus. HALL. G. 331. Moldavica Betonicae folio, floribus minimis pallide coeruleis. AMMANN. Ruth. 46.

h. Dracocephalum. (Moldavica) foliis cordato-triangularibus, serratis corymbis geminis pedunculatis axillaribus. HALL. G. 333. Comment. Gatt. 1752. ^{Sibirica} Nepeta ~~Sibirica~~. LINN. Sp. 11. Cataria

montana, foliis veronicae pratensis. EVXB. Cent. III.
p. 27. T. 50. f. 1. In Sibiria.

PRASIVM. LINN. 655.

H. *Prasium* foliis ovato-oblongis ferratis. *maius.*
LINN. Sp. p. 601. *Teucrium fruticans, amplo &*
albo flore, italicum. BARREL. ic. 895. *Lamium*
arboreum perenne Creticum. ZANON. Hist. 112. T.
46. In Italia. Fr.

MOLVCELLA. LINN.

H. *Molucella* calycibus quinquedentatis, denti-
culis aequalibus. *laevis,* LINN. Sp. 1. *Molucca lae-*
vis. DOD. Pempt. 92. *Melissa Moluccana odorata.*
C. B. In Syria. A.

H. *Molucella* calycibus septemdentatis. *spinosa,*
LINN. Sp. 2. *Melissa Moluccana spinosa.* DOD. l. c.
HALL. G. 339. A.

MELITTIS. LINN. 650.

H. *Melittis. Melissophyllum.* LINN. Sp. 597. *Me-*
lissa. HALL. H. 650. *Melissa Fuchsi.* CAMER.
Hort. p. 99. n. 30. LOBEL. ic. p. 515. *Lamium*
Pannonicum I. CLVS. XXXVII. In montosis Germa-
niae, Helvetiae. P.

HORMINVM. LINN. 649.

H. *Horminum* foliis cuneiformi-oblongis, caule
bifolio. *Virginicum.* LINN. Sp. 2. *Melissa atroru-*
bens, bugulae folio. DILL. Elth. T. 175. f. 216.
in Virginia. Nobis semper apetalis visa, semina ta-
men perficit. A.

MELISSA. LINN. 647. Calamintha.

HALL.

H. Melissa floribus ex alis inferioribus subsessilibus.
 LINN. *Vps.* 163. *Mat. med.* 294. *officinalis*. LINN.
Sp. 1. Melissa BLACKW. *T.* 27. *bortensis*. B. *Pin.*
 229. P.

^{51. Calamintha}
 H. Melissa foliis ovatis obtusis, caule procum-
 bente. HALL. *H. n.* 2. Calamintha *pulegii* odore.
 B. *Pin.* 228. Calamintha *mont. praealta*, *pulegii*
odore. BOCC. *Mus.* II. p. 45. *T.* 40. In Anglia.
 Belgio. P. *Nel. Nepeta* Linn. 2. *per* *

H. Melissa racemis terminalibus, pedunculis so-
 litariis brevissimis. *Cretica*. LINN. *Sp.* 5. Calamin-
 tha *incana*, *ocymi foliis*. C. B. 569. Calamintha
Pulegii odore, flore minori, erecta, *Virginiana*.
 HALL. G. 339.

CLINOPODIVM. HALL. 653.

LINN. 644. *Thymus*

Clinopodium verticillis densissimis, dissitis, fo-
 liis tomentosis ovatis acuminatis. HALL. *H.* 653. *n.* 1.
 LINN. *Sp.* 1. Clinopodium *Origano simile*. C. B. 224.
 I. R. H. Circa Lützelengen.

Clinopodium verticillis paucifloris, ab imo ad
 summum dispersis. HALL. *H.* 654. *n.* 3. *Thymus*
 LINN. *Sp.* 4. Acinus RIV. 47. Clinopodium *ar-*
vense, *Ocymi facie*. C. B. 225. *vulgare*. LOB. *ic.*
 506. In collibus lapidosis. v. g. circa *Ples*.

H. Clinopodium ¹⁶⁴⁷subrotundo-acuminatis nervo-
 sis subserratis, caule procumbente, floribus ver-
 ticil-

ciliatis. Clinopodium perenne, *Pulegii* odore, *Majoranae* folio. BOCCON. HALL. Gott. 341. *Thymus vulgaris* Willd. *an Candia Thymus* 1222 p. 117.

THYMVS. LINN. 646. Serpillum.

HALL. 654.

H. Thymus erectus, foliis revolutis, ovatis, floribus verticillato-spicatis. LINN. Sp. 2. Thymus vulgaris. RIV. T. 41. & officinarum. Fr.

Thymus floribus capitatis, foliis planis obtusis basi ciliatis, staminibus tubum floris excedentibus. Serpillum HALL. H. 1. Thymus LINN. 1. β. Serpillum vulgare maius. In apricis glareosis. Fr.

Thymus floribus capitatis, foliis planis obtusis basi ciliatis, staminibus in tubo floris latentibus. Serpillum HALL. H. 3. Thymus LINN. Sp. α. Serpillum vulgare minus. Similibus locis. Flores duplo minores ac in priori. Fr.

β. Serpillum foliis Citri odore.

Thymus foliis ellipticis & caule hirsutis. HALL. H. 2. Serpyllum saxatile hirsutum, thymifolium, incanum flore rubello & flore albo. VAILL. T. XXXI. f. 40. Serpyllum Pannonicum III. CLVS. Pann. p. 626. In fabuletis Cellensibus. Fr. v. Willd. Sp. 1. p. 57.

H. *Thymus Asiaticus* p. 514. *Thymus Arvensis*.

SATVREIA. LINN. 626.

H. Satureia pedunculis bifloris. *hortensis*. LINN. Sp. 7. Mat. med. 284. Satureia RIV. T. 44. *sativa*. 1. B. III. p. 272. In Italia, Gallia. A.

H. Satureia pedunculis lateralibus solitariis, trifloris, foliis carinatis, punctatis, ciliatis, mucronatis.

natis. Caulis ramosissimus. Calyx angulatus, bilobatus, ore ciliato, labio superiori trifido, inferiori bifido. Bracteae calycis longitudine. Flos magnus. Segmenta labii inferioris subaequalia, labium superius minus vix emarginatum. *Satureia montana* L.

H. *Satureia* pedunculis lateralibus solitariis multifloris, foliis ovato-lanceolatis, ciliatis, margine reflexis. Calyx cylindricus incanus, segmentis setaceis ciliatis subaequalibus erectis, fauce villis clausa. Stamina tubo floris longitudine. (An *Satureia montana*, LINN. Sp. 6. perennis. RIV. T. 41.?)

H. *Satureia* verticillis lanuginosis, dentibus calycinis setaceis pilosis. *Mastichina*. LINN. Sp. 4. *Thymus* Vps. 2. *Thymbra* Hisp. maioranae folio. 1. R. H. 197. *Samolus* f. *Marum mastichen redolens*. B. Pin. 224. *Thymus mastichina* Lm. p. 8. 222.

H. *Satureia* foliis ovato-lanceolatis, nervosis, capitulis terminalibus. Habitus omnino *Satureiae Virginianae*. LINN. Sp. 1. HERM. Par. 218., nisi huic folia lineari-lanceolata pingerentur. Caulis & rami brachiati. Capitula densa terminalia, foliolis succincta, densissima, plana. Calyx tubulosus, ore ciliato, quinquedentato, subaequali. Floris albi labium superius subintegrum, rectum, segmenta labii inferioris subaequalia, medium inflexum, punctis purpureis adpersum. Stamina quatuor minima in tubo floris latent vix conspicua. Odor fortis, ferè *Pulegii*.

LAVANDULA. LINN. 630.

H. *Lavandula* foliis lanceolatis integris, spicis nudis. *Spica*. LINN. *Sp.* 1. *α.* *Lavandula angustifolia*. C. B. 216. BLACKW. T. 294.

β. *Lavandula latifolia*. C. B. BLACKW. T. 295. In *Europa* australi. Fr.

H. *Lavandula* foliis duplicato-pinnatifidis. LINN. *Sp.* 2. Lav. *folio dissecto*. C. B. *Pin. l. c.* *Lavand. multifido folio*. CLVS. *Hist.* I. 345. In *Hispania*. Fr.

H. *Lavandula* foliis serratis, spicis soliosis. HALL. G. 342. LINN. *Sp.* 3. *Stoechas folio serrato*. C. B. *Stoechas II. crispo folio*. CLVS. H. I. 346. *Stoechas serratifolia*. RIV. T. 55. In *Hispania*.

H. *Lavandula* foliis lanceolato-linearibus, spica comosa. LINN. *Sp.* 4. HALL. G. I. c. *Stoechas Arabica*. RIV. T. 55. *purpurea*. C. B. I. c. CLVS. I. 344.

ORIGANVM. LINN. 645.

Origanum spicis subrotundis paniculatis conglomeratis, bracteis calyce longioribus ovatis. *vulgare*. LINN. *Sp.* 7. HALL. H. 656. *Origanum sylv. f. vulgare*. TABERN. 346. BLACKW. T. 280. *Stramina ex tubo floris eminent*. Ad sepes & in collibus.

H. *Origanum* spicis longis pedunculatis aggregatis, bracteis longitudine calycum. LINN. *Sp.* 6. *Origanum straminibus in tubo floris latentibus*. HALL. G. 343. *Orig. Heracleoticum*. TABERN. 345. LOB. ic. 492. In *Europa* australi.

DICTAM-

(DICTAMNVS. TOVRN.)

H. Dictamnus Creticus. HALL. G. 343. LOBEL. ic. 502. B. Pin. 222. *Origanum foliis inferioribus tomentosis, spicis nutantibus*. ^(dictamnus) LINN. Sp. 2. Bractæe in spica laxa imbricata magnae coloratae. Calycis labium inferius vix ullum, superius ovatum integrum. Fr.

(MAIORANA. TOVRN.)

H. Maiorana. HALL. G. 343. BLACKW. T. 319. RIV. T. 58. vulgaris annua. β. perennis. *Origanum foliis ovalibus obtusis, spicis subrotundis compactis pubescentibus*. ^{Maiorana} LINN. Sp. 10. Bractæae decolores in spicis densis imbricatis tetragonis, calyx bifolius.

MENTHA. LINN. 633.

H. *Mentha floribus spicatis, foliis cordatis dentatis undulatis sessilibus*. ^{crispa} LINN. Sp. 1. *Mentha crispa Danica*. MORIS. III. S. XI. T. 6. f. 5. *Mentha crispa*. RIV. 47.

Mentha foliis elliptico-lanceolatis, verticillis in spicam nudam continuatis. HALL. H. n. 2. *Mentha spicata*. ^{sylvensis} LINN. Sp. 2. β. *longifolia*. *Mentha sylvestris longiore folio*. C. B. *Menthastrum*. LOBEL. ic. 509. BLACKW. T. 292. In aquosis vbique.

H. *Mentha foliis ovatis tomentosis, verticillis spicatis, staminibus tubo floris aequalibus*. *Mentha* HALL. H. 1. *Mentha sylv. rotundiore folio*. I. R. E. LINN. Sp. 2. In fossis *Alsatiae*. *Mentha rotundifolia* 2 spec. 3.

H. Men-

T. 21. *Marrubium palustre glabrum*. C. B. Ad fossas.

β. *Marrubium palustre hirsutum, tenuius laciniatum*.

AMETHYSTINA. HALL. LINN. 32.

H. *Amethystina* *Comment. Gott. T. I. Tab. X. p. 209. Act. Vps. 1742. p. 51. f. 1. Amethystea*. LINN. *Sp. p. 21. H. Vps. 9. Amoen. Acad. 386. * Amethystina montana erecta, foliis exiguis digitatis trifidis ferratis, strobulis cum coma e coerulesco-janthinis*. AMM. *Ruth. p. 54. n. 70. In Sibiria. A.*

VERBENA. LINN. 30.

Verbena tetrandra, spicis filiformibus paniculatis foliis multifido-lacinatis, caule solitario. *officinalis*. LINN. *Sp. 13. Verbena foliis lacinatis, superioribus tripartitis, spicis angustis longissimis*. HALL. *Helv. 661. Verbena communis coerulesco flore*. C. B. 1. R. H. BLACKW. *T. 41. Vbiq. ad vias. A.*

H. *Verbena tetrandra*, spicis filiformibus, foliis multifido-lacinatis, caulibus numerosis. *spuria*. LINN. *Sp. 12. H. Vps. 8. Verbena urticae folio, Canadensis, foliis incis, flore maiora*. 1. R. H. HALL. *G. 345. In Canada. R.*

H. *Verbena* spicis filiformibus paniculatis, foliis ovato-lanceolatis ferratis petiolatis. *urticifolia*. LINN. *Sp. 11. HALL. G. 346. Verbena recta Canadensis, urticae folio*. MORIS. *III. p. 418. S. XI. T. 25. f. 3. ZANON. 227. T. 175.*

H. Ver-

H. Verbena foliis ovatis ferratis sessilibus, spicis filiformibus paniculatis, floribus minimis. Verbena *Caroliniana*, *Melissae folio*. DILL. *Eleb. T.* 301. HALL. G. l. cit. *Verb. Carolina 2 spec. 12.*

H. Verbena spicis longis acuminatis, foliis ferratis inferioribus hastatis, superioribus lineari-lanceolatis. *hastata*. LINN. *Sp.* 10. Verbena *Amer. altissima*, *spica multiplici*, *verticillae foliis angustis*, *floribus coeruleis*. HERM. *Parad. T.* 242. HALL. G. l. c. P.

H. Verbena spicis fasciculatis, foliis lanceolatis amplexicaulibus. *Bonariensis*. LINN. *Sp.* 9. Verbena *Bonariensis altissima*, *Lavendulae Canariensis folio*, *spica multiplici*. DILL. *T.* 300. HALL. G. 347.

H. Verbena spicis capitato-conicis, foliis ferratis, caule repente ^{nodiflora}. LINN. *Sp.* 8. Verbena *foliis verticilliter ovatis*, *spicis solitariis ovatis*. ROYEN. 327. Verbena *nodiflora*. C. B. *Pin.* 269. *Prodr.* 125. In *Virginia*.

H. *Verbena mexicana*.
(caule tetragono)

H. *Verb. imariensis*.

TETRADYNAMIAE SILIQVOSAE.

Classi huic naturali calyx est quadrifolius, pene semper caducus, petala quatuor ex longiori vñgue in bracteam reflexa, more caryophyllaeo. Stamina sex, quatuor maiora, duo minora, ad horum basin glandulae virides saepe adsident. Fructus plerumque bilocularis est aut brevis, subrotundus, oblongus, *silicula* dictus, & stylo stigmate obruso instructus; aut est longior, teres, *siliqua* dictus, stigma

te fere seffili. Vires omnibus sunt resolventes, antiscorbuticae, sapor acer, natura alcalescens.

ISATIS. LINN. 738.

Silicula oblonga anceps unilocularis.

H. *Isatis* foliis radicalibus crenatis, caulinis sagittatis, siliculis oblongis. *tinctoria*, LINN. Sp. 1. *Isatis sylvestris*. C. B. BLACKW. 246. Circa *Jenam*. *β. sativa*. Vid. icon. SCHREBERI in proprio libello.

CRAMBE. LINN. 739.

Stamina apice bifido. Capsula rotunda.

H. *Crambe* foliis subcordatis, cauleque scabris. *hispanica*, LINN. Sp. 3. Rapiſtrum *maximum rotundifolium monospermum*. CORNVT. *Canad. T.* 148. MORIS. II. S. III. T. 13. f. 1. A.

BUNIAS. LINN. 737. Erucago.

TOURN. 193.

Silicula irregularis, tetragona aut gibba.

H. *Bunias* siliculis tetragonis; angulis bicristatis. *Erucago*, LINN. Sp. 2. *foliis antrorsum sinuatis*. Cliff. 340. *Erucago Monspel. siliqua quadrangula echinata*. B. Prodr. T. 41. A.

H. *Bunias* siliculis ovatis gibbis verrucosis. LINN. Sp. 3. *foliis retrorsum sinuatis*. Cliff. H. Vpf. 186. *Crambe orient. dentis leonis folio, erucaginis facie*, L. R. H. Cor. 14. P.

H. *Bunias* siliculis ovatis laevibus ancipitibus. *Cakile*. LINN. Spec. 4. *Crambe foliis succulentis lineae-*

linearibus pinnato-dentatis. SAUVAG. *Monsp.* 280.
Eruca marit. Italica, siliqua hastae cuspidi simili.
 B. *Pin.* 99. *Cakile Serapionis.* LOB. *ic.* 223. Ad
 littora maris. A.

ANASTATICA. LINN. 715.

Silicula figura ungulae bovis, stylo medio
 longiori.

H. *Anastatica.* LINN. *Sp.* 641. H. *Vps.* 183.
Rosa Hierobuntica. CAMER. H. T. 41. C. 2. Ad
 littora maris rubri. A.

VELLA. LINN. 714.

Silicula globosa, stylo ovato capsula longiori.

H. *Vella* foliis pinnatifidis, filiculis pendulis. *an-*
nuua. LINN. *Sp.* 2. *Nasturtium sylv. Valentinum.*
 CLVS. CXXX. In *Hispania.* A.

MYAGRVM. LINN. 713.

Silicula difformis bi- aut trilocularis, loculo uni-
 co seminifero, reliquis abortivis.

H. *Myagrvm* filiculis biarticulatis dispermis, foliis
 extrorsum sinuatis denticulatis. LINN. *Sp.* 1. H.
Vps. 182. *. *Myagrvm siliquae primo nodo strigio-*
ri, altero globoso striato. HALL. H. 536. Gott. 239.
Rapistrum monospermum. B. *Prodr.* T. 37. MAPP,
Alsat. T. 266. A.

H. *Myagrvm* filiculis cordatis subsessilibus, foliis
 amplexicaulibus. *perfoliatum.* LINN. *Sp.* 5. H. *Vps.*
 182. *. *Myagrvm filiculis utrinque gibbosis mono-*

Spermis. HALL. G. l. c. *Myagrum monospermum*, B. Prodr. p. 52. Inter segetes *Helvetiae*. A.

H. *Myagrum filiculis globosis compressis punctato-rugosis.* ^{*auriculatis*} LINN. Sp. 7. HALL. H. 537. *Myagrum similis siliqua rotunda*, B. Prodr. T. 52. Circa *Berolinum*. | *inter clypeos v. r. r.*

H. *Myagrum latissimum, inter clypeos et linum v. r. r.*

ALYSSVM. HALL. 537. LINN. 722.

Silicula subrotunda, stylo longiori. Habitus incanus.

(*Alysson foliis scabris sagittatis, leviter dentatis.* HALL. G. 537. *Myagrum* ^{*latissimum*} LINN. Sp. 6. *Alysson segetum, foliis auriculatis, acutis.* I. R. H. *Myagrum l. Camaelina.* Inter linum abunde. A.)

G. *Myagrum (Alyssum) foliis auriculatis, glabris profunde dentatis.* HALL. H. l. c. n. 2. Ad folias.

H. *Alyssum caule erecto, foliis lanceolatis incanis integerimis, floribus corymbosis. incanum.* LINN. Sp. 5. *Alyssum fruticosum incanum.* I. R. H. *Thlaspi incanum Mechliniense.* LOB. ic. 216. CLVS. CXXXII. *Petala bifida.* In arenosis, apricis, v. g. circa *Noribergam, Berolinum, Cellam.*

H. *Alyssum caule erecto herbaceo, filiculis sessilibus ovalibus compresso-planis, petalis acuminatis. clypeatum.* LINN. Sp. 7. *Leucoium alyssoides clypeatum maius.* B. Pin. 201. In *Europa australi.*

H. *Alyssum caule herbaceo, foliis lanceolatis dentatis, filiculis inflatis.* ^{*auriculatis*} LINN. Sp. 8. *Alyssum siliquis inflatis, foliis tomentosis.* HALL. G. 241. A-lyssi-

lystoides incanum, foliis sinuatis. 1. R. H. Eruca peregrina. CLVS. CXXXIV. Petala bifida.

Alyssum foliis imbricatis, lanceolato-ovatis integerrimis, filiculis tetraspermis, calyce persistente. *Clypeola* LINN. *Sp. 11* HALL. *H. 1.* Alyssum minimum. CLVS. CXXXIII. Alyssum incanum luteum, Serpylli folio maius. 1. R. H. A.

H. Alyssum foliis imbricatis lanceolato-ovatis, punctatis, filiculis tetraspermis, calyce caduco. HALL. *H. 2.* Gott. 240. Jonthlaspi luteo flore, incanum, discoides, umbellatum, montanum. COLVMN. Ecphr. T. 280. Omnia maiora, stylus longior. In arenosis ad Rhenum. P.

CLYPEOLA. LINN. 723.

Silicula orbiculata, plana, hilo modice excisa.

H. *Clypeola* filiculis unilocularibus, monospermis. LINN. *Sp. 1.* Thlaspi clypeatum, serpylli folio. LOB. *ic. 214.* Jonthlaspi minimum spicatum lunatum, COLVMN. Ecphr. T. 284.

DRABA. LINN. 717.

Silicula elliptica compressa, stylo minimo.

Draba caulibus nudis, foliis incis. LINN. *Sp. 2.* HALL. *H. 538.* Alysson vulgare, polygoni folio, caule nudo. 1. R. H. SEGV. Veron. I. T. 4. f. 3. Alsine minima. TABERN. 708. Verno tempore in macilentis. A.

LVNARIA. LINN. 725.

Silicula elliptica, plana maxima. Semina reniformia.

β. foliis serratis. Cochlearia Britannica. LOBEL. ic. 294. Ad littora maris septentrionalis.

H. Cochlearia foliis caulinis cordato-sagittatis amplexicaulibus. *glastifolia.* LINN. Sp. 7. Lepidium *glastifolium.* C. B. MORIS. II. S. III. T. 27. f. 34 A.

Cochlearia foliis radicalibus lanceolatis crenatis, caulinis incis. *Armoracia.* LINN. Sp. 6. Cochlearia *folio cubitali.* I. R. H. Raphanus *rusticanus.* C. B. In prato prope Herperhausen. P. *Echid. Danica.*

CORONOPVS. HALL. H. 542.

Silicula transversim lata, verrucosa, loculis monospermis.

Coronopus *Ruellii.* BLACKW. T. 120. Nasturtium *sylvestre, capsulis cristatis.* I. R. H. Cochlearia *foliis pinnatifidis.* LINN. Sp. 5. Hinter der Mäschmühle am Weg gegen den Berg hin, & ad fontes falsos. A.

IBERIS. LINN. 721.

Petala exteriora maiora.

H. Iberis frutescens, foliis cuneiformibus integerrimis, obtusis. *semperflorens.* LINN. Sp. 1. Thlaspi *Persicum.* RIV. T. 110. Thlaspi *fruticosum Persicum folio Cbeiri.* ZANON. T. 165. Fr.

H. Iberis frutescens, foliis linearibus acutis integerrimis. *sempervirens.* LINN. Sp. 2. Thlaspi *fruticosum sempervirens, flore albo Cretici.* BARREL. 734. In Creta.

H. Iberis foliis lanceolatis acutis subdentatis, floribus racemosis. LINN. Sp. 7. Iberis *foliis sursum*

Osyridis folio. C. B. Iberis. BLACKW. T. 312. In muris v. g. Auf der Albaner Kirchhof-Mauer. A.

H. Nasturtium (*Lepidium*) floribus diandris minutissimis, foliis linearibus pinnatis. LINN. *Sp.* 11. Iberis humilior, annua, ramosior, Virginiana. MORIS. II. S. III. T. 21. f. 2. A.

LEPIDIVM. LINN.

Silicula subcordata, apice non emarginata; stylus capitatus.

H. *Lepidium* foliis caulinis pinnato-multifidis, ramiferis cordatis amplexicaulibus integris. LINN. *Sp.* 1. HALL. Gott. 245. Thlaspi verum Dioscoridis. ZANON. T. 129. Nasturtium Orient. foliis inferioribus Millefolii, superioribus Persfoliatae. I. R. H. A.

H. *Lepidium* foliis pulposis ovato-lanceolatis ferratis. LINN. *Sp.* 8. *Lepidium latifolium* C. B. I. R. H. Circa Spiram; in Anglia. P.

H. *Lepidium* foliis lanceolatis amplexicaulibus dentatis. LINN. *Sp.* 14. *Draba umbellata*. C. B. *Draba l. vulgaris*. CLVS. II. CXXIII. In Germania australi. P. *Lepidium sativum*, *ruderale*, *virginianum*.

II. Fructu longiori f. filiqua.

SISYMBRIVM. HALL. 547. LINN. 728.

Erysimum. LINN. 729.

Calyx deorsum non gibbus. Stylus brevissimus.

Sisymbrium filiquis declinatis, foliis pinnatis, foliolis subcordatis. LINN. *Sp.* 1. *Sisymbrium foliis pinnatis, angulosis, floribus super folia vix elatis.*

HALL.

HALL. H. *Nasturtium aquaticum*. C. B. BLACKW. T. 260. Ad fontes. P.

Sisymbrium filiquis declinatis, foliis pinnatis, foliolis lanceolatis ferratis. LINN. Sp. 2. HALL. H. *Sisymbrium palustre repens*, *Nasturtii folio*. I. R. H. *Eruca sylv. luteo flore*. C. B. Ad vias & fossas.

Sisymbrium filiquis declinatis oblongo-ovatis, foliis pinnatifidis ferratis. LINN. Sp. 3. HALL. H. 3. *Raphanus aquaticus*, *foliis in profundas lacinias divisis*. B. Prodr. T. 38.

β. Sisymb. *foliis imis integris ovatis, serratis, superioribus pinnatis*. HALL. H. 2.

γ. Sisymb. *foliis simplicibus dentatis serratis*. HALL. H. 1. *Raphanus aquat. alter*. B. Prodr. 37. Ad rivos & fossas. P.

h. Sisymbrium filiquis axillaribus sessilibus subulatis aggregatis, foliis repando dentatis. LINN. Sp. 5. HALL. H. 10. *Erysimum polyceratum*, *f. corniculatum*. B. Pin. 101.

Sisymbrium petalis calyce minoribus, foliis decomposito-pinnatis. LINN. Sp. 11. *Sisymbrium pinnis foliorum, pinnatis, pinnulis subhirsutis linearibus*. HALL. H. 7. *Nasturtium sylv. tenuissime divisum*. C. B. *Sophia Chirurgorum*. BLACKW. T. 440. Ad muros vias. A.

h. Sisymbrium foliis pinnatis, extrema pinna triangulari dentata, petiolis longis hirsutis. HALL. G. 248. *Sisymbrium foliis pinnato-hastatis, acutis hirtis, caule retrorsum hispido*. LINN. Cent. Pl. I. n. 51. *Erysimum mont. Irionis folio, macrolepto-*

ceraton. COLVMN. Ecphr. I. T. 268. In testis stramineis Eislebiae.

H. *Sisymbrium* foliis pinnato-hastatis flaccidis, foliolis sublinearibus integerrimis, pedunculis laxis, *altissimum. LINN. Sp. 13. Erysimum Monspeffulanum, Sinapios folio. WALTH. Hort. 55. T. 22. Rapistrum sylvestre Irionis folio. COLVMN. Ecphr. I. T. 268. In Gallia. A.*

H. *Sisymbrium* foliis pinnato-hastatis tomentosis, caule laevi, ^{Orientali} *LINN. Cent. Pl. II. n. 173. Erysimum Orient. sonchi folio, flore sulphureo, siliquis longissimis. BOERH. I. p. 14. HALL. G. 248. A.*

H. *Sisymbrium* foliis pinnatis, extrema pinna amplissima, siliquis praelongis. *HALL. G. 247. Hesperis foliis dentato-pinnatifidis, caule laevi, ^{dentato} LINN. Sp. 6. Hesperis flore albo minimo, siliqua longa, folio profunde dentato. BILL. Elch. T. 148. f. 177. In Sicilia. ^{Hesperis maritima. H. ~~et. arvensis~~ 238.}*

Erysimum
Sisymbrium foliis lyratis, pinna extrema amplissima, subrotunda. *HALL. H. 6. Erysimum LINN. 2. Barbarea femina. TABERN. 452. Ad fossas. A.*

Sisymbrium caule ramosissimo, foliis inferioribus angulatis sinuatis, superioribus dentatis, siliquis nodosis. *Sisymbrium foliis imis Barbareae, superioribus integris dentatis. HALL. It. Hercyn. c. ic. Opusc. bot. p. 100. n. XL. Ad rivos bercynicos. Arabis ^{Barbarea} ~~Italiana~~ ^{Linnaeus 2 spec. 30. 1}*

Sisymbrium (Erysimum) siliquis cauli adpressis. HALL. H. 9. LINN. Sp. 1. Erysimum BLACKW. T. 28. vulgare. C. B. I. R. H. Ad vias. A.

Cent. Pl. II. n. 174. Eruca sylvestris, flore albo, Italica. BARREL ic. 132. Caulis retrorsum scaber; pedunculi similes patentes. Calyx & ungues petalorum alborum exsiccati violacei evadunt; stigma bicolle.

SINAPIS. LINN. 735.

Calyx patens. Glandulae virides quatuor. Stylus anceps, siliquae aequalis.

Sinapis filiquis multangulis toroso-turgidis, rostro longioribus. *arvensis*. LINN. *Sp. 1.* HALL. *H. 3.* Sinapi *arvensis praecox*, semine nigro. I. R. H. In agris & hortis. A.

H. Sinapis filiquis retrorsum hispida, apice sub-tetragonis compressis. *orientalis*. LINN. *Cent. Pl. I. n. 55.* Sinapi *Orient. maximum*, rapi folio. I. R. H. Cor. 17. HALL. G. 250.

Sinapis filiquis hispida, rostro obliquo longissimo. *alba*. LINN. *Sp. 2.* HALL. *H. 2.* Sinapi album. BLACKW. T. 29. Sin. *filiqua hirsuta*, semine albo & rufo. I. R. H. Sin. *apii folio*. C. B. In agris & hortis. A.

Sinapis filiquis glabris tetragonis. *nigra*. LINN. *Sp. 3. Mat. med. 326.* HALL. *H. 1.* BLACKW. 466. Sinapi rapi folio. C. B. I. R. H. In ruderalis. Haec species ob usum medicum & dieteticum celebris. A.

H. Sinapis ramis fasciculatis, foliis inferioribus ovatis laceris, summis lanceolatis integerrimis. LINN. *Sp. 4. H. Vpf. 191. **. Sinapi *Ind. maximum*, lactuca

afflucae folio. HERM. Par. T. 230. *Sinapi Chinenſe,*
Acanthi folio. GMELIN. A.

H. *Sinapi* *apiculata* 351.

RAPHANVS. LINN. 736.

Calyx erectus, gibbus. Siliqua acuminata, articulata.

H. *Raphanus* filiquis teretibus torosis biloculari-
bus. *sativus.* LINN. Sp. 1. *Raphanus hortensis,*
BLACKW. T. 81. *oblongus rotundus.* C. B. A.

Raphanus filiquis teretibus articulatis laevibus
unilocularibus. ^{*Raphanistrum*} LINN. Sp. 2. *Raphanistrum arvenſe,*
flore albo, & *luteo.* I. R. H. *Rapiſtrum flore albo,*
luteo. C. B. A.

β. Variet. *flore albo venoſo.* *Lampſana apula* Pli-
nii & *Diſcoridis.* COLUMN. Ecphr. l. p. 261. In
agris auf dem Hagen, und hinter Grohnde. A.

DENTARIA. LINN. 726.

Dentaria foliis inferioribus pinnatis, superiori-
bus ſimplicibus. *bulbifera.* LINN. Sp. 2. HALL. H.
3. *Dentaria heptaphyllor baccifera.* CLVS. CXXI.
In aliſ foliorum profert bulbos nigros congeſtos, qui
de lapſi ſpeciem propagant; ſiliquae raro matureſcunt.
In ſylvis circa Gottingam & bercynicis.

BRASSICA. LINN. 734.

Brassica radice cauleque tenui, foliis caulinis uni-
formibus cordatis ſeſſilibus. *campeſtris.* LINN. Sp.

2. *Brassica arvenſis perfoliata, flore luteo.* Inter fe-
geres. A. — *cap. h2c, 127 Pir. ornata;* *h2c et h2c h2c h2c h2c h2c*

Brassica radice cauleſcente fuſiformi. LINN. Sp. 3.
Napus ſylveſtris, ſativa. C. B. In agris ſeritur. A.

H. Brass

Cardamine foliis pinnatis, pinnis laciniatis. HALL.
H. 1. *impatiens.* LINN. Sp. 8. *Cardamine annua,*
exiguo flore. I. R. H. In sylvis circa *Göttingam* &
Hercyniis.

H. *Cardamine foliis pinnatis, foliolis lanceolatis*
basi videntatis. *Virginica.* LINN. Sp. 14. *Nastur-*
tium bursae pastoris folio, Virginianum, flore albo,
aliqua compressa. FLVKN. T. 101. f. 4. A.

TURRITIS. LINN. 733.

Calyx erectus. Siliquae longissimae, compressae,
cauli parallelae.

Turritis foliis radicalibus dentatis hispidis, cauli-
nis integerrimis, amplexicaulibus glabris. ^{*glabra*} LINN. Sp.
1. HALL. H. *Turritis foliis inferioribus Cichora-*
cis, superioribus Persiliatae. I. R. H. In sylvis. A.

Turriti: foliis omnibus asperis, dentatis, caulinis
amplexicaulibus. ^{*hirsuta*} LINN. Sp. 2. HALL. H. 3. *Ery-*
simo similis hirsuta, non laciniata, alba. BAVH.
Prodr. T. 42. In lapidosis. A.

ARABIS. LINN. 732.

Calycis folia deorsum gibba, in quibus glandu-
lae. Siliquae laxae dispositae.

H. *Arabis folijs amplexicaulibus dentatis, caule*
diffuso ramoso. ^{*arvensis*} LINN. Sp. 1. β. HALL. H. 1. *Dr-*
aba II. GLV. CXXIV. *Draba alba siliquosa.* C. B. In
alpibus helveticis. P.

Arabis foliis ovatis, vix dentatis, scabris, ad
terram confertis. HALL, H. p. 562. *Arabis foliis*
petio-

petiolatis lanceolatis integerrimis. (Thaliana
 LINN. *Sp.* 3. *Pilotella filiquata minor.* TH.
Harc. 84. *T.* 7. *f.* D. In agris sterilibus M. Hagen
 in vallecule inter Waaken & Besigehausen. A.

(H. Arabis (Sisymbrium) foliis lyrato-serratis f
bris radicalibus, caule subnudo ramoso. LINN. Sp.
Sisymbrium HALL. 14. *Eruca coerulea in arena*
proveniens. C. B. I. R. H. In Helvetia. A.)

H. Arabis foliis amplexicaulibus, siliquis anci
 tibus linearibus, calycibus subpilosis. *pendu*
 LINN. *Sp.* 6. *H. Vps.* 191. *. *Turritis latifolia h*
suta, siliquis pendulis. AMMANN. *Ruth.* 58. HALL
G. 253. In Sibiria. A.

Arabis Halleri p. 330.

CHEIRANTHVS. LINN. 730.

Petala calyce gibbo longiora. Siliqua compre
 stigmate bifido.

Cheiranthus foliis lanceolatis, acutis, glabris. Che
 LINN. *Sp.* 2. HALL. *H.* 563. *Leucoium luteu*
vulgare. C. B. I. R. H. Leucoium aureum. MAT
877. Sponte interdum in muris. v. g. bey dem C
servatorio. A.

β. Var. flore pleno. C. B.

H. Cheiranthus caule suffruticoso, foliis den
 congestis, obverse ovatis, lanceolatis, glabris, in
 gerrimis. *Rami sub floribus dense foliis tecti, im*
rius nudi. Calyx ruber. Flores grandes, spec
kermesini coloris, fere unciales. Siliqua viria
compressa, saepe undulata. An Cheiranthus sin
fere nomine HALL. *G.* 255. Videtur ex descripti

ne florum, sed synonyma aliena esse videntur. Optime nostram exprimit *Leucoium hyemale diu durans*, folio viridi & livido glabro. I. B. III. 877. etsi florem album dicat, nec abluderet *Leucoium marinum majus*. CLVS. 298. & *Keiri marin. flore candido*. LOB. ic. 332. nisi CLVSIVS annum illam diceret. Fr.

H. *Cheiranthus* foliis lanceolatis integerrimis obtusis incanis, siliquis apice truncatis, compressis, caule suffruticoso. LINN. Sp. 4. *Leucoium incano folio, hortense*. C. B. flore albo, purpureo, violaceo, pleno. *Viola alba & purpurea*. LOB. ic. 329. Fr.

H. *Cheiranthus* caule ramoso diffuso, foliis lanceolato ovatis retusis, raris, integris, siliquis apice subulatis. An *Cheiranthus Chius*. LINN. Sp. 3. *Leucoium thlaspioides facie, flore albo*. HERM. Parad. T. 193. optime nostram exprimit. Minus convenit *Hesperis siliquis hirsutis, flore parvorubello*. DILL. Elth. T. 147. f. 178. A.

H. *Cheiranthus* filicularum apicibus tridentatis. LINN. Sp. 9. *Leucoium marinum*. GAMER. hort. 87. T. 24. MORIS. II. p. 242. S. III. T. 8. f. 13. In maritimis. A.

Cheiranthus sylvestris 328.

HESPERIS. LINN. 731.

H. *Hesperis* caule ramoso, foliis lanceolatis, dentatis scabris, corollis minimis, siliquis longissimis, teretibus, sessilibus. An *Hesperis Africana*. LINN. Sp. 4. *Hesperis Afric. hieracii folio hirsuto, flore minimo purpurascens*. NISSOL. Act. Par. An *Hesperis sylvestris, parva flore*. II. Prodr. 103. Folia

X

den-

dentata, incisuris raris, vno alterove denticulo notatis. Siliquae longissimae, teretes, fere sessiles, hispidae, stylo vix villo, apice acuto. A.

H. Hesperis caule hispido ramoso patente. *trifolius*. LINN. Sp. 1. Hesperis *pergrina*, siliquis articulat. I. R. H. Hesp. *Syriaca* CAMER. Hort. ic. 19. Hesperis *montana pallida* odoratissima. C. B. Hesperis. I. CLVS. 296.

Hesperis caule simplici erecto, foliis ovato-lanceolatis denticulatis, petalis mucrone emarginatis. *matronalis*. LINN. Sp. 2. Hesp. *sylv.* III. CLVS. 297. Hesp. *sylv. inodora*. I. R. H. HALL. Rupp. Fl. Jen. p. 78. n. a. T. 1. In ruderosis ad arcem Pless & Grubenhagen. *Salter Hist. n. a. 449. Non lat. p. 1. Plaque*
112/1000. non end.

β. Hesperis *hortensis*. C. B. *Viola matronalis*. DODON. Pempt. 161. & hortulanorum, flore albo, purpureo, utroque pleno.

H. Hesperis caule erecto ramoso, foliis cordatis amplexicaulibus serratis villosis. LINN. Sp. 7. *Leucoium marinum alterum latifolium purpureo-violeum*. LOB. ic. 333. *Leucoium marit. latifolium*. C. B. MORIS. Hist. II. p. 241. S. III. T. 8. f. 5.

(Hesperis caule recto, foliis scabris, lanceolatis, rariter dentatis, siliquis tetragonis. *Cheiranthus erysimoides*. LINN. Sp. 1. *Leucoium luteum sylvestre bieracifolium*. C. B. HALL. Rupp. Fl. Jen. p. 77. n. a. *Leucoium sylvestre*. CLVS. Hist. 299. HALL. G. 257. In M. Altenstollberg.)

H. Hesperis caule ramosissimo, folioso, foliis linearibus, rariter dentatis, glabris, siliquis teretibus, corollis vix calyce maioribus. *Inter folia longe linearis*.

H. *Astragalus caulescens*, erectiusculus, floribus
 spicatis, leguminibus erectiusculis nudis tumidis
 tereti-depressis; mucrone reflexo. *uliginosus*. LINN.
Sp. 8. *H. Vpf.* 226. n. 3. *. *Astrag. pedunculis folio*
supparibus, floribus pendulis, siliquis erectis &
longis parum compressis. Descripti hanc plantam in
Comm. Gott. T. V. P.

H. *Astragalus caulescens erectus laevis*, peduncu-
 lis spicatis, leguminibus ovato-cylindricis stylo acu-
 minatis. LINN. *Sp.* 9. *Astrag. procerior non repens,*
flore viridi flavescente. DILL. *Elth. T.* 39. f. 45.
In Virginia. P.

H. *Astragalus caulescens prostratus*, leguminibus
 subglobosis inflatis mucronatis pilosis. LINN. *Sp.* 11.
H. Vpf. 226. *Astrag. HALL. H.* 11. *Astrag. luteus,*
perennis, siliqua gemella, rotunda, vesicam refe-
rente. I. R. H. *Cicer sylvestre, foliis oblongis, bi-*
spidis, maius. C. B. *Circa Jenam & in Sibiria. P.*

Astragalus caulescens prostratus, leguminibus
 subtriquetris arcuatis, foliis ovalibus pedunculo lon-
 gioribus. LINN. *Sp.* 13. *Astragal. HALL. H.* 1.
Astrag. luteus perennis procumbens vulgaris. I. R. H.
Glaux vulgaris, Glycyrrhiza sylv. Gesneri. CLVS.
 CCXXXIII. *In pratis & sylvis. P.*

H. *Astragalus caulescens procumbens*, legumini-
 bus subulatis recurvatis glabris. LINN. *Sp.* 14. *H.*
Vpf. 5. *Astrag. luteus annuus Monspeliacus, pro-*
cumbens. I. R. H. *Astragalus Monspeliacus,* CLVS.
 CCXXXIV. *Monspeli.* A.

H. Astragalus caulescens procumbens, leguminibus contorduplicatis canaliculatis, villosis. ^{contorduplicatis} LINN. Sp. 15. H. Vpf. 227. n. 8. Astrag. *siliquis recurvis depressis hirsutis*, HALL. Comm. Gott. T. II. p. 339. Tab. XI. In Sibiria.

H. Astragalus caulescens procumbens, spicis pedunculatis, leguminibus prismaticis rectis triquetris apice uncinatis. ^{baeticus} LINN. Sp. 16. Astrag. *triangularis*. MÜNTING. ic, p. 110. Astrag. *Baeticus*. RIV. T. 105. A.

H. Astragalus caulescens diffusus, capitulis subsessilibus lateralibus, leguminibus erectis subulatis acumine reflexis. ^{colymneus} LINN. Sp. 17. *Vicia Sesamacea apula*. COLUMN. Ecphr. I. p. 303. T. 301. In Gallia Narbonensi. A.

H. Astragalus caulescens procumbens, leguminibus capitatis cordatis acutis reflexis complicatis. ^{cruciale} LINN. Sp. 18. Astragalus *Hisp. siliqua epiglottidi simili, flore purpureo, maior*. HERM. Lugdb. 74. T. 75. *Glaux Hispanica*. RIV. T. 109. A.

PHACA. LINN. 798.

H. Phaca caulescens erecta tota glabra sulcata, pedunculis folio longioribus, leguminibus erectis subtriquetris. In Comm. Gott. T. V. descripti, addita ieme. *Astragalus caulescens erectus, caule glabro sulcato*. ^{fulcrata} LINN. Sp. 6. *Foliola 8. f. 9. parium, lineari-lanceolata glabra. Pedunculi, floribus delapsis, foliis triplo longiores. Spica laxa pauciflora flores parvi, violacei. Legumina unilocularia, subtri-*

triquetra, inflata percurrit fascia, tanquam septi rudimentum. P.

H. Phaca pedunculis radicatis, hirsutissimis, foliolis ex ovato acutis saepe imbricatis. *Comm. Gott. T. V. c. 10.* descripsi. *Astragalus acaulis, scapo erecto foliis longiore, leguminibus subulatis inflatis, villosis erectis.* ^{Vrncea} LINN. *Sp.* 27. H. *Vps.* *Astragalus non ramosus villosus & incanus spicatus, floribus purpureo-violaceis.* AMMANN. *Ruth.* 167.

PSORALEA. LINN. 801.

H. Pforalea foliis pinnatis, foliolis linearibus acuminatis, floribus axillaribus. *pinnata.* LINN. *Sp.* 1. Genitae affinis *arbor Afric. monospermos, flore coeruleo, foliis pinnatis.* HERM. *Eugdb.* T. 273. In *Aethiopia.* Fr.

H. Pforalea foliis omnibus ternatis, pedunculis spicatis folio longioribus. ^{betulinifolia} LINN. *Sp.* 4. *Vulneraria foliis ternatis, floribus umbellatis.* HALL. *G.* 261. *Trifolium bituminosum.* DOD. *Pempt.* 566. RIV. T. 14. In *Italia, Gallia australi.* Fr.

H. Pforalea foliis simplicibus ovatis. *corylifolia.* LINN. *Sp.* 6. H. *Vps.* 225. *Pfor. folio integro subrotundo.* JUSSIEV. *Act. Par.* 1744. p. 382. *Loto affinis coryli folio.* PLVKN. *Alm.* 227. T. 96. f. 5. In *India.* A.

VVLNERARIA. HALL. 569. TOURN. 211.

Anthyllis LINN. 773.

H. *Vulneraria (Anthyllis)* herbacea, foliis quaternopinnatis, floribus lateralibus. LINN. *Sp.* 1. H.

Vps. 221. n. 1. *Vulneraria pentaphyllos.* HALL G. 261. *Anthyllis Hispanica.* RIV. T. 18. *Trifolium balicacabum.* CAMER. hort. 171. T. 47. In Italia. A.

Vulneraria foliis ad terram simplicibus, ovatis, ad caulem pinnatis. HALL H. 2. p. 570. *Anthyllis herbacea, foliis pinnatis inaequalibus, capitulo dupplicato.* LINN. Sp. 2. *Vulneraria rustica.* 1. R. H.

β. *Vulneraria supina, flore coccineo.* DILL. *Elth.* 431. T. 320. f. 413. In Helvetia. P.

h. *Vulneraria foliis pinnatis sericeis nitentibus.* HALL. G. 261. *Anthyllis fruticosa, foliis pinnatis aequalibus, floribus capitat.* LINN. Sp. 6. H. *Vps.* n. 3. *Barba Iovis.* C. B. RIV. T. 4.

DORYCNIVM. LINN. ed. pr. 602.

TOVRN. 211.

h. *Dorycnium foliis quinatis digitatis sessilibus.* LINN. *Cliff.* 371. HALL. H. p. 570. *Lotus* LINN. Sp. 15. *Dorycnium Hispanicum.* CLVS. 100. *Monspeliensum.* LOBEL. ic. 51. In Europa australi. A.

LOTVS. LINN. 803. HALL. H. 571.

h. *Lotus leguminibus solitariis membranaceo-quadrangulatis, bracteis ovatis, (caule erecto.)* LINN. Sp. 2. *Lotus pulcherrima tetragonolobos.* CAMER. Hort. 91. T. 26. In Sicilia. A.

h. *Lotus leguminibus solitariis membranaceo-quadrangulatis, bracteis ovato-lanceolatis, caule pro-*

procumbente. HALL. H. 1. Lotus filiquosus luteus minor & mollior. I. R. H. Lotus tetragonolobus. RIV. T. 39. In Thuringia. A. Lotus filiquosus spec. Ed. 2.

H. Lotus leguminibus subbinatis linearibus compressis nutantibus. LINN. Sp. 6. Lotus filiquis geminis, peregrinis. BOERH. Lugdb. II. p. 38. HALL. G. 262. In Europa australi, A.

H. Lotus leguminibus subquinatis arcuatis compressis, caulibus diffusis. LINN. Sp. 8. Lotus filiquis Ornithopodii. C. B. CAMER. Hort. 91. T. 25. In Sicilia.

H. Lotus leguminibus subternatis, caule herbaceo erecto, foliolis linearibus. LINN. Sp. 9. Lotus angustifolia, flore luteo purpurascente, insulae S. Jacobi. COMMED. H. II. p. 169. T. 83.

H. Lotus capitulis dimidiatis, caule fruticoso, foliis nitidis. LINN. Sp. 10. Lotus argentea Cretica. PLYKN. Alm. 226. T. 43. In Syria. Creta.

H. Lotus capitulis hirsutis, caule erecto hirsuto, leguminibus ovatis. LINN. Sp. 11. Lotus pentaphyllus, filiquosus, hirsutus. I. R. H. HALL. G. 262. In Gallia Narbonensi.

Lotus capitulis depressis, caulibus decumbentibus, leguminibus cylindricis. LINN. Sp. 13. Lotus floribus umbellatis, filiquis pendulis, caule erecto. HALL. Helv. 1. In dumetis bey der Papiermühle.

β. Lotus floribus umbellatis, filiquis pendulis, caule procumbente. HALL. H. 2.

H. Lotus capitulis subglobosis, caule erecto, leguminibus rectis glabris, ^{rectis} LINN. Sp. 12. Lotus villosus, flore altissimo, glomerato. I. R. H. HALL. G. 262.

H. Lotus oligoceros, Cretica, lutea, filiquis longis erectis, MORIS. HALL. G. 263.

ORNITHOPVS. LINN. 790. Ornithopodium. TOURN.

H. Ornithopus foliis pinnatis, leguminibus articulatis levibus subarcuatis. HALL. H. 572. ^{neruosis} LINN. Sp. 1. Ornithopodium maius & minus. C. B. I. R. H. In Helvetia. A. ad p. 4. avaras inter Annovera et Colly.

H. Ornithopus foliis pinnatis, leguminibus compressis rugosis. LINN. Sp. 2. Ornithopodium scorpioides, filiqua compressa. I. R. H. HALL. G. 264. In Italia. A.

H. Ornithopus foliis pinnatis, leguminibus semicircularibus levissimis. HALL. G. 264. c. descr.

H. Ornithopus foliis ternatis subsessilibus appendiculatis: impari maximo. ^{corpulentis} LINN. Sp. 3. Ornithopodium portulacae folio. I. R. H.

HIPPOCREPIS. LINN. 791.

H. Hippocrepis leguminibus sessilibus solitariis, (vtrinque excissis) ^{unifloris} LINN. Sp. 1. HALL. Gott. 265. Ferrum equinum filiqua singulari. C. B. GARD. DEL. T. 114. Ferrum equinum vulgare. COLUMN. Ecphr. I. T. 300. In Italia. A.

H. Hip-

Hippocrepis leguminibus pedunculatis confertis, margine exteriore repandis. LINN. *Sp.* 3.
 Ferrum equinum *siliquis umbellatis, utrinque emarginatis.* HALL. *Helv.* 572. Ferrum equinum *capitatum.* COLUMN. *Ecphr.* I. T. 301. In apricis M. *Plesi.* P.

SCORPIURVS. LINN. 792.

H. Scorpiurus pedunculis unifloris, leguminibus testis undique squamis obtusis. *vermiculata.* LINN. *Sp.* 1. Scorpioides *siliqua crassa boelii.* 1. R. H. 402. A.

H. Scorpiurus pedunculis bifloris, leguminibus extrorsum obtuse aculeatis. *muricata.* LINN. *Sp.* 2. Hanc refert MORIS. II. p. 127. S. II. T. 11. f. 4.

H. Scorpiurus pedunculis subtrifloris, leguminibus extrorsum spinis distinctis acutis. *fulcata.* LINN. *Sp.* 3.

H. Scorpiurus pedunculis subquadrifloris, leguminibus extrorsum spinis confertis acutis. LINN. *Sp.* 4. Scorpioides *corniculis magis asperis in se contortis & convolutis.* MORIS. II. p. 127. S. II. T. 11. f. 2.

AESCHYNOMENE. LINN. 769.

H. Aeschynomene caule hispido, foliolis acuminatis, leguminum articulis suborbiculatis. LINN. *Spec.* 2. ROY. *Lugdb.* 384. HALL. *G.* 265. c. *descr.* In *Jamaica.*

HEDYSARVM, LINN. 793.

H. *Hedysarum* foliis simplicibus lanceolatis obtusis, caule fruticoso spinoso. *Alhagi*, LINN. *Sp.* 1. In *Oriente*.

H. *Hedysarum* foliis simplicibus ternatisque, floribus racemosis. *Canadense*, LINN. *Sp.* 15. *Hedysarum triphyllum Canadense*, CORNV. *Canad.* T. 44. P.

H. *Hedysarum* foliis pinnatis, leguminibus articulatis aculeatis nudis rectis, caule diffuso, LINN. *Sp.* 28. *Hedys. clypeatum flore suaviter rubente*, I. R. H. 401.

β. *flore albo*, I. R. H. Fr.

H. *Hedysarum* foliis pinnatis, leguminibus articulatis aculeatis flexuosis, caule diffuso, LINN. *Sp.* 30. *Hedysarum annuum, siliqua aspera undulata, intorta*, I. R. H. *Hedys. siliquis undulatis*, RIV. T. 99. In *Italia*.

H. *Hedysarum* foliis pinnatis, leguminibus monospermis aculeatis, corollarum alis calyce brevioribus, LINN. *Sp.* 32. *Onobrychis spicis densioribus, siliquis asperis*, HALL. *Helv.* 577. *Onobrychis foliis viciae, fructu echinato maior*, C. B. *Esparsette* vulgo. Copiose feritur. Sponte in apricis Germaniae, Helvetiae. P.

H. *Hedysarum* foliis pinnatis, leguminibus monospermis aculeatis, petalis aequalibus, LINN. *Sp.* 33. *Onobrychis spicis rarioribus, siliquis aculeatis*, HALL. G. 267. *Onobrychis Cret. foliis Viciae*, fructu

fructu magno, cristato & aculeato. I. R. H. In Gal-
tia calidiori.

CORONILLA. LINN. 789.

H. *Coronilla fruticosa*, corollarum vnguibus ca-
lyce duplo longioribus. *Emerus*. LINN. Sp. 1.
Coronilla frutescens, siliquis longissimis obscure ar-
ticulatis. HALL. H. 4. *Emerus* I. R. H. *Coronilla*
scorpioides maior. C. B. In *Helvetia* & *Europa* au-
trali. Fr.

H. *Coronilla fruticosa*, stipulis subrotundis. LINN.
Sp. 3. *Polygala Valentina*. CLVS. 98. *Coronilla*
RIV. T. 92. In *Europa* australi.

H. *Coronilla foliolis denis*, leguminibus numero-
is articulatis pendulis. HALL. G. 268. *Helv. n. 4.*
Coronilla scorpioides altera. CLVS. 96. *Coronilla*
montana. RIV. T. 93. In *Helvetia*. *Austria*.

H. *Coronilla herbacea*, leguminibus falcato-gla-
tiatis. LINN. Sp. 5. *Securidaca*. HALL. H. 576.
Securidaca lutea maior. C. B. In *Italia*. *Hispania*. A.

Coronilla herbacea, leguminibus erectis tere-
bibus torosis numerosis, foliis glabris. *varia*. LINN.
Sp. 6. *Coronilla herbacea*, flore vario. I. R. H. *Se-*
curidacae II. *prior species*. CLVS. 237. Inter se-
getes. P.

H. *Coronilla herbacea* leguminibus quinis erectis
erectibus articulatis. LINN. Sp. 7. *Coronilla Creti-*
ca herbacea, flore parvo purpurascente. I. R. H.

COLVTEA. LINN. 776.

H. Colutea arborea foliolis obverse ovatis oblongis, apice emarginatis. Colutea ^{arborescens} *arborea foliolis obverse cordatis*. LINN. Sp. I. HALL. H. 575. Colutea *vesicaria*. C. B. I. R. H. In *Helvetia. Italia. Foliola duplo longiora, ac lata.*

H. Colutea fruticosa, foliolis subrotundis, apice retusis. Colutea *Orientalis, flore sanguinei coloris, lutea macula notato*. I. R. H. Cor. 44. Diversa priori videtur. *Foliola subrotunda, aequae lata ut longa. Vexillum duabus maculis flavis notatum.* F.

H. Colutea fruticosa, foliolis ovato-oblongis ^{fruticosa}. LINN. Sp. 2. Colutea *Aethiopica, barbae Jovis folio, flore phoeniceo*. I. R. H. Colutea *Aeth. fl. purpureo*. BREYN. Cent. T. 29. P.

H. Colutea herbacea, foliis linearibus. ^{herbacea} LINN. 9.
3. Colutea *Afr. annua, vesiculis compressis, foliis parvis mucronatis & cordatis*. COMM. H. II. T. 44. VOLKAM. Norib. p. 118.

ROBINIA. LINN. 775.

H. Robinia pedunculis racemosis, foliis imparipinnatis. ^{sub-acacia} LINN. Sp. 1. Pseudo-Acacia. RIV. T. 63. Acacia AMER. *Robini*. CORNV. *Canad.* 177. In *Virginia. Fr.*

H. *Robinia laricina*. H. *Robinia frutescens*. 70.

GALEGA. LINN. 770.

* *Galega*. ^{animalis} LINN. Spec. 715. BLACKW. T. 91. *vulgaris*. CLVS. CCXXXIII. I. R. H. *Ruta capraea* MATH. 739. *flore albo*. β . *flore subcoerulea*. In *Helvetia. Italia. P.*

INDIGOFERA. LINN. 794.

H. *Indigofera* leguminibus arcuatis incanis, racemosis folio brevioribus. *tinctoria*. LINN. Sp. 1. *Amoen.* Acad. I. p. 408. Anil 1. B. II. p. 945. In *India*. Fr.

GLYCIRRHIZA. LINN. 788.

H. *Glycyrrhiza* leguminibus glabris. ^{glabra} LINN. Sp. 2. Riv. T. 64. *filiquosa* & *Germanica*. B. Pin. 312. *vulgaris*. DODON. *Pempt.* 341. Colitur prope *Bambergam*. Fr.

H. *Glycyrrhiza* *estivata*

OROBVS. LINN. 780.

Orobus caule simplicissimo, foliolis pluribus ovatis acutis, stipulis semisagittatis. LINN. *Cliff.* 366. HALL. H. 1. *vernus*. LINN. Sp. 4. *Orobus sylvaticus purpureus vernus*. C. B. Verno tempore ubique in sylvis. A.

Orobus caule simplici, foliis imis ovatis, superioribus ellipticis acutis. HALL. H. 3. *tuberosus*. LINN. Sp. 5. *Orobus sylvaticus foliis oblongis glabris*. I. R. H. *Astragalus sylvaticus*. THAL. *Herc.* 7. T. 1. In sylva *Hercynica* circa *Eibingerode*.

Orobus caule ramoso, foliis sexiugis ovato-oblongis. ^{ovatus} LINN. Sp. 7. HALL. H. 4. *Orobus sylvaticus, viciae foliis*. C. B. *Orobus Pannonius*. II. CLVS. CCXXX. Circa *Rheinhausen*.

(H. *Orobus* fruticosus, pedunculis unifloris, foliolis plurimis subtus tomentosis. ^{Caragana} *Robinia*. LINN. Sp. 3. *Aspalathus arborescens, pinis foliorum crebrioribus oblongis*. AMMANN. *Ruth.* 285. *Caragana Sibirica*. ROYEN. 537. Fr.)

(H. Orobus pedunculis uni- & bifloris, foliis quaternatis cuneiformibus, glabris. Robinia. LINN. Sp. 4. Aspalathus frutescens maior latifolius, caerulea aureo. AMMANN. Ruth. 283. In Siberia. Vtrique species a Robiniae definitione generica maxime aliena, fabrica calycis, corollae, & legumine proxime ad Orobum accedit. Fr.)

H. Orobus foliis coniugatis subfessilibus, stipulis dentatis. Lathyroides. LINN. Sp. 1. Lathyroides erecta, folio ovato acuminato, coeruleis viciae floribus & siliquis. AMMANN. Ruth. 151. T. 7. f. 2. P.

PISVM. LINN. 779.

H. Pisum stipulis crenatis, pedunculis multifloris. LINN. Cliff. 369. sativum. Sp. 1. BLACKW. T. 89. Pisum bortenfe maius. G. B. 3. Pisum namm. RIV. T. 39. A.

H. Pisum petiolis decurrentibus membranaceis diphyllis, pedunculis unifloris. Ochrus. LINN. Sp. 4. Ochrus folio integro capreolum emittente. C. L. In Italia.

LATHYRVS. LINN. 781.

H. Lathyrus pedunculis unifloris, cirrhis aphyllis, stipulis sagittato cordatis, ^{achae} LINN. Sp. 1. Aphaca. HALL. H. 597. LOBEL. ic. II. p. 70. In Helvetia. Gallia. A.

H. Lathyrus pedunculis unifloris, foliis simplicibus, stipulis subulatis. Nissolia. LINN. Sp. 2. Lathyrus foliis singularibus. HALL. H. 1. Nissolia vulgaris. I. R. H. In Gallia. A.

H. Lr

α. *Lathyrus* pedunculis unifloris, cirrhis diphyllis, leguminibus ovatis compressis dorso bimarginatis. LINN. *Sp.* 5. *Lathyrus* HALL. *H.* 3. *Lathyrus sativus flore frustuque albo.* 1. R. H. Seritur in *Thuringia* & alibi. A.

β. flore coeruleo. *Cicerula Aegyptia.* CLVS. CCXXVI.

γ. flore ex albo & coeruleo vario.

δ. *Lathyrus* pedunculis unifloris, calyce brevioribus, cirrhis diphyllis, foliolis lanceolatis. LINN. *Sp.* 6. *Lathyrus orientalis, flore vix conspicuo.* BOERH. II. p. 42. HALL. *Gott.* 289. A.

ε. *Lathyrus* flore miniato, siliquis & seminibus maculatis. HALL. *G.* 289.

ζ. *Lathyrus* pedunculis subunifloris, cirrhis polyphyllis, foliolis alternis. *articulatus.* LINN. *Sp.* 10. *Clymenum Hispanicum, folio vario, siliqua articulata.* 1. R. H. In *Baetica.* A.

η. *Lathyrus* pedunculis bifloris, cirrhis diphyllis, foliolis ensiformibus, leguminibus glabris, stipulis bipartitis. LINN. *Amoen. Acad.* III. p. 417. *Lathyrus Hispanicus flore luteo.* RIV. *T.* 44.

θ. *Lathyrus* pedunculis bifloris, cirrhis diphyllis, foliis ovato-oblongis, leguminibus hirsutis: *odoratus.* LINN. *Sp.* 11. *Lathyrus distoplatyphyllos hirsutus mollis, flore odore magno.* COMMEL. II. *T.* 80. In *Sicilia.* A.

ι. flore ex albo & rubro vario.

κ. *Lathyrus* pedunculis bifloris, cirrhis diphyllis, foliolis lineari-lanceolatis, leguminibus hirsutis,

feminibus scabris. ^{hirsutus} LINN. Sp. 12. HALL. H. 5. Lathyrus *angustifolius*, *siliqua hirsuta*. C. B. In *Helvetica. Alsatia*. A.

H. Lathyrus pedunculis bifloris, cirrhis diphyllis, foliolis alternis lanceolatis. ^{hispanicus} LINN. Sp. 13. Lathyrus *Tingitanus*, *siliqua* Orobis, *flore amplo ruberrimo*. MORIS. II. p. 5. Lathyrus *pedunculis bifloris*, *cirrhis polyphyllis*, *stipulis dentatis*. LINN. Sp. 14. Clymenum *Hispanicum*, *flore vario*, *siliqua plana*. L. R. H

Lathyrus pedunculis multifloris, cirrhis diphyllis, foliolis valibus, internodiis nudis. ^{tuberosus} LINN. Sp. 15. HALL. H. 9. Lathyrus *arvensis repens tuberosus*. C. B. Terrae glandes DOD. *Hist.* 550. In acclivi M. *Heymberg*. A.

Lathyrus pedunculis multifloris, cirrhis diphyllis, foliolis lanceolatis, cirrhis simplicissimis. *pratensis*. LINN. Sp. 16. HALL. H. 10. Lathyrus *sylv. lutescus*, *folio Viciae*. C. B. In *pratibus*. P.

Lathyrus pedunculis multifloris, cirrhis diphyllis, foliis ensiformibus. ^{sylvestris} LINN. Sp. 17. Lathyrus *foliis binatis ellipticis*, *floribus racemosis*. HALL. H. 6. *Gott.* p. 290. Lathyrus *sylvestris*. C. B. CLVS. CXXIX. In *sylva ultra Geismar*.

Lathyrus pedunculis multifloris, cirrhis diphyllis, foliolis lanceolatis, internodiis membranaceis. ^{latifolius} LINN. Sp. 18. HALL. H. 8. Lathyrus *latifolius*. C. B.

β. Varietas *angustifolia*. Lathyrus *Narbonensis maior angustifolius*. C. B. Circa *Blankenburg*.

H. *Lathyrus pedunculis multifloris*, cirrhis polyphyllis, stipulis lanceolatis, *palustris*. LINN. *Sp.* 20. *Lathyrus foliis suis linearibus*. HALL *G.* 288. Clymenum *Parisiense flore coeruleo*. I. R. H. In *Thuringia*. P.

H. *Lathyrus paribus foliorum quinque*, floribus racemosis, filiquis incurvis. HALL. *Comm. Gott. T.* II. 1752. p. 341. An *Lathyrus pedunculis multifloris*, cirrhis polyphyllis, stipulis ovatis, basi acutis. LINN. *Sp.* 21. H. *Vpf.* 217.?

VICIA. LINN. 782.

(H. *Vicia foliis quaternis*, ovatis, floribus solitariis petiolatis. HALL. *Gott.* 291. *Lathyrus*. LINN. 9. *Clymenum Bithynicum*, filiqua singulari, flore minore. I. R. H. Habitus Lathyri, flos viciae.)

Vicia pedunculis multifloris, foliolis octonis ovatis amplissimis. LINN. *Sp.* 1. HALL. *H.* 7. *Pisum sylvestre*. CLVS. CXXIX. Cracca *flore ochroleuco*. RIV. *T.* 52. An der Lengelerburg versus Makenrode. A.

Vicia pendunculis multifloris, foliolis reflexis ovato mucronatis, stipulis subdentatis. LINN. *Sp.* 2. HALL. *H.* 5. *Vicia maxima dumetorum*. C. B. In dumetis ad prata prope Herperbausen, im Eulenloch. T. 1. 171.

Vicia foliolis ovalibus, stipulis denticulatis, filiquis racematim pendulis. HALL. *H.* 6. *T.* XIV. LINN. *Sp.* 3. *Viciae pulchrum genus*, multifolium. I. B. II. p. 316. lisdem locis cum priori. T. 1. 171.

H. *Vicia pedunculis multifloris, foliolis ovatis, stipulis maximis folia aequantibus. Folia octona, sena, ovato-elliptica, glabra. Stipulae maximae, crenulatae, acutae. Petioli duodecim-flori & ultra. Calyx coeruleus. Vexillum venosum. Legumina longa, angusta, glabra.* *Lathyrus hispanicus. Webb. Bot. 1. p. 69*

Vicia foliolis linearibus subtus sericeis, siliquis glabris, dense spicatis, pendulis, stipulis integris. HALL. H. 8. LINN. Sp. 5. *Vicia multiflora.* C. B. In pratis vdis.

H. *Vicia pedunculis subquinquefloris, foliolis integerrimis, stipulis integris, leguminibus erectiusculis.* LINN. Sp. 9. *Vicia Benghalensis hirsuta & incana, siliquis Pisi.* HERM. Lugdb. T. 625. A.

Vicia leguminibus sessilibus subbinatis erectis, foliis retusis, stipulis notatis. LINN. Sp. 10. HALL. H. 1. *Vicia vulgaris sativa.* C. B. 1. R. H. BLACKW. T. 423. Inter segetes; & copiose colitur.

Vicia leguminibus sessilibus erectis, foliis imis ovatis, superioribus linearibus. HALL. H. 3. *Vicia vulg. angustiore folio.* C. B. 1. R. H. *angustifolia.* RIV. T. 55. Inter segetes circa Lenglern & M. Hagen. *cf. var. Vicia sativa p. hagen*

H. *Vicia leguminibus sessilibus reflexis pilosis foliariis pentaspermis, corollae vexillis glabris.* LINN. Sp. 12. *Vicia sylvestris lutea, siliqua hirsuta.* 1. B. II. p. 313. HALL. G. 292.

H. *Vicia leguminibus sessilibus reflexis pilosis polyspermis, corollae vexillis pilosis.* An *Vicia*

LINN.

LINN. Sp. 13. *Vicia sylv. lutea cum galea fusca.*
I. B. II. p. 314. HALL. G. l. c.

β. *Vicia orient. carina & alis albis, vexillo reflexo subfusco.* DILL. HALL. G. l. c.

h. *Vicia leguminibus subsessilibus solitariis pendulis glabris tetraspermis, foliolis linearibus emarginatis.* LINN. Sp. 14. *Vicia angustifolia purpureo-violacea, siliqua lata glabra.* MAGNOL. Bot. Monsp. 276, HALL. G. 293. *Semina sphaerica.*

h. *Vicia leguminibus sessilibus solitariis glabris nodosis, seminibus lentiformibus.* HALL. G. 293.

Vicia leguminibus subquaternis erectis, foliis ex ovatis obtuse decrescentibus. HALL. H. 4. LINN. Sp. 15. *Vicia sepium.* RIV. T. 56. *folia rotundiore acuto.* C. B. I. R. H. Ad sepes & in sylvis.

h. *Vicia leguminibus subsessilibus solitariis erectis, foliolis fenis subovatis, stipulis denticulatis.* LINN. Sp. 16. *Vicia Narbonensis.* RIV. T. 57. HALL. G. 291. *Bona sylvestris.* DOD. Pempt. 516. *Folia primo bina, deinde quaterna, suprema sena.* In Europa australi.

Vicia foliis linearibus obtusis, siliquis gemellis polyspermis. HALL. H. 9. *Ervum* LINN. 2. *Vicia minima cum siliquis glabris.* I. R. H. In agris. A.

h. *Vicia floribus solitariis, seminibus compressis.* HALL. G. 296. *Ervum* LINN. 4. *Lens monanthos.* HERM. Lugdb. 360. A.

{ FABA. TOVRN. 212. }

h. *Faba.* C. B. I. R. H. *Faba maior.* BLACKW. T. 19. *Vicia caule erecto, petiolis absque cirrbis.* LINN.

Sp. 17. Varietates ex diversitate floris & feminis. Copiose feritur. *habitat non procul a mari Caspio in*
saffordii Paph. Linn. Syst. nat. 12. p. 128.

CICER. LINN. 783.

Cicer foliis ferratis. ^{arietinum} LINN. Sp. 1. Cicer *arietinum*. DOD. *Pempt.* 525. *sativum*. C. B. Inter segetes Hispaniae.

ERVVM. LINN. 784. Lens TOURN.

H. *Ervum* seminibus compressis convexis. LINN. Sp. 1. *Lens*. HALL. H. 601. RIV. T. 35. *Lens vulgaris*. C. B. I. R. H.

β *Lens maior*. RIV. T. 35. I. R. H. diversa non videtur. A.

Ervum pedunculis multifloris, seminibus globosis. *hirsutum*. LINN. Sp. 3. *Vicia filiquis hirsutis, pendulis, dispermis*. HALL. H. 10. *Vicia segetum cum filiquis plurimis hirsutis*. C. B. In agris. A.

H. *Ervum* foliolis linearibus retusis octonis, pedunculis bifloris. *Vicia multiflora cum filiquis latis*. HALL. G. 296. In *Pensylvania*. A.

H. *Ervum* leguminibus undato-plicatis. ^{crusile} LINN. Sp. 5. *Ervum leguminibus pendulis*. HALL. H. 601. *Ervum verum*. I. R. H. In hortis rarius feritur. A.

H. *Ervum* caule, & foliolis obverse ovatis, hirsutis, petiolis mucronatis, leguminibus solitariis polyspermis erectis sessilibus. Circa *Berolinum*. A *Vicia minima*. RIV. RVPP. I. R. H. diversa videtur, quod pedunculi neque foliis longiores, neque cirrhosti sint, quales huic tribuit ^{isplenicafe} LINNÆVS *Cent. Pl. II. n. 185*.

Ervum tetrahermanum
H. *Ervum monanthos* } p. 357.

ONO-

ONONIS. LINN. 772. Anonis. HALL.

Ononis floribus racemosis geminatis, foliis ternatis, ramis spinosis. ^{innata} LINN. Sp. 1. Anonis floribus pene sessilibus, vaginis foliorum amplissimis HALL. H. 4. Anonis, *vesta bovis*. TABERN. p. 528. Anonis spinosa, flore purpureo. C. B. I. R. H. BLACKW. T. 301. Ad vias. Fr.

β. flore albo. In colle prope Bovenden.

γ. spinis destituta iunior.

Η. Ononis spicis foliosis, foliis simplicibus obtusis. ^{alopocurroides} LINN. Sp. 5. Anonis *alopocurroides* Sicula. I. R. H. HALL. G. 284. In Europa australiori. A.

Η. Ononis pedunculis unifloris, filo terminatis, foliis ternatis. ^{patens} LINN. Sp. 7. Anonis pedunculis longis unifloris, leguminibus pendulis, teretibus hirsutis. HALL. H. 6. Anonis spinis carens, lutea maior. I. R. H. Natrix RIV. T. 68.

Η. Ononis pedunculis bifloris filo terminatis. ^{ornithopodioides} LINN. Sp. 9. Anonis pedunculis longis, leguminibus pendulis articulatis compressis. HALL. G. 284. Anonis siliquis Ornithopodii. I. R. H. In Sicilia.

Et ^{innata} *Anonis viscosa*

SPARTIVM. LINN. 765.

Η. Spartium inerme, ramis oppositis teretibus, rariter foliosis, apice floriferis. *iunceum*. LINN. Sp. 1. Spartium *Dioscoridis*. CLVS. 101. Genista *transalpina*. CAMER. H. ic. XIV. Genista *Hispanica*. RIV. T. 66. In Provincia. Hispania. Fr.

Μ. Spartium foliis ternatis solitariisque, ramis inermibus angulatis. Spartium *scoparium*. LINN. Sp.

7. Genista TABERN. 1509, RIVIN. T. 65. Genista *angulosa* & *sc paria*. C.B. In arenosis *Marchiae* & circa *Noribergam*. Fr. *Apex Mundum ad rivum*
Julio Henrichs de Hertenberg

H. Spartium foliis solitariis ternatisque, ramis sexangularibus apice floriferis. ^{angulatum} LINN. Sp. 5. Non dum floruit, sed ex angulis tribus, per ramos a singulo folio decurrentibus, facile cognoscitur. Fr.

GENISTA. LINN. 766, HALL. 592.

H. Genista ramulis striatis & foliis ternatis villosis. *Canariensis*. LINN. Sp. 1. Cytisus *minoribus foliis*, ramulis *tenellis villosis*. B. Pin. 390. Cytisus. L. CLVS. 94. Cytisus *Canariensis sempervirens* & *incanus*. COMMEL. H. II. T. 52. Fr.

H. Genista foliorum parte prima cauli adnata, elliptica, altera patente lanceolata. HALL. H. 4. Genista *ramis angustibus articulatis*, foliis *ovato-lanceolatis*. ^{lanceolata} LINN. Sp. 2. Chamaegenista *altera*. CLVS. 104. Genista *sagittalis Pannonica*. CAMER. H. ic. XIII. In australiori *Germanica*. Fr.

Genista foliis confertis lanceolatis, ramis striatis *erectis*, floribus sessilibus spicatis. HALL. H. 1. ^{ramulosa} LINN. Sp. 4. Genista *tinctoria Germanica*. C. B. 395. *vulgaris*. CLVS. 101. In sylvis & ad oras agrorum. Fr.

Genista foliis lanceolato-obtusis villosis, caule tuberculato decumbente. *pilosa*. LINN. Sp. 5. HALL. H. 2. Genista *ramosa foliis Hyperici*. C. B. I. R. H. Chamaegenista *prima*. CLVS. 103. In sylva *bercyrica* circa M. *Bruderum*. Fr.

H. Ge-

H. *Genista spinis simplicibus*, in ramis florigeris, foliis levibus lanceolatis solitariis. HALL. G. 186.
^{fragilis} LINN. *Sp.* 6. *Genista minor asphaltoides*. C. B. *Prodr.* 157. In arenosis *Cellensis*, inter colias et *Soleptra*.

Genista foliis ovato lanceolatis, hirsutis, spinis ramosis in ramis senescentibus. HALL. H. ^{Germanica} 3. LINN. *Sp.* 7. *Genista spinosa minor Germanica*. C. B. In sylva ultra *Rheinhausen*. Fr.

ARACHIS. LINN. 787.

Arachis ^{typica} LINN. *Sp.* 1. *Arachidnoides Americana*. NISSELE *At. Par.* 1727. HALL. G. 287.

CYTISVS. LINN. 786.

Cytisus racemis simplicibus pendulis, foliolis ovato oblongis. *Laburnum*. LINN. *Sp.* 1. HALL. H. 1. p. 591. *Laburnum*. RIV. T. 62. Circa arcem *Grubenbagen*. Fr.

CROTALARIA. LINN. 771.

H. *Crotalaria* foliis ternatis ovatis, stipulis nullis, leguminibus hirsutis. *incana*. LINN. *Sp.* 11. *Crotalaria foliis ternatis, petiolis nudis*. CLIFF. 358. c. *descr.*

ERYTHRINA. LINN. 762.

H. *Erythrina* foliis ternatis, caule arboreo aculeato. *Corallodendron*. LINN. *Sp.* 2. *Cliff.* 354. *Corall arbor Americana*. COMMEL. H. 1. T. 108. In *America*. Fr.

LUPINVS. LINN. 774.

H. Lupinus calycibus alternis inappendiculatis, labio superiore integro, inferiore tridentato. ^{albus} LINN. Sp. 2. Lupinus flore albo. CLVS. CCXXIX. RIV. T. 25. A.

H. Lupinus calycibus semiverticillatis, labio superiore bifido, inferiore subtridentato. varius. LINN. Sp. 3. Lupinus sylv. flore purpureo, semine rotundo vario. I. B. II. p. 291. Lupinus maior, flore purpureo. HALL. G. 270. A.

H. Lupinus calycibus verticillatis appendiculatis, labio superiore & inferiore integris. hirsutus. LINN. Sp. 4. Lupinus villosus coeruleus maior. B. Prodr. 148. Lup. exoticus hirsutissimus. I. B. Hist. II. p. 280. flore coeruleo. RIV. T. 27. A.

H. Lupinus calycibus alternis appendiculatis, labio superiore bipartito, inferiore integro. angustifolius. LINN. Sp. 6. Lupinus flore coeruleo minore. RIV. T. 28. Indicus CORNVT. p. 207. A.

H. Lupinus calycibus verticillatis appendiculatis, labio superiore bipartito, inferiore tridentato. luteus. LINN. Sp. 7. Lupinus sylv. flore lutea. C. B. Pin. 348. RIVIN. T. 26. A.

PHASEOLVS. LINN. 777.

H. Phaseolus caule volubili, floribus racemosis geminis, bracteis calyce maioribus, leguminibus pendulis. ^{vulgaris} LINN. Sp. 1. Phaseolus vulgaris. LOB. ic. 59. Smilax bortensis f. Phaseolus maior. B. Pin. 339. Mire ludit colore & mole seminis.

H. Phaseolus caule volubili, pedunculis subquadrisfloris, bracteis calyce minoribus, leguminibus sublunatis superne granulatis. Phaseolus HALL. G. 271. n. 2. *Vexillum viride. Alae & carina albae, exsiccatae flavescentes.* An Phaseolus caule volubili, leguminibus acinaciformibus sublunatis laevibus. LINN. Sp. 3. ? Videtur.

H. Phaseolus caule volubili, floribus racemosis geminis, bracteis calyce brevioribus, leguminibus pendulis (^{vulgare coccineus} scabris). LINN. Sp. 2. Phaseolus puniceo flore. CORNV. Canad. 184. *Flos priori triplo maior, semper coccineus. Legumina scabra, nodosa.*

H. Phaseolus caule volubili, petiolis hirsutis alatis, bracteis calyce maioribus, corolla concolore. *Caulis scaber. Pedunculus communis hirsutus, parciales laeves. Petala omnia flava, bractee cum calyce ex viridi flavescentes.* An Phaseolus caule volubili, florum vexillo revoluti, calycibus concolore. LINN. Sp. 4. ^{inam viridis} Cliff. 359. *?

H. Phaseolus caule volubili, vexillis carinaque spiraliter convolutis. *Caracalla.* LINN. Sp. 9. Phaseolus Amer. flore cochleato, odorato, vulgo Caracalla. HALL. Gott. 271. Phaseolus Indicus, flore cochleato, TRIUMPH. Obs. T. 94.

DOLICHOS. LINN. 778.

H. Dolichos filiquis latissimis, superne verrucosis. HALL. G. 272. Dolichos leguminibus ovato-acinaciformibus, seminibus ^{scabris} ovatis bilo arcuato versus alteram extremitatem. LINN. Sp. 1. Phaseolus Aegyptius

Aegyptius, nigro semine. I. R. H. Phaseolus niger Lablab. ALPIN. *Aeg.* T. 75. A.

H. Dolichos foliis glabris, leguminibus angustis, erectis rigentibus. HALL. G. 272. Phaseolus orthocarpus. RIVIN. T. 32. Vexillum extus subluteum, intus violaceum. Alae violaceae.

H. Dolichos foliis glabris, intermedio subhastato, floribus gemellis, leguminibus cylindraceis, falcatis, deorsum recurvis. Folia amplissima, lateralia extrorsum maxime gibba, angulo interiori plane destituta: intermedium utrinque gibbum, finem versus auminatum. Stipulae lanceolatae. Vexillum amplum extus ex luteo album intus obsolete purpureum, in origine duabus maculis flavis notatum. Alae inflatae purpureae. Carina alba. Semina violacea. A.

H. Dolichos caule volubili perenni, foliis tomentosis, mollissimis, intermedio rhomboideo. Nondum floruit; ex habitu huc retuli. Fr.

H. Dolichos caule perenni, pedunculis capitatis, leguminibus strictis linearibus. LINN. *Sp.* 9. Dolichos caule perenni lignoso. H. Cliff. 360. T. 20. HALL. G. 273. Fr.

H. Dolichos volubilis annuus. HALL. G. 273.

TRIGONELLA. LINN. 804. Foenum graecum. TOURN.

H. Trigonella leguminibus pedunculatis congestis pendulis ovalibus compressis, caule diffuso, foliis subrotundis. LINN. *Sp.* 2. Melilotus supina latifolia,

ifolia, filiqua lata membranacea compressa. HALL. *Zamm. Gott. T. I. p. 213. T. XIII. Ad. Petrop. 8. p. 109. T. 12. In Sibiria. A.*

H. *Trigonella* leguminibus sessilibus subsolitariis rectis ensiformibus. ^{*Foenum sativum*} LINN. *Sp.* 6. *Foenum graecum, filiquis ensiformibus.* HALL. *G.* 274. *Foenum graecum sativum.* C. B.

H. *Trigonella* leguminibus paucioribus congestis rectis, linearibus, obtuse flexis. ^{*polyceratia*} LINN. *Spec.* 3. *Foenum graecum filiquis binis angustis, obtuse flexis.* HALL. *G. l. c. A.*

H. *Trigonella* (*Foenum graecum*) leguminibus pedunculatis quinque congestis declinatis subfalcatis compressis. HALL. *H. l. c. spinosa.* LINN. *Sp.* 4. *Foenum graecum sylv. polyceratum creticum maius.* BREYN. *Cent. T. 33. f. 1. A.*

H. *Trigonella* (*Foenum graecum*) leguminibus pedunculatis numerosis sessilibus declinatis subfalcatis glabris. HALL. *l. c. non fusilica* LINN. *Sp.* 5. *Foenum graecum sylv. minus polyceratum Monspeliense.* BREYN. *Cent. T. 33 f. 2. A.*

H. *Trigonella* (*Foenum graecum*) leguminibus spicatis numerosis falcatis declinatis subhirsutis. HALL. *l. c. corniculata* LINN. *Sp.* 7. *Trifolium corniculatum.* II. DOD. *Pempt. 573. In Italia. A.*

MEDICAGO. LINN. 805.

H. *Medicago* leguminibus lunatis margine integerrimis, caule arboreo. ^{*arborescens*} LINN. *Sp.* 1. *Medicago risolia frutescens, incana.* I. R. H. HALL. *G.* 275. *Cy.*

Cytisus Marantae. LOB. II. p. 46. MARTYN. *Georg. Virgil*. p. 232. qui pro vero *Cytiso* veterum habet. In regno *Neapolitano*. Fr.

H. *Medicago* foliis pinnatis, pinna extrema maxima, leguminibus reniformibus margine dentatis. ^{*renata*} LINN. Sp. 4. *Medicago Vulnerariae facie Hispanica*. I. R. H. In *Hispania*. A.

H. *Medicago* leguminibus reniformibus margine dentatis, foliis ternatis. ^{*radiata*} LINN. Sp. 3. *Medica filiqua reniformi, ciliata, amplissima*. I. R. H. *Lunaria radiata Itatorum*. LOB. II. p. 38. In *Italia*. A.

Medicago pedunculis racemosis, leguminibus lunatis, caule prostrato. ^{*falcata*} LINN. Sp. 6. *Medica* HALL. H. 3. *Falcata* RIVIN. T. 84. *Medica flavo flore* CLVS. II. CCXXXIII. In pratis. A.

Medicago spicis ovalibus, leguminibus reniformibus monospermis, caule procumbente. ^{*lupulina*} LINN. Sp. 7. *Medica* HALL. H. 4. *Trifolium pratense, capitulo brevior*. C. B. Pin. 328. *Melilotus minima*. RIV. T. 8. In pratis. A.

H. *Medicago* caule erecto, pedunculis racemosis, leguminibus contortis. ^{*sativa*} LINN. Sp. 5. *Medica* HALL. H. 2. *Medica sativa vulgo Luzerne*. In *Germania australiori & Helvetia*. P.

H. *Medicago* pedunculis racemosis, leguminibus cochleatis spinosis, caule procumbente tomentoso. ^{*marina*} LINN. Sp. 8. *Medica marina*. CLVS. CCXXXIII.

H. *Medicago* leguminibus cochleatis, stipulis dentatis, foliis serratis, caule diffuso. ^{*subserotina*} LINN. Sp. 9. Ad hanc refert LINNAEVS tanquam varietates omnes

eliquas cochleatas, quae ad varias species reduci posse videntur.

1. Medicago leguminibus solitariis cochleatis compressis planis, stipulis ciliatis, caule diffuso. SAUVAG. Medica fructu orbiculato, compresso, plano. RIV. HALL. 275. *orbicularis*.

2. Medicago leguminibus solitariis cochleatis compressis umbilicatis. Medica fructu scutellato. HALL. H. 276.

3. Medicago fructu tornatili. RIV. HALL. l. c. *tornata*.

4. Medicago fructu turbinato. RIV. β. Medica fructibus rugosis, compressis, floribus racemosis; HALL. l. c. γ. Medica dicarpis, fructu turbinato, rugoso. MORIS II. p. 153. *turbinata*.

5. Medicago fructu coronulam vtrinque dentatam referente. SAUVAG. Medica coronata. HALL. l. c.

δ. Medicago fructu echinato. Medica fructu turbinato, breviori, brevissime echinato. HALL. l. c. 277. *echinata*.

β. Medica fructu polygono. HALL. 278.

γ. Medica turbinata echinata. RIV.

δ Medica fructu turbinato, echinato, rigidiori. HALL. l. c.

ε. Medica fructu longius echinato. RIV.

7. Medicago leguminibus spinosis. Medica spinosa, fructu conico truncato, spinis dispersis. HALL. 279.

β. Medica fructu ovato, angustiori, hirsuto, spinis se decussantibus. HALL. l. c.

γ. Medica

γ. *Medica cochleata spinosa maior*, capsulae spinis longioribus sursum & deorsum tendentibus. MORIS. I. p. 153. S. II. T. 15. f. 8. 9.

8. *Medicago fructu spinoso, globoso, foliis oblongis, argute dentatis.* HALL. G. 278. *Trifolium cochleatum spinosum Syriacum, foliis laciniatis.* BREYN. Cent. T. 34.

MELILOTUS. HALL. 587. Trifolii

Sp. LINN.

H. *Melilotus (Trifolium) spicis oblongis, leguminibus feminudis mucronatis, caule erecto.* LINN. *Sp. 1. Melilotus maior sativa violacea.* HALL. G. 281. *Lotus hortensis odorata.* B. Pin. 330. *Lotus urbana.* BLACKW. T. 284. In Bohemia. A.

H. *Melilotus floribus minimis dense spicatis, leguminibus monospermis rugosis, caule erecto.* *Melilotus lutea minor, floribus & filiculis minoribus, spicatis & dense dispositis.* MORIS. II. p. 167. A.

Melilotus (Trifolium) leguminibus racemosis, nudis, rugosis, dispermis, caule erecto. LINN. *Sp. 4. HALL. H. 1. Melilotus officinarum.* C. B. I. R. H. BLACKW. T. 80. flore flavo & albo. Vbiq̃ue inter segetes.

H. *Melilotus foliis hirsutis rariter dentatis, scape paucifloro.* HALL. *Comm. Gott. 1752. T. II. p. A.*

H. *Melilotus (Trifolium) leguminibus racemosis nudis dispermis rugosis obtusis, caule erecto, foliolis integris.* LINN. *Sp. 5. Melilotus Italica folliculis rotundis.* C. B. *Melilotus Italica.* CAMER. Hort. ic. 29. In Italia.

H. Me-

h. Melilotus (*Trifolium*) leguminibus racemosis nudis dispermis membranaceis, caule erectiuscullo. LINN. Sp. 6. Melilotus ^{cretica} Cretica, fructu orbiculari plano, maximo. 1. R. H. *Trifolium peltatum Creticum*. C. B. Prodr. T. 147.

TRIFOLIUM. LINN. 802.

Trifolium caule procumbente, & florigeris & foliosis petiolis longissimis. HALL. Helv. 5. *Trifolium capitulis umbellaribus, leguminibus tetraspermis, caule repente*. LINN. Sp. 12. *Trifol. prat. flore albo*. 1. R. H. In pratis.

h. *Trifolium capitulis villosis quinquefloris, involucro centrali reflexo rigido fructum involvente. subterraneum*. LINN. Sp. 15. *Trifolium album triococcum subterraneum gastonium reticulatum*. MORIS. II. p. 132. S. II. T. 14. f. 5. In Gallia. A.

h. *Trifolium spicis villosis longis, corollis monopetalis, caule erecto, foliis serrulatis*. LINN. Sp. 17. *Trifolium spicis villosis, calyce glabro, dentibus villosis, uno longissimo*. HALL. H. 11. *Trifol. mont. spica longissima rubente*. C. B. 338. In Thuringia.

Trifolium spicis subvillosis, cinctis stipulis oppositis membranaceis, corollis monopetalis. LINN. Sp. 18. *Trifol. spicis villosis, foliis insidentibus, vaginarum caudis capillaceis*. HALL. H. 14. *Trifol. prat. purpureum*. C. B. In pratis, & ob pabulum feritur. P.

Trifolium spicis villosis, foliis insidentibus, vaginarum caudis latioribus. HALL. H. 13. *Trifol.*

mont. purpureum, C. B. Hanc LINNAEVS cūm priori iungit. *alpestre* *Sp. 21. Lagopus 74. Bot. M. p. 14. 2.*

H. *Trifolium spicis villosis oblongis obtusis aphyllis*, calyce tomentoso, foliolis subrotundis. LINN. *Sp. 21. Trifol. spica f. brotunda rubra*, C. B. *Lagopus maximus*, CIVS. CCXLVI.

Trifolium spicis villosis conico-oblongis, dentibus calycinis setaceis subaequalibus, foliolis linearibus. LINN. *Sp. 23. HALL. H. 9. Trifolium arvense humile spicatum f. Lagopus*, C. B. *Pin. 328. In agris.*

H. *Trifolium spicis ovatis pilosis. denticulis calycinis lanceolatis patulis*. LINN. *Cliff. 375. *stellatum*. LINN. *Sp. 24. Trifolium stellatum*, C. B. *Pin. 329. Prodr. 143. In Europa calidiori. A.*

H. *Trifolium spicis ovatis, calycibus patulis: lacinia infima maxima lanceolata, foliis ovatis*. LINN. *Sp. 25. Trifol. clypeatum argenteum*. ALPIN. *Exot. T. 306. In Oriente. A.*

Trifolium caule erecto, spicis albis brevibusque terminato, foliis nervosis, inferne subhirsutis lanceolatis ferratis. HALL. *H. 8. montanum*. LINN. *Sp. 29. Trifolium mont. album*. C. B. I. R. H. *In pratis siccis. v. g. prope Hesselndreisch.*

H. *Trifolium spicis ovatis, calycibus inflatis glabris quinquedentatis, involucris vniuersalibus pentaphyllis*. LINN. *Sp. 32. HALL. G. 282.*

H. *Trifolium spicis subovatis, calycibus inflatis dorso gibbis, caulibus prostratis*. LINN. *Sp. 33. Trifol. pratense folliculatum*. I. R. H.

Trifolium capitulis subrotundis, calycibus inflatis bidentatis reflexis, caulibus repentibus. LINN. Sp. 35. Trifol. ^{fragiferum} ~~fragiferum~~ purpureum folio oblongo. I. R. H. fragiferum Friscum. C. B. CLVS. Cur. post. T. 39. In pratis gegen die Mäsbmühle.

Trifolium spicis ovalibus imbricatis, vexillis deflexis persistentibus, calycibus nudis, caule erecto. LINN. Spec. 36. HALL. H. 1. Trifolium ^{lupulinum} ~~lupulinum~~ alcorum minus R. & H. In pratis. *Nota spec. magn.*

Trifolium spicis ovalibus imbricatis, vexillis deflexis persistentibus, calycibus pilosis, caule erecto. LINN. Sp. 37. HALL. H. 3. Trifol. ^{montanum} ~~montanum~~ lupulinum. C. B. Prodr. 140. In sylvis.

Trifolium spicis ovalibus imbricatis, vexillis deflexis persistentibus, caulibus procumbentibus. LINN. Sp. 38. Trifol. ^{pratense} ~~pratense~~ luteum, capitulo lupuli s. agrarium. C. B. I. R. H. In siccis. *Nota spec. magn.*

H. Trifolium foliis quinis ciliatis sessilibus, leguminibus polyspermis. LINN. Sp. 9. Lupinaster ^{foliis} ~~foliis~~ purpureis & albis, siliquis minoribus. AMMANN. Rutb. 143. 144. In Sibiria.

PAPILIONACEIS ADFINES.

Cum Papilionaceis habitu, similitudine quadam floris, & fructu conveniunt, vario tamen modo colla, staminibus, & situ partium ab illis differunt.

AMORPHA. LINN. 768.

Vexillum solum adest, alae & carinae deficiunt.

H. *Amorpha* ^{fruticosa} LINN. Sp. p. 713. H. Cliff. 353.
T. 19. Barba Jovis Amer. *pseudo-acaciae foliis*,
flosculis purpureis minimis. H. Angl. In Carolina.

SOPHORA. LINN. 456.

Stamina decem distincta.

H. *Sophora* foliis pinnatis; foliolis numerosis
villosis oblongis. ^{alopon xidi} LINN. Sp. 1. *Ervum Orient. alo-*
peturoides perenne, fructu longissimo. DILL. Elth.
136. T. 112.

CERCIS. LINN. 458.

Corolla pentapetala florem papilionaceum refe-
rens. Stamina distincta.

H. *Cercis* foliis cordato-orbiculatis glabris. *Sili-*
quastrum LINN. Spec. 1. *Siliqua sylv. rotundifo-*
lia. G. B. Arbor Judae. DODON. Pempt. 786. In
Europa australiori; Palaestina; Aegypto.

CASSIA. LINN. 461.

H. *Cassia* foliolis triiugatis quadriiugatisque sub-
ovatis. ^{senaa} LINN. Sp. 8. *Senna Italica foliis obtusi-*
MORIS. II. p. 200. S. II. T. 24. f. 2. In *Aegypto*. Fr.

H. *Cassia* foliolis quinqueiugatis ovato-lanceola-
tis margine scabris, exterioribus maioribus, glandula
baseos petiolorum. ^{accidentia} LINN. Sp. 10. Fr.

H. *Cassia* foliolis quinqueiugatis ovato-lanceola-
tis glabris, glandula baseos petiolorum. ^{an-afra} LINN. Sp.
11. ROYEN. Lugdb. 468. HALL. Gott. 299.
deser. Fr.

H. Cassia foliolis multiugatis, calycibus acutis, floribus pentandris. ^{Acrotiana} LINN. Sp. 25. H. Cliff. 497. T. 36. HALL. G. l. c. c. descr. Senna spuria Virginiana, Mimosae foliis, floribus parvis nictitantibus. FLVKN. T. 314. f. 3. Fr.

MIMOSA. LINN. 597.

H. Mimosa foliis subdigitatis pinnatis, caule aculeato hispido. ^{rupea} LINN. Sp. 14. Mimosa spinosa III. siliquis parvis echinatis. BREYN. Cent. p. 40. T. 18. Fr.

H. Mimosa inermis, foliis bipinnatis, spicis cernuis, floribus decandris, inferioribus castratis apetalis, siliquis linearibus glabris. ^{sericata} LINN. Sp. 17. Mimosa spuria de Fernambouque, dicta Mimosa Italica. ZANON. T. 123. Fr.

H. Mimosa floribus superioribus pentapetalis decandris, inferioribus decapetalis sterilibus. HALL. Gott. 300. c. descr. Fr.

H. Mimosa spinis geminis distinctis, foliis duplicato-pinnatis, partialibus ytrinque quinopluribus. LINN. Cliff. 209. Mimosa. ^{cinerea} LINN. Sp. 29. Acacia ALP. Aegypt. T. 9. Fr.

H. Mimosa spinis geminis distinctis, foliis bipinnatis, partialibus octoiugatis. ^{farneisiana} LINN. Sp. 30. Acacia Indica Farneisiana. ALDIN. Farnef. 2. Fr.

PARKINSONIA. LINN. 460.

H. Parkinsonia ^{aculeata} LINN. Sp. p. 375. H. Cliff. T. 13. p. 157. Parkinsonia aculeata, foliis Mimosae uni costae affixis. WALTH. 6. T. 13. In America. Fr.

GLEDITSIA. LINN. 1025.

H. *Gleditsia triacanthos*. LINN. *Sp. p.* 1056
H. Vps. 298. GRONOV. *Virg.* 183. DV HAMEL
Tr. des arbres. T. I. p. 265. *sqq.* T. 105. Fr.

FVMARIA. LINN. 760.

Vexillum calcaratum. Filamentum duobus lateralibus, singula tribus antheris.

Fumaria caule simplici bifolio, radice cava, foliis inter flores integris. HALL. *H. p.* 604. *Fumaria bulbosa*, radice cava, maior. C. B. Ubique verno tempore in dumetis. ~~bulbosa a caua~~

Fumaria caule simplici multifolio, radice bulbosa solida, foliis inter flores palmatis. HALL. *H.* 605. *Fumaria bulbosa*, radice non cava, maior. C. B. Rarius in hortis. Circa Noribergam. *Fumaria caule simplici, bracteis longitudine florum.* LINN. *Sp. 3.* diversissimas hasce species iungit.

H. *Fumaria* siliquis linearibus paniculatis polyspermis, caule erecto. ^{*Aspermaria*} LINN. *Sp. 4.* Capnoides. I. R. H. *Fumaria siliquosa Canad. flore carneo & auct.* BARREL. *ic.* 103. A.

H. *Fumaria* siliquis linearibus tetragonis, caulibus diffusis acutangulis, radice perenni. ^{*Aspermaria*} LINN. *Sp. 5.* *Fumaria calyce donata, siliquis oblongis latiusculis.* HALL. *G.* 302. c. *descr. floris.* *Fumaria lutea.* C. B. *Pseudofumaria flore luteo.* RIVIN. *Tetrap. irreg. T.* 74. In Italia. P. | ^{*Aspermaria*} *Aspermaria* *fl. lutea. T. 7. 165.* *Aspermaria* *fl. lutea. T. 7. 165.*

Fumaria pericarpis monospermis racemosis, caule diffuso, foliolis subrotundis. ^{*Aspermaria*} LINN. *Sp. 7.* HALL. H.

H. 3. Fumaria offic. & Dioscoridis. C. B. BLACKW. T. 237. In agris & hortis.

H. Fumaria pericarpiis monospermis spicatis, caule erecto, foliolis capillaribus. LINN. *Sp. 8* HALL. *G. l. c. Fumaria minor, tenuifolia.* C. B. Capnos *tenuifolia.* CLVS. CCVIII. *Fumaria tenuifolia erecta Hisp. purpurea.* BARREL. *it. 41.* β . *flore albo, roseo.* In *Europa australi.*

H. Fumaria pericarpiis monospermis, caulibus se convolventibus. HALL. *G. 303* LINN. *Sp. 9. Fumaria viticulis & capreolis plantis vicinis adhaerens.* C. B. *Folia extrema subcirrhosa.*

H. Fumaria siliquis ovatis inflatis polyspermis, foliis cirrhiferis. LINN. *Sp. 11. Cysticapnos Afric. scandens.* BOERH. *Lugdb. I. T. 310.* In *Aethiopia.* A.

H. Fumaria floribus albis. HALL. *G. 301.*

POLYGALA. LINN. 761.

Carina sine anteriori appendiculata. Stamina octo connata.

Polygala floribus cristatis racemosis, caulibus herbaceis simplicibus procumbentibus, foliis linearilanceolatis. LINN. *Sp. 2. a. Polygala foliis omnibus acutis.* HALL. *H. 606. n. l. vulgaris maior.* C. B. CLVS. 324. In pratis & sylvis.

β . *Polygala foliis imis subrotundis, superioribus angustis acutis.* HALL. *Helv. 2. Polygala minor.* HALL. In locis aridis apricis. Varietas prioris. *Polygala minor L. 2 (per. 1. 787).*

H. *Polygala floribus imberbibus sparsis* : carina apice subrotundo, caule fruticoso, foliis lanceolatis. ^{*Chamaebuxus*} LINN. Sp. 11. *Polygaloides* HALL. H. 607. *Chamaebuxus flore Coluteae*. C. B. *Anonymos flore Coluteae*. CLVS. L. p. 105. Circa *Noribergam*.

AGGREGATAE. S. DIPSACEAE.

In plurimis flores in capitulum congesti, & calyce communi excepti, in nonnullis flores tantum paniculati, singuli squamis distincti, & in multis receptaculo communi innati. In omnibus, excepta *Globularia*, semini unica innascitur flos in plurimis monopetalus, in paucioribus pentapetalus. Stamina quatuor aut quinque corollae affixa, antheris distinctis. Tuba unica.

STATICE. LINN. 348. *Limonium*. T.

Statice caule nudo simplici capitato. *Armeria*. LINN. Sp. 1. *Statice*. RVPP. 96. *Limonium ophyllocaulon gramineum globosum*. MORIS. III. S. XV. T. 1. f. 29. *Caryophyllus mont. maior, flore globoso*. C. B. 211. *flore rubro & albo*. In sabulosis Germaniae. v. g. circa *Herzberg*. &c.

β. *Statice montana minor*. I. R. H. *Caulis humilis foliis aequalis. Flores maiores*.

H. *Statice* caule nudo paniculato tereti, foliis laevibus, obverse ovatis. ^{*Limonium*} LINN. Sp. 2. *Limonium maritimum maius*. C. B. LOB. 295. Ad littora maris.

H. *Statice* caule ancipiti dichotomo nudo, foliis lanceolatis mucronatis radicalibus, floribus alternis di-

distans. ^{purpurea} LINN. Sp. 6. HALL. G. 349. GMEL. Sib. II p 223. T. 92. *Limonium elatius*, *Plantaginifolius* foliis procumbentibus in aculeum terminatis, floribus albis spicatis. AMMANN. Ruth. n. 130. In Sibiria. P.

H. Statice foliis radicalibus alternatim pinnatifidis: caulinis ternis triquetris subulatis decurrentibus. ^{annua} LINN. Sp. 13. *Limonium peregrinum* *Asplenii folio*. C. B. *Limonium Rauwolfianum*. CLVS. Cur. post. p. 33. In Palaestina. A.

H. Statice caule fruticoso folioso, floribus solitariis, foliis lanceolatis vaginantibus. *monopetala*, LINN. Sp. 11. *Limonium lignosum*. BOCC. Sic. T. 17. Fr.

LAGOECIA. LINN. 251.

H. *Lagoecia*. LINN. Sp. 203. *Cuminoides*. I. R. H. *Cuminum sylvestre*. C. B. 186. CAMER. Epit. 519. In Oriente. A.

VALERIANA. LINN. 43.

Semine pappo plumoso. *Valerianae*. T.

H. *Valeriana* floribus monandris calcare donatis, foliis lanceolatis integerrimis. ^{ris Gra} LINN. Sp. 1. HALL. H. 6. *Valeriana rubra*. DOD. Pempt. 351. I. R. H. In Italiae, Helvetiae rupibus. A.

H. *Valeriana* floribus monandris, foliis imis integris, superioribus pinnatifidis. ^{valerianae} LINN. Sp. 2. *Valeriana folio Calcitrapae*. C. B. I. R. H. A.

^{divisa} *Valeriana* floribus triandris dioicis, foliis pinnatis. LINN. Sp. 4. *Valeriana foliis imis ovalibus*,

superioribus pinnatis, sexu distincta. HALL. H. 2.
α. Mar. Valeriana palustris minor. C. B. I. R. H.
 flos maior. *Femina β. Valeriana palustris inodora,*
parum laciniata. C. B. I. R. H. Flores minores. In
 paludosis.

Valeriana floribus triandris, foliis omnibus pin-
 natis. *officinalis.* LINN. Sp. 5. HALL. H. 1. Vale-
 riana *sylvestris.* I. CLVS. LX. *maior.* C. B. I. R. H.
 BLACKW. 271. In vdis sepibus & sylvis.

H. Valeriana floribus triandris, foliis radicalibus
 indivisis, caulinis pinnatis. *Ala.* LINN. Sp. 6. Valer-
burtenfis. C. B. Caulis non falcatus. In Gallia.

H. Valeriana floribus triandris, foliis omnibus
 integris ovato-oblongis, subdentatis, caule sim-
 plici. *montana.* LINN. Sp. 9. HALL. H. 4. Valer. *alpina,*
fol. Scrophulariae C. B. Prodr. 87. In Helvetiae
 montibus.

Semine non papposo. *Valerianellae.* TOWN-
 HALL.

Valeriana caule dichotomo, foliis oblongis rari-
 ter incis, feminis mucrone simplici. Valerianella
 HALL. H. 1. Valeriana. *Coccyza a. glauca.* LINN. Sp. 14. *α. Va-*
lerianella arvensis praecox humilis, semine compressa.
 I. R. H. *Lactuca agnina.* TABERN. Vbiqve verno
 tempore in gramineis. A.

Valeriana (*Valerianella*) caule dichotomo, fo-
 liis plerumque serratis, feminis corona tridentata.
 HALL. H. 2. *dentata.* LINN. Sp. cit. *δ. Valerianella arven-*
sis, serotina, altior, semine turgidiore. I. R. H.
 Caulis

Caulis magis ramosus. Squamulae flores excipientes glabrae. Serius floret.

H. Valeriana caule dichotomo, foliis pinnatifidis, feminis corona radiatim dentata. Valerianella *semine stellato*. J. R. H. LINN. *Sp. cit.* var. *γ. coronata*. *Squamulae floribus subiectae hirsutae, fructus hirsutus.*

H. Valeriana caule dichotomo, foliis lanceolatis serratis, feminis corona globosa inflata. Valerianella *Cretica fructu vesicato*. J. R. H. *Cer.* 6. BOERH. I. T. 75. *Semen subhirsutum in apice coronatur membranula in vesiculam globosam inflata, orificio contracto dentato, in cuius fundum apex seminis prominet.* *Valeriana. Lin. p. 23. 16.*

H. Valeriana floribus diandris ringentibus, foliis ovatis sessilibus. ^{*cornucopioidea*} LINN. *Sp.* 2. Valerianella *Indica rubra*. C. B. Valerianella *Cornucopioidea echinata*. COLUMN. *Ecphr.* 306. *Semen calyce quinquedentato coronatum. In calidiori Europa.*

H. Valeriana floribus tetrandris, foliis pinnatifidis, seminibus paleae ovali adnatis. ^{*Sibirica*} LINN. *Spec.* 16. Valerianella *tetrasemon*, *semine foliolo insidente*. HALL. *Comment. Gott.* 1751. p. 207. T. IX. In *Sibiria*.

BOERHAAVIA. LINN. 9.

H. *Boerhaavia* caule erecto, pedunculis folio glabro longioribus. An *Boerhaavia erecta*. LINN. *Sp.* 1. ? HALL. G. 352.

SCABIOSA. LINN. 108. HALL. Succisa,
HALL.

Corollae quadrifidae; involucrum feminis dentatum ciliatum.

Scabiosa (*Succisa*) foliis ad terram ovatis, ad caulem oblongis parum dentatis. HALL. H. 671. n. 1. Scabiosa corollulis quadrifidis aequalibus, caule simplici, ramis approximatis, foliis lanceolato-ovatis. ^{Succisa} LINN. Sp. 4. Succisa glabra. C. B. Succisa f. Morsus Diaboli. MATH. 623. BLACKW. T. 142. P. In pratis vdis, circa Harste, Bovenden & hercynicis.

H. Scabiosa foliis subhirsutis, pinnatifidis, lanceolatis, squamis calycinis hirsutis acutis, flore brevioribus. Calycis folia exteriora lineari-lanceolata, pilosa, squamae interiores apice fusco rigido aristatae. Involucrum feminis brevissime ciliatum. Segmenta floris albi omnia obtusa, inferius maius, apice viridi. A. An Scabiosa alpina? Linn. 2 spec.

H. Scabiosa (*Succisa*) capitulis nudis globosis, foliis glabris pinnatis solidis. HALL. Comm. Gott. T. I. p. 225. En. Gott. 354. Secundum descriptionem a priori diversa videtur.

H. Scabiosa foliis pinnatis, foliolis lanceolatis serratis, capitulis globosis hispidis congestis. HALL. H. 672. n. 2. ^{alpina} LINN. Sp. 1. Scabiosa alpina, foliis Centaurii maioris. C. B. In Helvetia. P.

H. Scabiosa corollulis radiantibus, caule hispido, foliis sessilibus serratis, squamis calycinis acutis. Omnino eadem esse videtur Scabiosa flosculis quadrifidis,

drifidis, foliis pinnatifidis, laciniis lateralibus erectiusculis. LINN. *Act. Vpsf.* 1744. p. 11. T. I. *Tatarica* LINN. *Sp.* 7. Nam etiam nostrae folia radicalia saepe sunt pinnatifida, lobis imbricatis. In *Tataria. Bienn.*

Scabiosa ^{armenica} corollulis radiantibus, caule hispido, foliis subhirsutis, imis ovatis, superioribus semipinnatis. HALL. *H.* 670. n. 1. LINN. *Sp.* 8. Scabiosa *pratensis hirsuta officinarum.* C. B. I. R. H. *sylvestris.* BLACKW. *T.* 185. In pratis & ad vias.

(ASTEROCEPHALVS. HALL. *H.* 668.)

Corollulae quinquefidae; feminis involucri annulo fimbriato plicatili coronatum; ex apice feminis setae quinque circa florem.

Asterocephalus (*Scabiosa*) foliis radicalibus ovatis crenatis, caulinis pinnatis setaceis, capitulis fructiferis globosis. HALL. *H.* 670. LINN. *Sp.* 10. Scabiosa *capitulo globoso minor.* C. B. In collibus siccis apricis. P.

n. Asterocephalus foliis pinnato-compositis, foliolis cuneiformibus incisis, superioribus linearibus, capitulis fructiferis subrotundis. Egregie hanc exprimit Scabiosa *glabra, carnosis foliis virentibus.* HERM. *Parad. Bat. ic.* p. 221. Nec alia esse videtur Scabiosa *mont. foliis scabiosae vulgaris.* C. B. I. R. H. Priori similis, diversa tamen *caule ramosissimo altitudinis humanae, foliis omnibus pinnatis, & setis circa florem fere semper paucioribus ac quinque.* P.

n. Astero-

H. *Astrocephalus* foliis bipinnatis linearibus, corollulis radiantibus, ochroleucis. Scabiola ^{egriobura} LINN. Sp. 17. Scabiola folio multifido, flore flavescens. VII. CLVS. III. MORIS. III. S. VI. T. 13. f. 23. III. HALLERVS dubitat, num a vulgari satis sit diversus.

H. *Astrocephalus* foliis lyratis, lobis profunde ferratis, capitulis globosis, annulo seminis amplissimo, parulo. Scabiola ^{stellata} LINN. I I. Sc. maior Hispanica. CLVS. I. stellata, folio lucinato, maior. C. B. I. R. H. HALL. G. 353. A.

H. *Astrocephalus* foliis inferioribus lyratis, serratis, superioribus linearibus integerrimis, capitulis cylindricis hispidis. Scabiola atropurpurea LINN. Sp. 12. Scab. peregrina rubra, capite oblongo. C. B. Scabiola VI. Indica. CLVS. III. Caulis herbaceus, annuus, diffusus. A.

H. *Astrocephalus* foliis inferioribus petiolatis ovatis serratis, superioribus pinnatis linearibus, capitulis cylindricis hispidis. Radix perennis, caulis firmior erectus ramosus. Folia glabra, nitida. Pedunculi hirti. An Scabiola Afr. frutescens maxima, foliis tenuissime incis. BOERH. Lugdb. I p. 128. In Africa. P.

H. *Astrocephalus* caule fruticoso, foliis petiolatis obovatis, crenatis, hirsutis, pedunculis hirsutis ramos terminantibus. Scabiola ^{Africana} LINN. Sp. 14. Scabiola Afr. frutescens, foliis rugosis & crenatis. HERM. Par. ic. p. 219. 220. Fr.

H. *Astrocephalus* foliis planis, carnosiss, inferioribus pinnatis, ramorum integerrimis, calyce florem

DIPSACEAE.

383

florem superante. GMEL. Sib. II. p. 213. T. 87.
HALL. G. 353. A.

H. *Asterocephalus* foliis carnosissimis hirsutis lanceolatis, calyce folioso florem excedente, corollulis aequalibus. *Planta humilis, caulis dichotome ramosus, folia sessilia, crassiuscula; folia calycis vncialia, hirsutissima, inaequalia; flos exiguus, corollulae omnes aequales, nec radiantes, limbo quinquesido regulari, carnaeae. Setae ex apice seminis rigidae, corollulis longiores.* Fufius illam descripsi in *Comment. Gott. T. V. A.*

DIPSACVS. LINN. 107.

Dipsacus foliis connatis ferratis, capitulis ovalibus, aristis calycinis rectis. HALL. *H. T.* LINN. *Dipsacus* *fullonum* a. *Sp. 1. a. Dipsacus sylv. aut virga pastoris maior.* c. H. BLACKW. T. 50. Ad oras agrorum, v. g. circa *Herperhausen.* A.

H. *Dipsacus* foliis sessilibus ferratis, capitulis ovalibus, aristis calycinis recurvis. HALL. G. 355. *Dipsacus sativus, fullonum.* c. B. In usum fullonum seritur.

H. *Dipsacus* foliis connatis sinuatis. LINN. *Sp. 3.* *Dips. folio laciniato.* c. B.

Dipsacus pilosus is a plant in Syria, that I believe is the same as the *Dipsacus pilosus* of the Arabs. LINN. *Sp. 3.*

KNAVTIA. LINN. 109.

H. *Knavtia* *orientalis.* LINN. *Sp. p. 101.* *Lychni-Scabiosa, flore rubro, annua.* BOERH. *Lugdb. I. p. 131.* In Oriente.

SYNGE-

SYNGENESIA. LINN. 2. Flores compositi, antheris quinque coalitis.

Plantae *flore composito* dicuntur, in quibus plures flosculi calyce communi comprehensi sunt. In omnibus hisce receptaculo communi innascuntur semina, quae singula ex apice proferunt florem monopetalum tubulosum, in aliis in limbum breviter quinquesidum, in aliis in ligulam planam extrorsum versam, terminatum. Ex pariete interno tubi floralis nascuntur quinque filamenta, antheris totidem linearibus in cylindrum connatis, per quem adscendit stylus, stigmate bifido ex tubo eminente.

Flosculis omnibus tubulosis, limbo quinquesido.

I. CAPITATAE. Tubus floris longissimus, limbi segmenta longiora; capitulum oblongum ventricosum.

ECHINOPS. LINN. 829.

Singulis flosculis in globum collectis calyx peculiaris.

H. *Echinops* calyculis unisloris, caule multifloro, foliis spinosis, supra nudis. *Sphaerocephalus*. LINN. *Sp.* 1. *maior*. 1. R. H. *Carduus sphaerocephalus latifolius vulgaris*. C. B. In Italia. P.

H. *Echinops* caule subunisloro, foliis duplicatopinnatifidis, foliolis latiusculis vicinis. Gmel. Sib. II. p. 100. ¹⁷⁵³ LINN. *Sp.* 2. H. *Vpf.* 248. *. In Sibiria.

H. *Echinops* calycibus fasciculatis: lateralibus sterilibus, foliis pinnatifidis supra strigosis. LINN. ¹⁷⁵³

- Sp. 3. *Echinopus minor, annuus, capite magno.*
 1. R. H. *Scabiosa Carduifolia annua.* HERM. *Par.*
T. 224. *Sphaerocephalus annuus.* CAMER. *Hort.*
c. 45. In *Hispania.*

Placenta alveolata.

ONOPORDON. LINN. 834.

Onopordon calycibus squarrosis, foliis decurren-
ibus ovato-oblongis sinuatis, caule brachiato. ^{*Acanthium*} LINN.

- Sp. 1. HALL. *H.* p. 674. *Spina alba tomentosa la-*
tifolia sylv. C. B. *Carduus alatus tomentosus, acan-*
thi folio, vulgaris. 1. R. H. Ad vias.

β. *Carduus luteus tomentosus, Acanthi folio*
flore magno. HALL. *G.* 357. A.

H. *Onopordon calycibus imbricatis, caule parum*
ramoso. ^{*Acanthium*} LINN. *Sp.* 3. *H.* *Vps.* 249. *Carduus to-*
mentosus, Acanthi folio, altissimus Lusitanicus.
 1. R. H. HALL *G.* 357.

H. *Onopordon calycibus squarrosis, foliis line-*
aribus pinnatifidis, vix caule latioribus. ^{*Acanthium*} LINN. *Sp.*
 2. *Acanthium Illyricum.* LOBEL. II. ic. I. In *Eu-*
ropa australi. A.

H. *Onopordon acaulon, capitulis squarrosis ner-*
vo foliorum insidentibus. HALL. *G.* 357. c. *deser.* B.

Placenta paleis aut pilis tecta.

CYNARA. LINN. 835.

H. *Cynara foliis subspinosis pinnatis indivisisque,*
calycinis squamis ovatis. ^{*Acanthium*} LINN. *Sp.* 1. *Cynara*
bort.

hort. foliis non aculeatis. C. B. Scolymus *Dioscori*
dir. CLVS. CLIII.

H. Cynara foliis spinosis: omnibus pinnatifidis,
calycinis squamis ovatis. ^{Cardunculus} LINN. Sp. 2. Cynara *spinosa*
curius cauliculi cillantur. C. B. Scolymus *aculeatus*.
TABERN. 1075.

LAPPA, HALL. 674. TOVRN. Arctium.
LINN 830.

Lappa capitulis tomentosis. HALL. *Helv.* 1. Arctium. LINN Sp. 1. β. Lappa *minor montana*, *capitulis tomentosis.* C. B. Vbiq̃ue ad vias.

Lappa capitulis glabris. HALL. H. 2. Arctium.
LINN Sp. 1. α. Lappa *maior.* f. *Arctium* *Dioscoridis.* C. B. 1. R. H. Bardana *officin.* Hisdem loci
cum priori.

Lappa capitulis glabris minimis. HALL. H. 3.

SERRATVLA. LINN. 831.

Serratula foliis pinnatifidis, petiolis multifloris,
calycibus glabris spinulosis. Serratula tinctoria.
LINN. Sp. 1. α. C. B. Carduus HALL. H. 5. Ser-
ratula *Mathioli.* CLVS. VIII. In M. arcis *Plesi.* P.

H Serratula foliis pinnatifidis, petiolis unifloris,
calycibus squalidis. Serratula ^{prostrata} ~~prostrata~~. LINN. Sp.
1. β. Carduus *inermis*, foliis glabri, pinnatis, la-
cinia extrema maxima, capitulis squalidis. Gmel
Sib. II. T. 20. In Sibiria. Planta priori omnibus
partibus multo maior. Squamæ calycinæ densè
conglutinatae & pulvere sordido fere tectæ, ut
ora alba, uti prior species, ornatae. P.

H. Serratula (*Carduus*) caule ramoso, foliis pin-
natifidis, foliolis inferiorum dentatis, squamis ca-
lycinis ex lanceolato-spinosis, appressis. G. MEL. Sib:
II. n. 38. T. 17.

Serratula foliis dentatis spinosis. ^{EDMUND.} LINN. Sp. 14.
Carduus foliis semipinnatis, spinosis, capitulis ob-
longis mitioribus. HALL. H. 5. *Carduus arvensis.*
TABERN. 700. In segetibus humidioribus.

H. Serratula caule ramosissimo, ramis patulis,
foliis petiolatis, ovato-acutis, nervosis, lunate
dentatis, flosculis calycem aequantibus. Sic interim
definire liceat plantulam, quae licet *figura foliorum*
& *capituli* ad *Cirsium alp. boni Herinci folio.*
HALL. H. T. 22. n. 16. accedere videatur, caule
tamen *ramosissimo*, aliisque notis longe ab illo se-
cedit. *Folia inferne incana, nec lanugine albida.*
Calycis parvi cylindrici squamae lanceolatae, cilia-
tae. Flosculi carnei coloris calycem non superant.
Pappus simplex, tubi floris longitudine. Placenta
villosa.

CARDVVS. LINN. 832. *Cirsium.*

HALL.

H. ^{Cirsium} (*Carduus*) foliis lanceolatis dentatis ciliatis de-
currentibus: spinis marginalibus duplicibus. LINN.
Sp. 1. *Acarna maior, caule folioso.* C. B. *Chamae-*
leon Salmaricensis. CLVS. CLV. *Squamae calycinæ*
in aculeum validissimum flavescentem abeunt.

Carduus (Cirsium) foliis decurrentibus spinosis,
pinnatis, laciniis divaricatis, capitulis ovatis tomen-
tosis spinosis, caule piloso. HALL. H. 7. LINN. Sp.

lis subquinis. ^{var. 14.} LINN. Sp. 14. *Cirsium spinosum, vennis foliorum lacteis*. HALL. G. 361. *Carduus lacteus Syriacus*. CAMER. Hort. 35. T. 10. In Syria. Hispania.

H. *Carduus foliis amplexicaulibus hastato-pinnatifidis spinosis, squamis calycinis margine apiceque spinosis*. LINN. Sp. 13. *Silybum nervis foliorum lacteis*. HALL. H. 684. *Carduus albis maculis notatus vulgaris*. C. B. I. R. H. *Carduus Mariae*. BLACKW. T. 79.

H. *Carduus foliis ex cordato-lanceolatis margine ferratis & spinosis, squamis calycum membranaceis laceris spinosis, capitulis nutantibus*. GMEL. Sib. II. p. 47. T. 19. *Cnicus foliis cordatis, petiolis crispis spinosis amplexicaulibus, floribus cernuis*. LINN. Sp. 5. H. Vpf. 251. *. In Sibiria.

H. *Carduus (Cirsium) foliis spinosis hirsutis, pinnis semibilobis, capitulis tomentosis sphaericis pungentibus*. HALL. H. 6. *Carduus eriophorus*. LINN. Sp. 15. Card. *Eriocephalus*. CLVS. CLIII. *Carduus capite rotundo tomentoso*. I. R. H. In montosis Sueviae & Helvetiae, P.

Carduus (Cirsium) inerme, radicibus teretibus congestis, pinnis foliorum semibilobis, ramis unifloris. HALL. G. 360. *Carduus tuberosus*. LINN. Sp. 118. *Carduus prat. Asphodeli radice, foliis tenuiter incisis*. C. B. flore purpureo. In sylvis. v. g. *Göttingensi, & Eulenloch. β. flore albo bey Harste & Herperhausen*.

H. *Carduus foliis lanceolatis amplexicaulibus, ferraturis spinoso-setaceis, capitulis triphyllis*.

Fatigue

LINN. Sp. 22. Omnia conveniunt, sed color floris in nostra planta *purpureus* est, quem *album* dicit.

Carduus *acaulis*, calyce glabro. *arab.* LINN. Sp. p. 1199. in app. Cirsum *caule floris longitudine*. HALL. 11. Cirsum *acaulos flore purpureo*. I. R. H. In siccis acclivibus. In horto caulis elongatur. P.

CNICVS. LINN. 833.

Cnicus caule erecto, foliis inferioribus laciniatis, superioribus integris concavis. *oleum* LINN. Sp. 2. Cirsum *inverme*, foliis *inver* *flavescentia folia congestis*. HALL. H. 4. Carduus *prat. latifolius*. C. B. LOB. II. p. 11. In pratis vdis.

Cnicus Arvensis in Sp. 2 p. 1156. Ann. Card. Arvensis
(H. Cnicus caule diffuso, foliis dentato-finuatis. LINN. Sp. 1. Cnicus *sylv. hirsutior s. Carduus benedictus*. C. B. offic. *Centauria benedicta* Linn. p. 92. 2. 2.)

CARTHAMVS. LINN. 838.

H. Carthamus foliis ovatis integris serrato-aculeatis. *tinctorius*. LINN. Sp. 1. Cnicus *sativus s. Carthamus officinarum*. C. B. Cnicus *vulgaris*. CLVS. CLII. In Aegypto. A.

H. Carthamus caule piloso, superne lanato, foliis interioribus pinnatifidis, summis amplexicaulibus dentatis. *lanatus* LINN. Sp. 2. Atractylis *succo sanguinea, flore luteo*. HALL. H. 685. Atractylis *lutea*. C. B. I. R. H. COLVMN. *Ephr.* p. 19. T. 23.

at Carthamus lanatus p. 390.

CARLINA.

CARLINA. LINN. 836.

Carlina caule uniflora. HALL. H. 686. n. 2.
^{acaules} LINN. Sp. 1. *Carlina acaulos, magno flore.* C. B.
 1. R. H. In collibus versus Allendorf.

β. *Carlina caulescens, magno flore albo.* C. B.
 1. R. H. Varietas *bortensis* eiusdem plantae, quae in
 loco culto caulem cubitalem profert.

Carlina caule multiflora, floribus terminalibus.
^{vulgaris} LINN. Sp. 4. *Carlina sylvestris vulgaris.* C. B. In
 apricis aridis.

Floribus in radio sterilibus.

CENTAVREA. LINN. 880.

1. *Rhapontica.* Calycibus laevibus inermibus.

H. *Centaurea calycibus inermibus*: squamis li-
 nearibus subulatis, foliis pinnatis serratis. LINN. Sp.
 1. *Rhaponticum foliis pinnatis ciliatis, calycibus*
acutis. HALL. G. 364. *Serratula annua semine cilia-*
ri elegantissimo. 1. R. H. *Senecio-Carduus apulus.*
 COLVMN. Ecphr. p. 34. ic. In Italia australi.

H. *Centaurea calycibus inermibus subrotundis*
glabris, squamis ovatis, foliis sinuatis. LINN. Sp.
 2. *Rhaponticum flosculis sterilibus quindecimfidis,*
squamis calycinis rotundis. HALL. l. c. *Cyanus ori-*
entalis maior moschatus. MORIS. III. S. VII. T. 25.
 f. 5. ZANON. T. 62. *Amberboi flore purpureo*
odorato. VAILL.

β. flore albo.

γ. flore luteo.

H. Centaurea squamis calycinis ovatis glabris, foliis pinnatis, foliolis decurrentibus, serratis & integris. GMEL. Sib. II. p. 89. T. 41. f. 2. ^{Rhaponticum} LINN. Sp. 6. Rhaponticum foliis pinnatis, pinnis serratis, squamis calycinis rotundis. HALL. Comm. Gott. 1751. p. 206. Centaurium maius L. vulg. CLVS. X. In Sibiria, & alpebus Italiae.

Centaurea calycibus squamosis, foliis lanceolatis radicalibus sinuato-dentatis, ramis angulatis. ^{Jacea} LINN. Sp. 26. Rhaponticum foliis squalidis tomentosis, inferioribus ad b sin sempinnatis, superioribus integris. HALL. H. 3. Jacea nigra prat. latifolia. C. B. In pratis.

H. Centaurea calycibus squamosis, foliis indivisis integerrimis decurrentibus. ^{plantula} LINN. Sp. 30. Rhaponticum foliis linguiformibus decurrentibus, capitulis argenteis. HALL. G. 364. Centaurium maius Orient. rectum glasti folio, flore luteo. COM-MEL. rar. T. 39. P.

Cyan. Squamis calycinis serrato-ciliatis.

Centaurea calycibus serratis, foliis linearibus integerrimis: infimis dentatis. ^{Cyanus} LINN. Sp. 12. Cyanus flosculis radialibus, latissimis, semiseptifidis. HALL. H. 4. Cyanus segetum. C. B. Inter segetes, β. flore purpureo. A.

H. Centaurea calycibus serratis, foliis lanceolatis decurrentibus, caule simplicissimo. ^{montana} LINN. Sp. 11. Cyanus caule alato, foliis integris subtus tomentosis. HALL. H. 5. Cyanus maior. BLACKW. T. 266. mont. latifolius. C. B. I. R. H. In montosis Helvetiae.

Centaurea (*Cyanus*) foliis ellipticis dentatis rigidis, squamis calycinis ferratis. HALL. G. 370. *Cnicus perennis coeruleus Tingitanus*. HERM. Lugdb. T. 163. *Carthamus foliis lanceolatis acuminato-ferratis*. LINN. Sp. 3.)

H. *Centaurea calycibus ciliatis, foliis pinnatifidis linearibus, caule paniculato*. LINN. Sp. 13. *Cyanus pinnis foliorum pinnatis, pinnis acutis gracilibus dentatis*. HALL. H. 1. *Jacea foliis candicantibus, laciniatis, calyculis non splendentibus*. I. R. H. *Stoebe Austria a elatior*. CLVS X. In *Thuringia*.

H. *Centaurea calycibus ciliatis, foliis tomentosis pinnatifidis, foliolis obtusis ovatis integerrimis, exterioribus maioribus*. LINN. Sp. 16. *Cyanus foliis pinnatis tomentosis*. HALL. G. 369. *Jacea arborea, argentea, Ragusina*. ALP. exot. 116. ZANNON. Hist. T. 92.

H. *Centaurea calycibus ciliatis subrotundis, foliis tomentosis indivisis pinnatifidisque integerrimis, caule declinato*. LINN. Sp. 19. GMEL. Sib. II. p. 96 T. 42. f. 2. *Cyanus foliis pinnatis, pinnis rotundis, superioribus ovatis, caule prostrato*. HALL. G. 368. *Comm. Gott. T. I. p. 204*. In *Sibiria*.

Centaurea calycibus margine ciliatis, foliis pinnatifidis, pinnis lanceolatis dentatis. LINN. Sp. 21. *Cyanus*. HALL. H. 2. *Jacea vulgaris laciniata*. I. R. H.

♂. flore albo.

♀. flore ochroleuco.

Bb 5

♂. flore

δ. flore flavo. Ex Sibiria. HALL. Comm. Gott. T. I. Tab. VII.

ε. flore helvolo.

η. foliis latioribus glaberrimis firmulis. Ex Sibiria.

H. Centaurea (*Cyanus*) calycibus ciliatis, foliis pinnatis tomentosis, pinnis subrotundis, flore helvolo. HALL. Comm. Gott. T. I. p. 203. En. Gott. 367. In Sibiria.

Centaurea calycibus recurvato-plumosis, foliis indivisis. ^{Phrygia} LINN. Sp. 7. Cyanus foliis ovato-lanceolatis subasperis, glumis calycis in reflexam plumulam desinentibus. HALL. H. 6. Jacea IV. Austr. villoso capite. CLVS. VII. Jacea cum squamis pennatis f. villoso capite. L. R. H. In sylvis bercynicis.

H. Centaurea calycibus recurvato-plumosis, foliis imis pinnatis dentatis: summis lanceolatis. ^{capillatis} LINN. Sp. 8. GMEL. Sib. II. p. 98. T. 43. Cyanus foliis calycinis plumosis reflexis, foliis pinnatis. HALL. Comm. Gott. T. I. p. 202. Tab. 6.

H. Centaurea (*Cyanus*) calycis plumulis reflexis, foliis linguiformibus subhirsutis. HALL. G. 369. Comm. Gott. T. I. p. 201. Centaurea calycibus ciliatis, ciliis setaceis reflexis, foliis integerrimis. GMEL. Sib. II. p. 100. Tab. 45. f. 1. 2.

Calcitrapae. Squamis calycinis spinosis.

H. Centaurea (*Rhoponticum*) calycibus glabris, simplicissime & breviter spinosis, foliis dentato pinnatifidis. HALL. Gott. 364. ^{calcarantia} LINN. Sp. 46. Jacea foliis

foliis eleboraceis, altissima, flore purpureo. I. R. H. Nostrae plantae folia omnia glabra, & laevia sunt. A.

H. Centaurea (*Rhaponticum*) calycibus hirsutis, simplicissime spinosis. HALL. G. 365.

H. Centaurea calycibus palmato-spinosis subaequalibus, foliis decurrentibus sinuatis spinulosis. ^{nanifolia} LINN. Sp. 37. H. Vpf. 272. Calcitrapa caulibus amplissime alatis, spinis calycinis palmatis. HALL. G. 367. Jacea Cyanoides, caule alato. HERM. Par. 189. ic. In Creta. A.

H. Centaurea calycibus duplicato-spinosis, foliis lanceolatis decurrentibus subdentatis, capitulis subsessilibus lateralibus. LINN. H. Vpf. 10. *Solstitialis*. LINN. Sp. 41. Calcitrapa lutea vulgaris, caule alato. VAILL. HALL. G. 366. Flores saepe conferti, foliis linearibus succinhti. A.

H. Centaurea calycibus palmato-spinosis, foliis decurrentibus spinulosis repando-dentatis. ^{sonchi folia} LINN. Sp. 32. Jacea laciniata sonchi folio. B. Prodr. 128. HERM. Par. T. 675. Carduus maritimus canescens, alato caule. I. R. H.

H. Centaurea calycibus duplicato-spinosis lanatis, foliis semidecurrentibus integris sinuatisque. ^{crispifolia} LINN. Sp. 39. Calcitrapa foliis linguiformibus, capitulis tomentosis. HALL. G. 366. Calcitrapa lutea, alato caule, capite eriophoro. VAILL. Art. Par. 1718. p. 212.

Centaurea (*Calcitrapa*) calycibus sessilibus duplicato-spinosis, aculeis rigidissimis, foliis pinnatifidis

dis linearibus dentatis. ^{Calceolaria} LINN. Sp. 40. HALL. H. 689.
Carduus stellatus. C. B. *Carduus muricatus* vulg.
Calcitrapa dictus. CLVS. VII. Circa Wernigerode.

Centaurea (Calcitrapa) foliis semipinnatis, spinis
 calycinis praelongis ciliatis. HALL. G. 365.

Centaurea benedicta p. 390.

II. CORYMBIFERAE.

Flosculi rubulosi breves, aut omnes androgyni
 limbo quinquesido, aut in ambitu feminini fistulosi,
 in capitulum planum, latum, non ventricosum
 collecti.

α. Semine nudo.

TANACETVM. LINN. 848.

Tanacetum foliis pinnatis, pinnis semipinnatis
 serratis, floribus umbellatis. HALL. H. 694. LINN.
 Sp. 7 *Tanacetum vulgare tutum*. C. B. I. R. H.
 BLACKW. T. 464. Ad ripas fossarum.

β. Var. foliis crispis. C. B.

γ. *Tanacetum* foliis ovatis integris serratis.
 LINN. Sp. 8. *Balsamita mas*. BLACKW. T. 98. Men-
 tha hort. *corymbifera*. C. B. In Narbona, P.

δ. *Tanacetum* foliis rigidis integris dentatis, sca-
 po unifloro. HALL. Gott. 370. *Chrysanthemum*
flosculis omnibus uniformibus hermaphroditis. LINN.
 Sp. 14. *Bellis spinosa*. ALPIN. Exot. T. 326. In
 Africa.)

ARTEMI-

ARTEMISIA. LINN. 849.

H. Artemisia caule fruticoso ramosissimo; foliis caulinis pinnato-multifidis linearibus, floralibus indivisis, spicis secundis reflexis. ^{Sanctonicum} LINN. Sp. 3. GMEL. Sib. II. p. 115. T. 51. *Planta fruticosa, foliis multifidis, pinnulis remotis, linearibus longis, capitulis subcylindricis sessilibus dense spicatis, squamis calycinis ora membranaceis, flosculis omnibus androgynis, paucioribus, vix ultra quinque.* Fr.

H. Artemisia foliis ramosissimis setaceis, caule recto suffruticoso. ^{Abrotanum} LINN. Sp. 2. *Abrotanum marginatifolium maius.* C. B.

H. Artemisia foliis planis longe petiolatis duplicato-pinnatis, caule procumbente laxè spicato. HALL. H. 6. ^{campestris} LINN. Sp. 4. *Abrotanum campestre.* C. B. Ad vias in *Thuringia. Marchia.*

H. Artemisia foliis pinnatis, caulibus adscendentibus hirsutis, floribus globosis cernuis subsolitariis; receptaculo villoso. ^{europaei} LINN. Sp. 8. *Fl. Suec. ed. alt.* n. 733. *Absinthium pumilum palmarum minus, argenteo sericeoque folio.* BOCC. Mus. II. T. 71. In *Siberiae & Sueciae* rupibus. Fr.

Artemisia foliis incanis pinnatis, pinnis bilobis; floribus subglobosis pendulis secundis: receptaculo villoso. ^{Absinthium} LINN. Sp. 12. HALL. H. 4. *Absinthium.* BLACKW. T. 17. *Absinthium Ponticum f. Romanum officinarum, Dioscoridis.* C. B. I. R. H. Passim in ruderosis spontanea quasi. P.

H. Artemisia foliis bipinnatis subtus tomentosis nitidis, pinnis transversis, racemis simplicibus, recepta-

receptaculo villoso. ^{*tanacetifolia*} LINN. Sp. 11. *Artemisia radice perenni, foliis pinnatis pinnatifidis incisis serratis, calycibus subrotundis viridibus nutantibus.* GMEL. Sib. II. p. 123. T. 58.

H. *Artemisia foliis simplicibus tomentosis obtusis lanceoatis; inferioribus cuneiformibus trilobis, receptaculo villoso.* ^{*chinesica*} LINN. Sp. 17. *Artemisia foliis radicalibus a caulinis diversis.* GMEL. Sib. II. p. 61. f. 1. 2. Fr.

Artemisia foliis pinnatifidis planis incisis, subtus tomentosis, racemis simplicibus, floribus ovatis erectis, radio quinquefloro. ^{*vulgaria*} LINN. Sp. 13. HALL. H. 5. *Artemisia vulgaris.* C. B. 1. R. H. BLACKW. T. 431. *rubra.* TABERN. p. 7.

H. *Artemisia foliis multipartitis tomentosis, floribus subrotundis nutantibus: receptaculo nudo.* ^{*montica*} LINN. Sp. 9. *Abinthium Ponticum tenuifolium incanum.* C. B. In Hungaria. Sibiria. P.

H. *Artemisia inodora, foliis incanis laxè pinnatis, pinnis longis linearibus, floribus dense spicatis erectis.* HALL. Gott 373. *Racemi plurimi erecti, cauli fere paralleli, floribus congestis tecti, calyce laxo, ex viridi fusco.*

H. *Artemisia foliis linguiformibus tomentosis, integris, lateralibus saepe palmatis, floribus cylindricis spicatis cernuis.* ^{*cerulea*} LINN. Sp. 15. *Abinthium marit. lavendulae folio.* C. B. *Abrotanum latifolium varius, artemisiae folio.* COLUMN. Ecpbr. II. T. 76. Ad littora maris.

H. *Artemisia* foliis lineari-lanceolatis integerrimis glabris. ^{*Tanacetum*} LINN. *Sp.* 16. GMEL. *Sib.* II. p. 126. T. 59. 60. Abrotanum lini folio acriori & odorato. J. R. H. Draco herba. DOD. *Pempt.* 709. In *Sibiria*. P.

H. *Artemisia* lini folio hirsuto, inodorum. AMMANN. *Ruth.* 147. Num satis a priori diversa?

COTVLA. LINN. 868.

H. *Cotula* foliis lanceolato-linearibus amplexicaulibus inferne dentatis. ^{*Coronopifolia*} LINN. *Sp.* 2. Ananthocyclus *Coronopi folio*. DILL. *Elrh.* 27. T. 23. f. 36. In *Frissiae* inundatis. A.

H. *Cotula* receptaculis subtus inflatis turbinatis. ^{*marginata*} LINN. *Sp.* 4. *Chamaemelum Aethiop. lanuginosum*. BREYN. *Cent.* 148. T. 73. A.

CARPESIVM. LINN. 853.

H. *Carpesium* floribus terminalibus. ^{*cerneuus*} LINN. *Sp.* 1. p. 859. Tanacetum flore singulari nutante, foliis longioribus cincto. HALL. H. 693. Gott. 371. Alter *cernuus*. COLVYN. *Ecpbr.* 251. it. In *Helvetia*. *Italia*. A.

SANTOLINA. LINN. 847.

H. *Santolina* pedunculis unisporis, foliis quadrifariam dentatis. *Chamaecyparissus*. LINN. *Sp.* 1. Abrotanum femina vulgare, foliis teretibus. G. P. CLVS. 341. In *Europa* australi.

H. *Santolina* pedunculis unisporis, foliis linearibus. ^{*chamaecyparissus*} LINN. *Sp.* 2. Abrotanum femina foliis. Rorisma-

rismarini. C. B. Abrotanum femina Narbon. CLVS.
342. Fr. *Athanasia*.

H. Santolina corymbis simplicibus coarctatis, foliis pinnatifidis dentatis. LINN. Sp. 4. Elichrysum inodorum glabrum Coronopi folio annuum. HERM. H. Lugdb. T. 227. Chrysanthemum corymbiferum. TRIVME. Obs. T. 86. Caulis ramosissimus. Folia pinnata, pinnis distantibus lineari-lanceolatis, apice saepe trifidis. Capitula oblonga; calycis squamae flavescentes. *Athanasia annua* Linn. p. 9. 122.

ANACYCLVS. LINN. 869.

H. Anacyclus foliis decompositis linearibus, lacinis divisis teretiusculis acutis, floribus flosculosis. LINN. Sp. 3. Chrysanthemum *Valentinum*. CLVS. 332.

β. Semine papposo.

GNAPHALIVM. LINN. 850. Filago.

HALL.

Gnaphalium sexibus tota planta distinctis. HALL. n. 1. *Gnaphalium caule simplicissimo, corymbo simplici terminali, sarmentis procumbentibus*. LINN. Sp. 1.

α. *Androgyna*, flore breviori. Elichrysum mont. flore & folio rotundiore. I. R. H.

β. *Femina*, flore longiori. Elichrysum mont. folio & flore longiore. I. R. H. In collibus siccis apricis.

H. *Gnaphalium* foliis lingulatis subtus tomentosis, caule umbella lactea terminato. HALL. H. 3. *Gnaphalium foliis lineari-lanceolatis acuminatis at-*

ternis

cornis, caule superne ramoso, corymbis fastigiatis.
^{*argenteum*} LINN. Sp. 3. Gnaph. *Americanum*. CLVS. 327.

H. Gnaphalium foliis amplexicaulibus ^{*perforatis*} integerrimis acutis subtus tomentosis, caule ramoso. LINN. Sp. 5. Gnaph. Afr. *latifolium foetidum, capitulo argenteo*. COMMEL. II. T. 56. VOLKAM. Fl. Nor. 194. ic. β. flore aureo. P.

H. Gnaphalium foliis lanceolatis, caule tomentoso ramoso, floribus terminalibus glomeratis conicis. ^{*obtusifolium*} LINN. Sp. 8. Elichrysum *obtusifolium, capitulis argenteis conglobatis*. DILL. Elth. T. 108. P.

H. Gnaphalium foliis lineari-lanceolatis, caule inferne ramoso, corymbo decomposito terminali. ^{*pubescens*} LINN. Sp. 20. Elichrysum Afr. *folio oblongo angusto, flore rubello, postea aureo*. DILL. Elth. T. 167. f. 127. P.

Gnaphalium foliis lingularis, caule herbaceo simplicissimo, floribus umbellatis flavis, flosculis omnibus androgynis. HALL. G. 376. Gnaphalium. ^{*argenteum*} LINN. 19. Stoechas *citrina latifolia offic.* C. B. Circa Rheinhausen & über der Papiermühle.

H. Gnaphalium foliis confertis angusto-lanceolatis, caule fruticoso: corymbo composito. ^{*orientale*} LINN. Sp. 13. Elichrysum Afr. *frutescens, angustis & longioribus foliis incanis*. COMMEL. II. T. 55. Fr.

H. Gnaphalium foliis linearibus, caule fruticoso ramoso, floribus corymbosis cylindricis. ^{*frutescens*} LINN. Sp. 24. Stoechas *citrina tenuifolia altera*. I. B. III. 155. In Europa calidiori.

Gnaphalium caule ramosissimo diffuso, floribus confertis terminalibus fuscis. ^{*filaginifolium*} LINN. Sp. 30. HALL. H. 9. Elichrysum *aquat. minus, capitulis foliatis*. I. R. H. Ad fossas, in agris vdis.

Gnaphalium caule simplicissimo, floribus spicatis sparsis axillaribus. *sylvaticum*. LINN. Sp. 31. HALL. H. 5. Elichrysum *spicatum*. I. R. H. In sylvis.

Filago.

Gnaphalium caule erecto ramosissimo, glomerulis ad alas foliorum longe sessilibus. HALL. H. 11. Gnaph. *paniculatum hirsutum*. ^{*Filago arvensis*} LINN. Sp. 32. Filago *incana tomentosa erecta*. VAILL. Locis aridis, e. g. in colle prope *Bovenden*.

Gnaphalium caule erecto ramoso, foliis pennatis, glomerulis distitis in summo caule sessilibus. HALL. H. 12. ^{*fil. montana*} *montanum*. LINN. Sp. 12. Filago *minor*. I. R. H. In sabulosis, v. g. *M. Hagen*.

Gnaphalium caule dichotomo erecto, capitulis globosis axillaribus terminalibusque. ^{*Fil. germanica*} LINN. Sp. 35. HALL. H. 10. Gnaph. *vulg. maius*. C. B. Filago *f. herba impia*. I. R. H. Locis ficcis & inter stipulas.

H. Gnaphalium foliis alternis linearibus acutis planis, pedunculis longissimis unisfloris. *saxatile*. LINN. Sp. 36. Elichryso *sylvestri similis, flore oblongo*. C. B. Prodr. T. 123. ^{*Fil. saxatile*} *saxatile* Linn.

H Gnaphalium (*Filago*) foliis ovatis undulatis, pedunculis unisfloris. HALL. G. 378. Elichrysum *syn. latifolium, flore magno singulari*. I. R. H.

BACCHARIS, LINN. 853.

H. Baccharis foliis lanceolatis serratis, floribus cymboſis, caule fruticoſo. LINN. *Sp.* 1. Erigeron. HALL. *G.* 379. n. 1. Conyza frut. ^{sua folia} foliis angustioribus nervosis. FEVILL. *Peruv.* T. 37. Senecio *Afr.* arbor. folio serrato. BOERH. In *America*. Fr.

H. Baccharis foliis obovatis, ſuperne emarginato-natis glaucis, floribus in alas congeſtis, caule fruticoſo. ^{halimifolia} LINN. *Sp.* 4. Erigeron HALL. *G.* 379. Senecio *Virginianus* arbor. atriplicis folio. HERM. *r. Bat.* T. 225. In *Virginia*. Fr.

ERIGERON, LINN. 855.

Erigeron floribus ex alis foliorum paniculatis, ſculis femininis omnibus lingulatis. HALL. *H. n.* 4. Erigeron caule floribusque paniculatis. ^{Canadense} LINN. *Sp.* 3. Conyza annua acris alba elatior, *Linariae* foliis. MCC. *Sic.* T. 46. Exoticae quidem originis nunc omnibus ſylvis frequentiffima. A.

H. Erigeron foliis inferioribus dentato-laciniatis, ſuperioribus integris. LINN. *Cliff.* 407. LINN. *Sp.* 4. Senecio *Bonariensis* purpurascens, foliis imis *Coniopi.* DILL. *Elth.* T. 257. f. 334. In *America* ſtrali. A.

H. Erigeron foliis lingulatis, floribus ſolitariis et ramos ſeſſilibus. HALL. *G.* 380. Erigeron ^{nuculea} ramos lateralibus multifloris, calycibus squarrosis. LINN. *Cent. Pl.* I. n. 81. Virga aurea minor, foliis *atinosis* & graveolentibus. I. R. H.

Erigeron pedunculis alternis uniſloris alaribus. LINN. *Sp.* 6. HALL. *H.* 3. Conyza coerulea. TA-



BERN. 861. After *arv. coeruleus acris*. I. R. H. In ficcis apricis. A.

CONYZA. LINN. 854.

Conyza foliis ovato-lanceolatis, floribus laxo corymbosis. HALL. H. 704. *squarrosa* LINN. Sp. 1. *Conyza maior vulgaris*. C. B. In M. *Pless* aliisque collibus ficcis. P.

H. *Conyza saxatilis* p. 202.

CHRYSOCOMA. LINN. 845.

H. *Chrysocoma herbacea*, foliis linearibus glabris, calycibus laxis. ^{*linifolia*} LINN. Sp. 4. *Osyris Austr.* CLVS. 325. *Conyza Linariae folio*. I. R. H. In *Helvetia*. P.

H. *Chrysocoma fruticosa*, foliis linearibus dorso decurrentibus. ^{*arvensis*} LINN. Sp. 1. *Coma aurea Afr. fruticans*, foliis *Linariae angustis*, maior. COMMEL. H. II. T. 45. In *Africa*.

TVSSILAGO. LINN. 856. *Petasites*, HALL. 706.
TOVRN.

Tussilago scapo imbricato unifloro, foliis subcordatis angulatis denticulatis. ^{*parviflora*} LINN. Sp. 3. *Petasites scapo unifloro, flosculis in ambitu lingulatis*. HALL. H. 6. *Tussilago s. Farfara*. I. R. H. BLACKW. T. 204. In argillosis vdis. P.

Tussilago (*Petasites*) ^{*anthelm.*} thyrsis ovato, flosculis omnibus androgynis. LINN. Sp. 6. HALL. H. 7. *Petasites maior & vulgaris*. C. B. BLACKW. T. 222. Ad *Leinam* & fossas. P.

Tussi-

Tussilago (*Petasites*) *thyrso oblongo*, flosculis
femineis nudis plurimis, paucissimis androgynis.
^{*Agrostis*} LINN. Sp. 6. HALL. H. 3. *Petasites maior*, flosculis
radunculis longis insidentibus. DILL. Elth. T. 230.
f. 297. *Bey der Weender Papiermühle; in der*
Contrescarpe.

^{*Tussilago alba*}

CACALIA. LINN. 84f.

H. *Cacalia foliis reniformibus acutis denticulatis*.
LINN. Sp. 10. *Cacalia*. HALL. H. 705. *Cacalia*
glabro folio. CLVS. CXV. In *Helvetiae montosis*.

H. *Cacalia* (*Kleinia*) *caule fruticoso*, foliis car-
nosis ovato-oblongis, petiolis basi linea triplici de-
ductis. ^{*Anteuphorbia*} LINN. Sp. 2. H. Cliff. 395. *Anteuphorbium*.
DOD. Pempt. 378. DILL. Elth. T. 55. f. 2. 3. Fr.

H. *Cacalia* (*Kleinia*) *caule fruticoso levi ventri-*
oso, foliis lanceolatis planis, petiolorum cicatri-
cibus obsoletis. ^{*Kleinia*} LINN. Sp. 3. Cliff. 395. *Cacalian-*
hemum folio Nerii glauco. DILL. Elth. T. 54.
f. 2. Fr.

H. *Cacalia* (*Kleinia*) *caule fruticoso terete*, fo-
liis carnosis compressis lanceolatis. ^{*Ficoidis*} LINN. Sp. 2.
Senecio Afr. arborescens, *Ficoidis folio & facie*.
COMMEL. rar. T. 40.

EVPATORIVM. LINN. 842.

Eupatorium foliis plerumque ternatis. HALL. H.
84. *Eupat. foliis digitatis*. ^{*annabina*} LINN. Sp. 10. *Eupat.*
annabinum, G. B. I. R. H. *Eup. Avicennae*. BLACKW.
T. 110. Ad rivas, P.

H. Eupatorium foliis ovatis ferratis petiolatis, ^{cili} glabro, calycibus simplicibus. Eupatorium *Scroph-*
lariae folio, floribus albis. HALL. G. 381. Ag-
eratum ^{albidum} LINN. Sp. 3. Valeriana *Vrticae folio, flore alb*,
CORNV. T. 21. P.

Semine aristis coronato.

AGERATVM. LINN. 843.

H. Ageratum caule piloso, foliis petiolatis hirs-
tis ovatis crenatis, stylis flore longioribus reflexis,
Ageratum. ^{longifolium} LINN. 1. *Flosculi albi.* An Eupato-
rium *Betonicae folio.* HERM. Par. ic. p. 162? Sed
folia sessilia ibi pinguntur. ^{non defuit}

H. Ageratum caule subhirsuto, foliis petiolatis
oblongis crenatis glabris ora ciliatis, stylis tubum
floris vix superantibus. An Eupatorium *Amer. her-*
baceum, Melissa folio, villosum, flore coerule.
HALL. G. 323?

H. *Ageratum albidum.*

XERANTHEMVM. LINN. 851.

H. *Xeranthemum* foliis lanceolatis patentibus, re-
ceptaculo paleaceo, seminum pappo quinqueseto.
^{nastrum} LINN. Sp. 1. *Parmica Austriaca.* CLVS. XI.

BIDENS. LINN. 840.

Bidens foliis tripartito divisis, calycibus subfo-
lios, seminibus erectis. ^{tripartita} LINN. Sp. 1. HALL. H. 709.
n. 1. *Cannabina aquat. folio tripartitim diviso.* C. B.
Ad fossas. β . Var. flore radiato.

Bidens

Bidens foliis longe ellipticis indivisis, seminibus erectis, floribus cernuis. HALL. H. ^{cernuis} LINN. Sp. 2.
Bidens folio non dissecto. 1. R. H.

β. Varietas flosculis in radio lingulatis, semine abortivo. *Coreopsis foliis lanceolatis serratis oppositis amplexicaulibus.* ^{Bidens} LINN. Sp. 7.

h. *Bidens* foliis pinnatis subpilosis, corollis flosculosis, calycibus simplicibus, seminibus divergentibus. ^{neolob.} LINN. Sp. 5. *Bidens latifolia hirsutior, semine angustiore radiato.* DILL. Elth. T. 43. f. 51.

h. *Bidens* foliis ovatis & tripteris, caulibus hirtis brachiatis. HALL. G. 383. *Bidens foliis ovatis serratis: inferioribus oppositis, superioribus ternatis; intermedio maiore.* ^{subulata} LINN. Sp. 11. c. descr. A.

RADIATAE.

Flores in disco tubulosi, quinquefidi, androgyni; in radio lingulati, feminini aut abortivi.

I. Placenta paleacea, semine paleis coronato.

VERBESINA. LINN. 874.

h. *Verbesina* foliis alternis decurrentibus vndulatis obtusis. ^{alata} LINN. Sp. 1. *Chrysanthemum Curassavicum, alato caule, floribus aurantiis.* HERM. Par. T. 125. In Surinamo. A.

h. *Verbesina* foliis oppositis lanceolatis serratis, floribus alternis geminis subsessilibus. *prostrata.* LINN. Sp. 8. *Eupatorio-phalacron menthae arvensis folio.* DILL. Elth. T. 113. f. 138. Eandem icone addita descripsi in *Comm. Soc. Gott. T. III. Tab. XI.*

COREOPSIS. LINN. 879.

^{Trientalis} H. Coreopsis foliis subternatis integerrimis.
LINN. *Spec.* 2. H. *Vpf.* 269. *. HALL. G. 385.
Chrysanthemum *Virgin.* folio acutiore lacvi trifol-
liato, *f. anagyridis folio.* MORIS. III. S. VI. T. 3.
f. 44. P.

H. Coreopsis foliis lanceolatis ferratis alternis per-
tiolatis decurrentibus. ^{alternifolia} LINN. *Sp.* 8. HALL. G. 385.
Chrysanthemum *caule alato Canadense.* ZANON.
T. 58.

^{Coreopsis} ~~Coreopsis~~ *3id. 1. 407*

HELIANTHVS. LINN. 877.

H. Helianthus radice annua, flore amplissimo
nutante. ^{annuus} LINN. *Cliff.* 419. foliis omnibus cordatis,
nervis pone basin unitis : extrorsum denudatis.
LINN. *Sp.* 1. Corona solis. I. R. H. A.

H. Helianthus foliis inferioribus cordatis : nervis
prope basin unitis denudatis : superioribus ovatis,
multiflorus. LINN. *Sp.* 2. Hel. *radice tereti inle-*
xa perenni. *Cliff.* 419. Corona Solis *minor femina.*
TABERN. *ic.* 764. P.

H. Helianthus foliis ovato-cordatis : nervis in-
fra folium unitis, radice ^{tuberosa} tuberosa. LINN. *Sp.* 3.
Corona Solis, *flore parvo, radice tuberosa.* I. R. H.
In *Brassilia*.

H. Helianthus foliis lanceolatis scabris, caule
stricto : inferne giabro, radice fusiformi. ^{gigantius} LINN. *Sp.*
6. Helianthus *maior Virginianus.* ZANON. *Hist.*
T. 58. P.

RVDBECKIA. LINN. 878.

H. Rudbeckia foliis pinnatis, extrema pinna amplissima. HALL. G. 386. *foliis compositis laciniatis*. LINN. Sp. 1. Obeliscotheca *Hydrophylli folio, foliorum lobis latioribus*. VAILL. β. Variet. *lobis angustioribus*. P.

H. Rudbeckia foliis asperis ovatis dentatis, radii petalis emarginatis. ^{Linna} LINN. Sp. 3. Obeliscotheca *integrifolia, radio aureo, umbone atrorubente*. DILL. Elth. T. 218. f. 285.

H. Rudbeckia ^{Linna} foliis oppositis hirsutis ovato-acutis, calyce imbricato cylindrico, radii petalis pistillatis. Sic interim definire liceat plantam, quae, si nulla, noui generis nomen meretur. *Calyx cylindricus uncialis, squamis rotundatis dense imbricatus. Flores radii apice emarginato, immarcescibiles. Discus conicus. Semina omnia longa triquetra, quae in radio, nuda, cum petalo cartilagineo unita; quae sub disci floribus, simplici aut duplici longiori trifida coronata; squamae calycis longitudine, semina distinguunt. A Rudbeckia, cuius habitum refert, petalis radii femininis & calyce differt. Vide iconem sub fine libri.* *Linna pauciflora* (Lutca) Linn. 2 p. p. 126.

H. *Linna multiflora* (coccinea - purpurascens) ibid. v. Icon in Linna 2 p. p. 17. *Linna multiflora* (Lutca) Linn. 2 p. p. 17.

SIEGESBECKIA. LINN. 873.

H. Siegesbeckia LINN. Cliff. 412. T. 23. *. H. 7 p. 267. HALL. G. 385. *petiolis sessilibus, calycibus incrassatis*. ^{orientalis} LINN. Sp. 1. Bidenti similis; foliis latissimis ferratis. BUXB. Cent. III. T. 52. In Media. A.

BUPHTHALMVM. LINN. 876.

H. *Bupthalmum* foliis oppositis lanceolatis, petiolis bidentatis. ^{*Leucocaulis*} LINN. *Sp.* 1. *Asteriscus frutescens*, *Leucoii foliis sericeis & incanis*. DILL. *Elth.* T. 38. f. 44. In *Virginia*. Fr.

H. *Bupthalmum* calycibus acute foliosis, ramis alternis, foliis lanceolatis amplexicaulibus integerrimis. *spinosum*, LINN. *Sp.* 2. *Aster.* I. CLVS. XIII. *Aster luteus, foliis ad florem rigidis*. C. B.

H. *Bupthalmum* calycibus obtuse foliosis sessilibus axillaribus, foliis oblongis obtusis. ^{*aguticatis*} LINN. *Sp.* 3. *Aster. annuus odoratus, Lusitanicus*. I. R. H. ZANON. *Hist.* T. 24. In *Creta. Lusitania*. A.

H. *Bupthalmum* foliis lineari-lanceolatis, serratis, villosis, calycibus nudis. HALL. *H.* 711. n. 2. *salicifolium*. LINN. *Spec.* 5. *Aster* III. *Austr.* I. CLVS. XIII. *Aster luteus maior, foliis succisae*. C. B. In *Helvetia*.

H. *Bupthalmum* calycibus foliosis, foliis cordatis serratis trinerviis basi hinc brevioribus. ^{*Helianthoides*} LINN. *Sp.* 7. HALL. *G.* 388. *Chrysanthemum Scrophulariae folio Americanum*. PLVKN. T. 99. f. 1. P.

— *Bupthalmum* foliis incanis pinnatis, pinnis argute serratis. HALL. *H.* 3. *Anthemis*. ^{*linctaria*} LINN. *Sp.* 16. *Chamaemelum Chrysanthemum*. I. B. III. p. 122. *Bupthalmum Tanacetii minoris foliis*. C. B. I. R. H. In sylvis & collibus.

H. *Bupthalmum* (*Anthemis*) caulibus simplicibus unifloris decumbentibus, foliis pinnato ^{*multifloris*} ~~multifloris~~.

LINN.

LINN. *Sp.* 12. *Pyrethrum officinarum*. BLACKW. *T.* 390. DOD. *Pempt.* 327.

ACHILLEA. LINN. 871.

H. Achillea foliis ellipticis obtusis acute ferratis fasciculatis, floribus umbellatis. HALL. *G.* 389. LINN. *Sp.* 2. ^{*serratis*} *Ageratum foliis ferratis*. C. B. *Ptarmica lutea suaveolens*. I. R. H. *Balsamita minor*. DOD. *Pempt.* 295. In Gallia Narbonensi.

Achillea foliis integris lanceolatis minutissime ferratis. ^{*Ptarmica*} LINN. *Cliff.* 412. HALL. *H.* 1. LINN. *Sp.* 10. *Ptarmica vulgaris*. CLVS. XII. BLACKW. *T.* 276. *foliis longis serratis, flore albo*. I. R. H. In pratis. v. g. *gegen die Masckmühle*. Var. hort. *flore pleno*.

Achillea foliis punctatis bipinnatis nudis, laciniis linearibus dentatis. ^{*Millefolium*} LINN. *Sp.* 14. HALL. *Helv.* 8. *Millefolium*. BLACKW. *T.* 18. *vulgare album*. C. B. I. R. H. β *vulgare purpureum minus*. I. R. H. Prior var. vbique in gramineis; altera potius *hortensis*. P.

Achillea foliis bipinnatis, pinnis primis remotis, vltimis confertis, pinnulis semipinnatis. HALL. *H.* 9. ^{*nobile*} LINN. *Sp.* 15. *Millefolium nobile*. I. R. H. Achillea *Mill. folia odorata*. I. B. III. p. 140. In M. *Altenstollberg*; circa *Stollberg & Kelbra* in muris. P.

H. Achillea pinnulis foliorum aequalibus confertis tomentosis, pinnulis imorum trifidis, summorum simplicibus. HALL. *H.* 11. ^{*tomentosa*} LINN. *Sp.* 4. *Millefolium tomentosum luteum*. C. B. *Stratiotes millefolia flavo flore*. CLVS. 330. In *Vallesia*. P.

H. Achillea foliis pinnatis, pinnis semipinnatis, foliolis lanceolatis incisive serratis subtus lanigeris. ^{montana} LINN. Sp. 5. Millefolium Orient. foliis Tanacetii incanis, radiis pallide luteis. BOERH. Lugdb. I. p. 212. I. R. H. Cor. 37.

ANTHEMIS. LINN. 870. Chamaemelum. T.

H. Anthemis caule ramoso, foliis pinnato-multifidis setaceis, calycibus villosis pedunculatis. ^{valentina} LINN. Spec. 13. Santolina foliis repetito-divisis planis, HALL. G. 375. Cotula flore luteo radiato. I. R. H.

H. Anthemis foliis pinnato-multifidis, foliorum apicibus subspinosis. ^{alpina} LINN. Sp. 2. Chamaemelum Leu anthemum Hisp. magno flore. C. B. Prodr. 70. In Gallia Narbonensi.

H. Anthemis foliis planis pulposis pinnatis, pinnis dentatis, caule ramoso. HALL. G. 391. ^{maritima} LINN. Sp. 3. Chamaemelum maritimum. I. B. III. p. 122.

H. Anthemis foliis pinnato-multifidis planis; laciniis linearibus acutis trifidis, pedunculis longissimis. ^{reclama} LINN. Sp. 10. Chamaemelum Orient. absinthii folio. I. R. H. Cor. 37.

H. Anthemis foliis pinnato-compositis linearibus acutis subvillosis. ^{nobile} LINN. Sp. 7. Chamaemelum foliis lanuginosis pinnatis, pinnis capillariter multifidis brevioribus. HALL. H. 2. Chamaemelum odoratum. DOD. Pempt. 270. Chamaem. nobile f. Leucanthemum odoratius. C. B. I. R. H. In Helvetia.

β. Var. flore pleno & semipleno.

H. Chamaemelum Chium vernum, folio crassiore, flore magno. I. R. H. Cor. 37. Flos maior. Folia lata & robusta; LINNAEVS tamen pro varietate prioris habet. *Anthem. Chier. L. with. v. 1. p. 76.*

Anthemis (*Chamaemelum*) foliis pene glabris, bipinnatis, pinnis latiusculis linearibus. HALL. H. 3. Anth. receptaculis conicis, paleis setaceis, seminibus nudis; ^{Cotula} LINN. Sp. 9. *Cotula foetida*. BLACKW. T. 67.

Anthemis tinctoria. H. Anth. *Pyethum* p. 40.

MATRICARIA. HALL. 718. LINN. 867, Chrysanthemum. LINN. 866.

H. Matricaria (*Chrysanthemum*) foliis lanceolatis superne ferratis vtrique acuminatis. ^{serotina} LINN. Sp. 2. Bellis Amer. procerior serotina ramosa; flore amplissimo. PLVKN. T. 17. f. 2.

Matricaria foliis imis petiolatis; caulinis amplexicaulibus ferratis. HALL. H. 3. *Chrysanthemum*. ^{lanceolatum} LINN. Sp. 3. Bellis sylv. caule folioso, maior. C. B. Bellis maior. CAMER. In gramineis.

Matricaria foliis glaucis amplexicaulibus, profunde dentatis. HALL. G. 393. *Chrysanthemum segetum*. LINN. Sp. 10. CLVS. 334. Chrysanth. folio minus secto glaucum. I. B. III. 103. Vbique in arenosis circa Hefeld, versus Seeburg: in ducatu Celsensi. A.

H. Matricaria (*Chrysanthemum*) foliis pinnatifidis incisicis extrorsum latioribus. ^{Cotula} LINN. Sp. 11. Chrysanth. maius folio valde laciniato, flore croceo. I. B. III. p. 104. Chrysanth. Creticum. CLVS. 335. β. Var. semisofsculis fistulosis. A.

H. *Chrysanthemum pleurolophum* p. 396.

H.

Chrys. corymbiferum.

H. *Chrys. fruticosum* p. 414.

RADIATAE.

Matricaria

H. *Matricaria* foliis pinnatis, pinnis distantibus parallelis semipinnatis. HALL. G. 392. *Matricaria Orient. folio Tanaceti, flore suaverubente.* I. R. H.

Matricaria foliis compositis planis, foliolis ovatis incisis, pedunculis ramosis. HALL. H. 2. ^{*Parthenium*} LINN. Sp. 1. *Matricaria.* BLACKW. T. 192. *vulgaris.* C. B. I. R. H. In cultis & ruderalis. In horto radios saepe deponit. — *in fine pinn*

Matricaria foliis pinnatis, pinnis laxae & capilliter multifidis, receptaculis haemisphaericis. HALL. H. 1. *Matricaria Chamomilla.* LINN. Sp. 3. *Chamaemelum vulgare* f. *Leucanthemum Dioscoridis.* C. B. I. R. H. BLACKW. T. 296. In agris antequam Albanam ad radicem M. Heymberg. Circa Nithbergam abunde.

PYRETHVM. HALL. 720. Chrysanthemi

Sp. LINN.

Pyrethrum foliis pinnatis, pinnis semipinnatis, serratis. HALL. H. 1. *Chrysanthemum*. ^{*corymbiferum*} LINN. Sp. 12. *Matricaria folio Tanaceti, flore maiore, semine umbilicato.* I. R. H. *Tanacetum inodorum.* L. CLVS. 338. In sylvis Montanis. P.

H. *Pyrethrum* foliis incanis triplicato-pinnatis tomentosis. HALL. G. 393. *Parmica Orient. foliis Tanaceti incani, semisofculis brevioribus.* I. R. H.

H. *Pyrethrum* foliis pulposis, rariter pinnatis, pinulis bidentatis & tridentatis. HALL. Gott. 393. *Chrysanthemum* ^{*vulgare*} LINN. 1. *Leucanthemum Canariense, sapore Pyrethri.* WALTH. Hort. T. 24.

CALEN-

CALENDULA. LINN. 885.

H. Calendula foliis ovatis integerrimis, floribus parvis pallentibus. HALL. H. 722. *Caltha arvensis*. C. B. 1. R. H. In *Helvetia & Thuringia* ad lacum alsum. A. *Calendula arvensis* Linn Sp. 1 622.

H. Calendula foliis ovatis acutis, flore magno aurantio. Calendula *feminibus radii cymbiformibus rechinatis: disci bicornibus* ^{officinalis} LINN. Sp. 1. utramque ungit.

H. Calendula foliis ovato-lanceolatis, dentatis hirsutis, pedunculis superne incrassatis. LINN. Sp. 3. Calendula *humilis* Afr. flore intus albo, extus violaceo. HERM. Lugdb. T. 105. Radii extus dilute violacei. A.

H. Calendula foliis linearibus subdenticulatis & integris, radiis floris distantibus. An Bellis Afr. florum pedunculis foliosis, foliis angustis & integris. COMMEL. H. II. T. 34. A. *Cal. graminifolia* L. Sp. 722.

OSTEOSPERMUM. LINN. 887.

H. Osteospermum foliis oppositis palmatis, petiolis foliosis. HALL. Gott. 395. c. *charact.* LINN. Sp. 1. Chrysanthemum perenne *Virgin. maius, Platani Orient. folio*. MORIS. III. S. VI. T. 7. f. 55. P.

Polymnia tuberosa Linn. Sp. 2. H. Osteospermum foliis ovatis rigidis dentatis. LINN. Sp. 3. Chrysanthemoides Afrum, Populi albae foliis. DILL. Elth. T. 68. f. 79. In *Aethiopia*.

ERIOCEPHALVS. LINN. 890.

H. *Eriocephalus Africanus*. LINN. *Sp.* 926. *Eriocephalus sempervirens, foliis fasciculatis & digitatis*. DILL. *Elb.* T. 110. f. 134. Fr.

Semine paleis coronato.

TAGETES. LINN. 865.

H. *Tagetes* caule subdiviso diffuso. *patula*. LINN. *Sp.* 1. Flos Africanus. DODON. *Pempt.* 255. *Tagetes minor flore fulvo maculato*. DILL. *Elb.* T. 279. f. 361. In Mexico. A.

H. *Tagetes* caule simplici erecto, pedunculis nudis unifloris. LINN. *Spec.* 2. *Tagetes maior, flore luteo pallescente*. VAILL.

β. *flore aurantis, simplici*.

γ. *flore pleno*.

δ. *radiis fistulosis*. *Tagetes Mexicanus, flore fistuloso, simplex*. COLUMN. *Ecpbr.* II. T. 46. f. 1. A.

ARCTOTIS. LINN. 886.

H. *Arctotis* foliis pinnatis, pinnis undulatis bipinnatis, subtus tomentosis. HALL. *G.* 406. LINN. *Sp.* 3. *Anemonospermum Afra, folio Jacobae tenuiter laciniato, flore aurantio pul. herim.* BOERH. *Lugdb.* T. 100. COMMEL. *H.* II. T. 22. Fr.

H. *Arctotis* foliis tomentosis pinnatis, pinna extrema maxima, flosculis radiantibus vicenis. HALL. *G.* 407. LINN. *Sp.* 1. *Anemonospermum Afra, foliis & facie Taraxaci incanis*. BREYN. *ic.* 27. T. 15. A.

HELE.

HELENIA. LINN. 863.

H. Helenia¹¹⁷⁴ foliis decurrentibus. ^{autumnale} LINN. Sp. 1.
 After *Floridanus aureus*, caule alato. PLVKN. T.
 372. f. 5. After *luteus alatus*. CORNVT. Canad.
 T. 63. P.

Semine papposo, placenta nuda.

BELLIS. LINN. 864.

Calyx haemisphaericus imbricatus ex duplici se-
 rie squamarum.

Bellis scapo nudo unifloro, semine villos^{parvulus &}o. LINN.
 Sp. 1. HALL. H. 722. Bellis minor sylvestris. C. B.
 1. R. H. BLACKW. T. 200. In omnis pratis. Semina
 brevi pappo semper coronata. ^{β. var. hort. flore pleno.}

(H. Bellis scapo nudo unifloro, semine papposo.
 Bellidastrium. HALL. Helv. 723. ^{Bellidastrium} Doronicum. LINN.
 Sp. 4. In montibus Helveticis.)

(H. Bellis caule diviso folioso, foliis lanceolatis.
 After LINN. Sp. 19. Bellis ramosa umbellifera. COR-
 NVT. Canad. T. 174. In Canada. A.)

INVLA. LINN. 860.

Calyx latus, laxus. Flosculi plurimi. Antherae
 inferne fetas demittunt.

H. Inula foliis ovatis rugosis, subtus tomentosis
 ferratis, calycum squamis ovatis. ^{Helelenium} LINN. Sp. 1.
 After HALL. H. 6. After *omnium maximus*, *Hele-*
nium dictus. 1. R. H. *Helenium vulgare*. C. B. In
Anglia. Helvetia. P.

Inula foliis radicalibus petiolatis lingulatis, caulinis subamplexicaulibus serratis, floribus corymbosis. After *luteus folio succiferae*. C. B. HALL. G. 397. Conyza III. *Austr.* cl. v s. XX. An Inula ^{o. m. y. x. p.} LINN. Sp. 3. Fufius hanc & fequentem defcripsi in *Comment. Gott. T. V. p. 13. sqq.* Circa Berolinum, Jenam. A.

H. Inula foliis lanceolato-ovatis serratis, asperis, subtus incanis, caule ^{hirs. c. l. p. 12.} tubulifloro. *Comm. Gott. T. V. l. c.* After HALL. H. 5. After *luteus, hirsuto salicis folio*. C. B. 266. Circa Bero:inum. A.

Inula foliis lanceolatis recurvis serrato-ciliatis glabris, floribus folitariis, ramis angulatis ^{hirsuta} LINN. Sp. 7. After HALL. H. 8. After *mont. luteus, salicis folio glabro*. C. B. In M. *Atenstollberg.* P.

Inula foliis oblongis amplexicaulibus, caule hirsuto paniculato, squamis calycinis setaceis. *dyfenterica*. LINN. Sp. 5. After. HALL. H. 3. After *prat. autumnalis, Conyzae folio*. I. R. H. Conyza *media*. C. B. LOB. *ic.* 345. Ad fossas. A.

Inula caule ramosissimo prostrato, foliis undulatis amplexicaulibus, radiis brevissimis ^{Pulicaria} LINN. Sp. 6. After HALL. H. 4. Conyza *pulicaria*. BLACKW. T. 103. After *palustris parvus, globoso flore*. I. R. H. In agris vdis & ad fossas.

H. Inula foliis linearibus carnosis tricuspidatis ^{crithmoideis} LINN. Sp. 11. After *marit. folio tereti crasso tridentato*. I. R. H. *Crithmum Chrysanthemum*. DOD. *Pempt.* 706.

ASTER. LINN. 858.

Calyx oblongus. Flosculi pauciores. Semiflosculi plurimi calyce longiores.

After

After foliis lanceolatis integerrimis carnosif glabris, ramis inaequatis, floribus corymbosis. ^{*Tripolium*} LINN.

Sp. 4. GMEL. Sib. II. p. 187. T. 80. f. 2. Tripodium *maius coerulcum*. C. B. DOD. *Pempt.* 379. In pratis falsis circa *Harste*.

A. After foliis lanceolatis scabris trinerviis integerrimis, supremis ovatis, pedunculis nudiusculis corymbosis, squamis calycinis obtusis. ^{*amelus*} LINN. Sp. 5. After HALL. H. 2. After *atticus coeruleus vulgaris*. C. B. I. R. H. BLACKW. T. 109. DOD. *Pempt.* 266. In M. *Plesi*. P.

H. After foliis omnibus ovato-lanceolatis scabris integerrimis, pedunculis corymbosis, squamis calycinis acutis ciliatis. HALL. G. 396. After *latifolius*, *Tripolii flore*. I. R. H. P.

H. After ramis divaricatis, foliis ovatis serratis, floralibus integerrimis, pedunculis vnifloris nudis. *divaricatus*. LINN. Sp. 6. GRON. *Virg.* 99. In *Virginia*. P.

H. After caulibus asperis, foliis lanceolatis integerrimis semiamplexicaulibus, floribus terminalibus, pedunculis foliosis. ^{*novae Angliae*} LINN. Sp. 15. HALL. G. 396. After *novae Angliae altissimus hirsutus*, floribus amplis purpureo-violaceis. HERM. Par. T. 98. VOLKAM. p. 50. ic. P.

H. After foliis amplexicaulibus integris lingularis glabris, pedunculis foliosis vnifloris, calycibus squarrosis. HALL. G. 396. After *novi Belgii laeifolius, paniculatus, floribus dilute violaceis*. HERM. *Lugdb.* T. 67. An After LINN. 25. ? P.

H. Aster (*Solidago*) foliis lanceolatis venosis amplexicaulibus glabris, pedunculis foliosis, calycis squamis lanceolatis appressis. ^{multiflorus} HALL. G. 400. ¹⁷³⁸ Aster novi Belgii latifolius, paniculatus, floribus saturate violaceis. HERM. Lugdb. p. 67. ic. P.

H. Aster (*Solidago*) foliis ovato-lanceolatis serratis petiolatis, caule paniculato, ramis multifloris. ^{cordifolius} HALL. G. 400. LINN. Sp. 17. Aster latifolius autumnalis. CORNV. Canad. T. 65. P.

H. Aster (*Solidago*) caule præalto ramosissimo, foliis linearibus integerrimis, floribus terminalibus. HALL. G. 1. ^{linearis} LINN. Sp. 7. Aster novae Angliae, *Linariae foliis*, *Chamaemeli floribus*. HERM. Par. T. 95. Flores albi. P.

H. Aster (*Solidago*) foliis linearibus trineerviis, floribus corymbosis. HALL. G. 400. Aster Tripolii flore. I. R. H. P.

H. Aster caule paniculato, pedunculis racemosis, pedicellis foliosis, foliolis linearibus integerrimis. *ericoides*, LINN. Sp. 16. Aster *ericoides dumosus*. DILL. Elth. T. 36. f. 40. P.

H. Aster foliis ovatis angulatis dentatis petiolatis, calycibus terminalibus patentibus foliosis. ^{Chenopodii} LINN. Sp. 29. Aster *Chenopodii folio annuus*, flore ingenti specioso. DILL. Elth. T. 34. f. 38. A.

H. Aster annuus p. 17.

SOLIDAGO LINN. 859.

Calyx oblongus, imbricatus. Semiflosculi pauciores, calyci subaequales. Flores plerumque racemosi.

H. *Solidago* foliis lanceolatis subcarnosis glaberrimis margine scabriusculis, panicula ^{Chamaemeli} corymbosa.

LINN.

LINN. Sp. 1. *Solidago maxima*. CORNVT. *Canad.*
168. *Virga aurea Novboracensis glabra, caulibus*
rubentibus, foliis angustis, glabris. HERM. Fl. 26. A.

H. *Solidago paniculato-corymbosa, racemis re-*
curvis, floribus adscendentibus, foliis trinerviis subser-
ratis scabris. LINN. Sp. 2. *Virga aurea angustifolia, pa-*
nicula speciosa, Canadensi. ^{Canadensis} PLVKN. T. 236. f. 1. P.

H. *Solidago paniculato-corymbosa, racemis re-*
curvis, floribus adscendentibus, foliis enerviis sub-
integerrimis, altissima. LINN. Sp. 3. *Virga aurea*
altissima serotina, panicula speciosa patula. MAR-
TYN. Cent. T. 14. P.

H. *Solidago caule obliquo, spicis secundis, pe-*
dunculis erectis foliolatis ramosis, foliis lanceolatis
integerrimis. ^{Medicinalis} LINN. Sp. 6. HALL. G. 399 *Virga*
aurea Limonii folio, panicula vno versu disposita.
I. R. H. PLVKN. T. 235. f. 2. In Mexico. P.

H. *Solidago caule flexuoso, foliis ovatis acuminatis*
ferratis, racemis lateralibus simplicibus. LINN. Sp. 7.
Virga aurea mont. Scrophulariae folio. PLVKN. T. 235.
f. 3. *Virga aurea Can. Asterisci folio*. HERM. Par. T. 244.

Solidago caule subflexuoso angulato, racemis
paniculatis erectis confertis. ^{Virga aurea} LINN. Sp. 9. HALL.
H. 729. *Virga aurea vulgaris latifolia serrata*.

U. B. I. R. H. BLACKW. T. 169. In fylvis.

^{Virga aurea} *SENECIO*. HALL. 730. LINN. 857. ^{Virga aurea} *Senecio*
varia Othonna. LINN. 888.

Calyx conicus ex squamis linearibus parallelis,
simplici serie in cylindrum conglutinatis, acceden-
tibus ad basin squamulis vagis.

H. Senecio corollis nudis, foliis amplexicaulibus laevibus laceris, caule herbaceo erecto. ^{heranifolius} LINN. Sp. 1. Senecio Amer. altissimus, blattariae f. bieracii folio. HERM. Par. T. 226. A.

Senecio corollis nudis, foliis pinnato-sinuatis amplexicaulibus, floribus sparsis. ^{vulgaris} LINN. Sp. 4. HALL. H. 7. Senecio minor vulgaris. C. B. 1. R. H. Erigeron Senecio. BLACKW. T. 132. In agris & hortis. A.

H. Senecio corollis revolutis, foliis sinuato-dentatis, inferioribus petiolatis, pedunculis trifloris folio longioribus. Senecio triflorus. LINN. Sp. 5. ROYEN. n. 13. Senecio Aegyptius, foliis Matricariae. 1. R. H. A.

H. Senecio corollis revolutis, foliis amplexicaulibus lanceolatis dentatis, squamis calycinis apice purpureis. Senecio lividus. LINN. Sp. 7. H. Vpf. 267. n. 7. Jacobaea Hisp. minus laciniata, petalis brevissimis. 1. R. H. Folia saepe subtus rubent. A.

H. Senecio corollis revolutis, foliis amplexicaulibus apice trilobis ferratis, squamis calycinis sphaecelatis laceris. ^{calobus} LINN. Sp. 8. H. Vpf. 4. c. descr. In Hispania. A.

Senecio corollis revolutis, foliis viscidis pinnatis, pinnis angulatis, squamis calycinis laxis longitudine perianthii. HALL. H. 5. ^{vulgaris} LINN. Sp. 9. Senecio hirsutus viscidus graveolens. DILL. Elzb. T. 258. f. 336. Senecio incanus pinguis. In sylvis caeduis, in primis areis carbonariorum. A.

β. *Jacobaea Senecionis folio incano*. I. R. H. HALL. *Rupp. Fl. Jen. T. III.* Senecio. LINN. *Sp.* 10. Flores minores, folia minime viscida.

Senecio corollis radiantibus, foliis pinnatis, pinnis multiformibus dentatis, floribus corymbosis. HALL. *H.* 3. ^{*Jacobaea*} LINN. *Sp.* 19. *Jacobaea vulgaris laciniata*. C. B. I. R. H. Ad fossias. P.

Senecio corollis radiantibus, foliis pinnatifidis, subtus tomentosis, pinnis linearibus patentissimis, rariter incis. MOEHRING. *Commerc. Nor.* 1743. p. 117. HALL. *H.* 4. ^{*erucifolia*} Senecio LINN. *Sp.* 15. *Jacobaea altissima, foliis Erucæ Artemisiæve similibus*. RVPP. *Fl. Jen.* 176. In sylvis. v. g. *Eulenloch*. P.

H. Senecio corollis radiantibus foliis semiamplexicaulibus, semipinnatis, pinnis aequalibus angulatis, HALL. *G.* 402. ^{*celastri*} LINN. *Sp.* 13. *Jacobaea Afr. annua, floribus suaverubentibus*. VOLK. *Noric. p.* 225. ic. A.

(H. Senecio foliis pinnatifidis tomentosis, laciniis sinuatis, caule fruticoso. ^{*Cissampelos maritima*} *Orthonna* LINN. *Sp.* 9. *Jacobaea maritima*. C. B. Cineraria. DOD. *Pempt.* 642.)

Senecio corollis radiantibus, floribus corymbosis, foliis utrinque lanceolatis, ferratis, glabriusculis. ^{*apocynifolia*} LINN. *Sp.* 23. HALL. *H.* 9. *Solidago Sarra-cenica*. DOD. 141. *Consolida aurea*. TABERN. 556. In sylvis. P.

Senecio corollis radiantibus, foliis ovato-lanceolatis subtus hirsutis, floribus corymbosis. HALL. *H.* 10. ^{*nemorosa*} LINN. *Sp.* 22. *Jacobaea nemorensis, latiore*

bus corymbosis. Doria Orient. *Limonii folio amplissimo*. DALL. *Elch. T.* 105.

ARNICA. LINN. 861.

Arnica foliis ovatis integris, caulinis geminis oppositis. LINN. *Sp.* 1. HALL. *H.* 737. *Doronicum Plantaginis folio, alterum*. C. B. 1. R. H. *Doronicum VI. Pannonicum*. CLVS. XVIII. *Arnica offic.* In omnibus pratis *Hercynicis*. P.

DORONICVM. LINN. 862.

H. *Doronicum foliis cordatis obtusis; radicalibus petiolatis, caulinis amplexicaulibus*. LINN. *Sp.* 1. *Doronicum Romanum*. BLACKW. 239. *Aconitum Pardalianches*. DOD. *Purg.* 305. In montibus *Helvetiae*. P.

H. *Doronicum Pellidistrum* p. 417.

PLANIPETALAE S. CICHORACEAE.

Flosculi omnes aequales lingulati androgyni, apice quinquedentato. Plantae huius classes omnes fauciatae lac stillant, amarae, saponaceae.

I. Semine papposo, placenta nuda.

TARAXACVM. HALL. 739. *Leontodon*. LINN. 817.

Taraxacum pinnis foliorum triangularibus, calyce inferne reflexo. HALL. *Helv.* 1. *Leontodon*. LINN. *Sp.* 1. *Dens Leonis, Taraxacum*. BLACKW. *T.* 1. C. B. 1. R. H. Vbique in gramineis. A.

Taraxacum foliis hispida triangulariter dentatis, caule erecto hispido, pappo plumoso. HALL. *H.* 6.

D d 5

Leon

^{hirsutum}
 Leontodon LINN. Sp. 6. Hieracium *asperum* ^{dentis} *leonis folio*. C. B. In sylvis & locis siccis. A.

^{Leontodon autumnale p. 422.}
 H. Taraxacum foliis semipinnatis, pinnis angulatis, floribus ante florescentiam nutantibus. ^{Crepis}
 HALL. G. 411. LINN. Sp. 6. Hieracium *apulum*, *flore suaverubente*. COLVMN. Ecphr. I. p. 242. it. Chondrilla *purpurascens foetida*. C. B. Prodr. T. 68. A.

H. Taraxacum (*Leontodon*) calyce toto erecto, inferioribus squamis siccis, foliis amplexicaulibus. GMEL. Sib. II. p. 16. T. 8. ^{Crepis} LINN. Sp. 7. Hieracioides *annua*, *Endiviae folio*, *calyce magnis latis ad basin cinctus*, *semina conica ex stylo longo papposa*, *placenta villosa*. Proprium genus constituere videntur.

H. Taraxacum (*Crepis*) foliis semipinnatis glabris, calycis foliis interioribus latissimis. HALL, G. 411. Fabrica priorum similiis videtur.

HIERACIVM. LINN. 818.

Hieracium foliis integerrimis ovatis, caule repente, scapo unifloro. ^{Pilosella} LINN. Sp. 5. HALL. H. 1. *Pilosella maior repens hirsuta*. C. B. I. R. H. BLACKW. T. 365. Vbique in gramineis.

Hieracium foliis virentibus pilosis ovato-lanceolatis, caule repente, scapo nudo multifloro HALL. H. 3. ^{subdum} LINN. Sp. 6. *Pilosella maior repens minus hirsuta*. C. B. In pratis. ^{for. in ang. ovatis} ^{fratibus repentibus, laxis nudo multifloro} ^{fr. 2. 2.} Hier-

Hieracium caule subnudo unifloro, foliis ellipticis dentatis & calyce longo villo barbato. HALL.

H. 8. ^{alpinum} LINN. Sp. 3. Hieracium alp. pumilum, folio ^{alpinum} anaginofo. C. B. In summo M. Bruellero.

Hieracium foliis longe lingulatis pilosis, caule ^{auricula} imo folioso, erecto, multifloro. LINN. Sp. 7.

Pilosella maior macrocaulis umbellifera. COLUMN.

Ecpbr. p. 249. ic. Pilosella maior erecta altera.

C. B. Vor dem Grobender Thor. ^{fr. in togerum laevis}

H. Hieracium foliis ovatis subdentatis tomentosis, scapo nudo racemoso. HALL. H. 7. ^{prostratum} LINN. Sp.

8. Hierac. prat. latifolium non sinuatum. C. B.

β. Var. angustifolia in M. Altenstollberg.

Hieracium foliis radicalibus ovatis dentatis pilosis, caule ^{in lacum} ramoso subnudo. HALL. Helv. 12.

LINN. Sp. 15. Hierac. murorum, folio pilosissimo.

C. B. I. R. H. Vbique ad muros & in sylvis.

Varietas, cui tumores pilosi ab infectis nati alis foliorum insident. Hieracium myophorum. RVPP.

Var. foliis longioribus, angustioribus, argute dentatis, caulibus magis foliosis. Circa Elbingerode.

H. Hieracium foliis ovatis integris, caule subnudo simplicissimo piloso corymbifero. aurantiacum. LINN. Sp. 9. HALL H. 6. Hierac. alp. non laciniatum, flore fusco. C. B. Pin. 128. Prodr. 65. In alpihus Helveticis.

H. Hieracium foliis radicalibus obovatis denticulatis, ^{caulinis} caulinis oblongis semiamplexicaulibus. LINN. Sp. 18. Hier. pyren. folio cerinthes latifolium.

lium. I. R. H. Hier. *pilosum*, foliis variis dentatis.
DILL. *Elth.* T. 149.

Hieracium foliis glabris amplexicaulibus dentatis, caule paniculato, calycibus hispidis. LINN. Sp. 16. HALL. H. 20. Hierac. *mont. latifolium glabrum minus*. C. B. In pratis vdis. v. g. *um die Papiermühle*.

(H. Hieracium caule ramoso, foliis firmis, infimis petiolatis, reliquis ex ovato-lanceolatis: omnibus sinuosis, petiolorum *instar* dentatis. GMEL. Sib. II. p. 26. T. 24. *Crepis* LINN. Sp. 9. In Sibiria. P.).

Hieracium solaudum. | *Hierac. umbellatum*. p. 44
CREPIS. LINN. § 19.

Crepis foliis ad terram ferratis, ad caulem pinnatis, pinnis alterne maioribus, caule nudo pene nudo multifloro. Hieracioides. HALL. *Helv.* 2. *Crepis*. LINN. Sp. 10. Hieracium *maius erectum angustifolium, caule laevi*. C. B. I. R. H. In agris.

Crepis (*Hieracioides*) foliis semipinnatis, subasperis amplexicaulibus, caule hispido folioso & multifloro. HALL. H. 3. *Crepis foliis bastato-lyratis, caule piloso*. LINN. Sp. 12. Hier. *Chondrillae folio hirsutum*. C. B. In agris & hortis.

Crepis foliis glabris integris & varie incisus, caule ramossimo, floribus minimis. HALL. H. 4. *Crepis*. LINN. Sp. 11. (*Hieracium chondrillae folio glabrum*. C. B.) In hortis.

Crepis foliis ovatis, caule brachiato, floribus perexiguis. HALL. G. 411. Hieracioides *annua* glu.

glutinosæ, floribus parvis. VAILL. *Act. Par.*
1721.

*Crepis foliis hirsutis pinnatis, pinnis angulatis
interne maioribus.* HALL. *H.* 1. ^{rebia} LINN. *Sp.* 13.

Hieracium amygdalarum amarum olens, f. odore Hier.

Apuli suave rubentis. I. R. H. *Über der Papier-
mühle, in M. Plessi sub arce. Ja monte prope Mafschmühle.*

H. crepis rubra ¹⁷ *crepis alpina* ^{p. 426.} *H. crepis sibirica* ^{p.}

INTYBUS, HALL. *Comment. Gott.* 1751.

p. 219.

Habitus Hieracii, sed placenta alveolata, & flores umbellati.

*Intybus caule folioso, foliis ovato-lanceolatis
dentatis hirsutis.* *Hieracium* ^{alpinum} HALL. *H.* 16. LINN.
Sp. 25. *Hier. frutic. folio subrotundo.* C. B. *Hiera-
cii Sabaudi varietas altera.* I. B. II. p. 1030. In
sylvis.

*Intybus caule folioso, foliis amplexicaulibus el-
lipticis acutis dentatis.* HALL. *G.* 419. *Helv.* p. 747.
Hierac. frutic. latifolium hirsutum. C. B.

*Intybus caule folioso, foliis linearibus lingulatis
dentatis firmis.* HALL. *Comm. Gott.* 1751. p. 219.
Hieracium. *Helv.* 22. LINN. ^{umbellatum} *Sp.* 26. *Hier. frutico-
sum angustifolium minus.* I. R. H. In dumetis.

β. Var. *foliis angustis vix dentatis.*

PICRIS. LINN. 812.

*Picris foliis amplexicaulibus lingulatis dentatis,
perianthiis imbricatis.* HALL. *H.* 751. ^{hieracoides} LINN. *Sp.* 2.
*Hieracium asperum maiori flore in agrorum limi-
bus.* I. R. H.

HEL-

((HELMINTHOTHECA. VAYLL. HALL.)

H. Helminthotheca. HALL. G. 413. *Picris perianthiis involacro maximo cinctis*. LINN. Sp. 1. *Hieracium echinoides, capitulis Cardui benedicti*. C. B. A.

SONCHVS. LINN. 813.

Sonchus pedunculis tomentosis, calycibus glabris. LINN. Sp. 4. *Sonchus foliis levibus teneris, calycibus glabris*. HALL. H. 3. *larvis laciniatus latifolius*. C. B. In hortis. A.

β. *Sonchus foliis rigidiusculis serratis, calycibus laevibus*. HALL. H. 2. *Sonchus asper laciniatus & non laciniatus*. C. B. In cultis. A.

Sonchus pedunculis calycibusque hispidis subumbellatis, foliis basi cordatis semipinnatis. LINN. Sp. 3. HALL. H. 1. *Sonchus repens, multis Hieracium maius*. I. R. H. In vdis segetibus. A.

H. *Sonchus* pedunculis calycibusque hispidis subumbellatis, foliis basi sagittatis. *palustris*. LINN. Sp. 2. *Sonchus asper arborescens*. C. B.

Sonchus pedunculis squamosis, floribus racemosis. LINN. Sp. 6. HALL. H. 4. *Sonchus laevis laciniatus coeruleus alpinus*. C. B. In sylvis Hercynicis circa M. Bruclerum. P.

H. *Sonchus* pedunculis subsquamosis, floribus racemosis, foliis lyrato-hastatis. LINN. Sp. 7. *Sonchus Canad. altissimus laciniatus, flore coerulescente*. VAYLL. Atl. Par. 1721. p. 198.

H. *Sonchus* pedunculis squamatis, foliis lanceolatis indivisis sessilibus. LINN. Sp. 8. GMEL. Sib. II.

P. 11. T. 3. *Lactuca Salicis folio, flore coerulea.*
AMMANN. *Ruth.* 211. In Sibiria.

PRENANTHES. LINN. 816.

Prenanthes flosculis quinis, foliis lyrato-hastatis.
^{mutata}
LINN. *Sp.* 4. **HALL.** *H.* 755. n. 1. *Lactuca sylv.*
murorum, flore luteo. I. B. II. p. 1004. *Sonchus*
laevior vulgaris. II. CLVS. CXLVI. In sylvis.

CHONDRILLA. LINN. 815.

Chondrilla ^{arvensis} **LINN.** *Sp.* 796. **HALL.** *H.* 755. 4
Chondrilla juncea viminea arvensis. **TABERN.** 487.
C. B. In apricis, v. g. über der Maschbrücke.

LACTVCA. LINN. 814.

H. *Lactuca foliis rotundatis, caule corymbofo.*
lativa. **LINN.** *Sp.* 2. **BLACKW.** *T.* 88. *capitata.* β.
non capitata. γ. *crispa.* In hortis cultura varie
 mutatur. A.

H. *Lactuca foliis ovatis dentatis, costa spinosa.*
HALL. *Gott.* 415. *Lactuca sylv. Italica, sanguineis*
maculis adpersa. **HERM.** *Par.* p. 194. *ic.* *Lactuca*
sylv. odore viroso. **C. B.** I. R. H. *L. scariola* L. 2 p. 1119.

Lactuca foliis semipinnatis, costa spinosa. **HALL.**
H. 3. *Lactuca sylv. costa spinosa.* **C. B.** I. R. H. In
 vallo vrbis. *Lactuca foliis carina aculeatis dentatis.*
^{crispa}
LINN. *Sp.* 3. ~~hanc cum priori iungit.~~

H. *Lactuca foliis lingulatis & semipinnatis, costae*
principio spinosae. **HALL.** *G. l. c. saligna* **LINN.** *Sp.*
S. *Lactuca sylv. altera, angusto foligno folio, costa*
 albi-

albicante. HALL. *Rupp. Jen.* 207. T. 4. In *Thuringia*. A.

H. *Lactuca* foliis glabris pinnatis, caule brachiato
HALL. H. 2. ^{*perennans*} LINN. *Sp.* 6. *Chondrilla coerulescens*.
TABERN. p. 176. In *Thuringia*.

SCORZONERA. LINN. 811.

Scorzonera foliis angustissimis semipinnatis & integris, caule folioso. HALL. H. 2. *Scorz. foliis linearibus dentatis*. LINN. *Sp.* 6. *Scorz. foliis laciniatis*. I. R. H. Ad vias. v. g. circa *Gescheröder Feld*.

β. *Scorzonera* foliis semipinnatis, caulibus erectis ramosis & foliosis. HALL G. 416.

(*Scorzonera* foliis semipinnatis, caule pene nudo, pappo sessili. HALL. H. 4. *Leontodon*. LINN. *Sp.* 4. *Hieracium minus praemorsa radice*. LOB. ic. 237. *Hierac. Chondrillae & Coronopi folio*. C. B. I. R. H. In pratis & gramineis. v. g. gegen der *Maschmühle*.)

H. *Scorzonera* foliis gramineis firmulis, caule brachiato multifloro. HALL. G. 417.

H. *Scorzonera* caule ramofo, foliis amplexicaulibus integris ferrulatis. LINN. *Sp.* 2. *Scorz. latifolia sinuata*. C. B. BLACKW. T. 406. *Scorz. maior* HISP. L. CLVS. CXXXVII. In *Hispania*. *Sibiria*.

H. *Scorzonera* caule subnudo, foliis nervosis planis ovato-lanceolatis. LINN. *Sp.* 1. *Mat. med.* 371. *Scorz. humilis latifolia nervosa*. C. B. I. R. H. CLVS. CXXXVII. In pratis & caeduis sylvarum. v. g. circa *Berolinum*. Spontanea caule *uniflora*, in hortis culta *multiflora*.

H. Scorzonera foliis omnibus antrorsum dentato-sinuatis amplexicaulibus, squamis calycis ovatis alatis. ^{linnæarum} LINN. Sp. 7. Chondrilla *Tingitana*, floribus luteis, papaveris bortenſis folio. HERM. Lugdb. p. 659. ic.

H. Scorzonera foliis glaucis, interioribus semipinnatis, superioribus amplexicaulibus integerrimis, squamis calycis cordatis alatis. ^{herzoides} LINN. Sp. 8. Crepis. HALL. G. 413. Vtraque modo dicta dubii generis.

^{H. scorzonera graminea folio. V. infra.}
TRAGOPOGON. LINN. 810.

Tragopogon calycibus corollae radium aequantibus, foliis carinatis amplexicaulibus gramineis. ^{pratensis} LINN. Sp. 1. HALL. H. 758. Tragopogon *pratense* luteum minus. C. B. 1. R. H. In vallo vrbis & pratis.

β. Var. flore viridi. Im Göttinger Wald.

H. Tragopogon calycibus corollae radio longioribus, foliis integris strictis, pedunculis superne incrassatis. ^{porri folium} LINN. Sp. 3. Tragop. *purpureo-coeruleum* Porri folio. C. B. Tragop. *crocifolium mont.* flore nigro purpureo. COLVMN. Ecphr. p. 230. ic.

H. Tragopogon calycibus corollae radio longioribus, foliis integris, seminibus laevibus, disci plumosis, radii setaceis. ^{anagyris (p. 1. 1.) hyemalis} LINN. Sp. 4. H. V. f. 243. *. Tragop. *gramineo folio, flore suaverubente.* COLVMN. Ecphr. I. T. 231. A.

H. Tragopogon calycibus corollae radium aequantibus, foliis lineari-ensiformibus integris carinatis. ^{graminis folio} Scorzonera. LINN. Sp. 3. Scorz. *Lusitanica, gramineo folio, flore pallide luteo.* BVXB. Gent. II. p. 26. T. 21.)

H. *Tragopogon calycibus corolla brevioribus inermibus, foliis lyrato-sinuatis*. ^{*calycibus*} LINN. *Sp.* 6. H. *Vps.* 244. *Chondrilla prior*. CLVS. CXLIII. A.

H. *Tragopogon calycibus corolla brevioribus aculeatis, foliis pinnato-hastatis*. ^{*pinnatis*} LINN. *Sp.* 7. HALL. G. 419. *Sonchus asper laciniatus Creticus*. C. B. *Prodr.* 60. A.

H. *Tragopogon calycibus corolla brevioribus hispida, foliis scabris caulinis oblongis, caule ramossimo*. ^{*asperum*} LINN. *Sp.* 8. *Sonchus asper subrotundo folio, maior & minor*. C. B. *Prodr.* 60. *Folia aspera, margine aculeis exiguis scabra. Folia calycis simplici serie, pilis rigidis hispida, accessoriis ad basin foliolis vagis laxis. Semina omnia incurva ex brevissimo stylo papposa, exteriora foliis cavinatis calycis involuta pappo breviori*. A.

Placenta pilosa, semine papposo.

ANDRYALA. LINN. 820.

H. *Andryala foliis integris*. ^{*in brevifolia*} LINN. *Sp.* 1. *Eriophoros foliis dentatis, flore extus melino, intus subrubente, & utrinque luteo*. VAILL. 1721. p. 277. *Sonchus lanatus*. DALECH.

Placenta paleacea, semine papposo.

HYPOCHAERIS. LINN. 822.

Hypochaeris foliis dentato-sinuatis scabris, caule ramoto nudo, pedunculis squamosis. ^{*radicata*} LINN. *Sp.* 6. HALL. H. 1. *Hieracium dentis leonis folio obtuso maius*. C. B. *Hierac. intybaceum*. TABERN. 489. In sylvis & gramineis. *sem. 62. Flor. Dan. T. 150.*

Hypo-

Hypochaeris glabra, calycibus oblongis imbricatis, caule nudo, foliis glabris dentato-sinuatis. LINN. Sp. 4. Hypochaeris *chondrillae folio, parvo flore.* VAILL. 1721. p. 14. Hyoseris *altera.* TABERN. 490. In gramineis M. Hagen.

Hypochaeris caule subnudo, ramo solitario, foliis ovato-oblongis dentatis hirsutis. LINN. Sp. 4. Hieracium *alp. latifolium, flore magno.* C. B. HALL. H. n. 2. T. 24. Hierac. *latifolium.* CLVS. CXXXIX. In pratis *bercynicis* circa *Elbingerode.* *ed. Fur. D. 147.*

H. Hypochaeris *calycibus* aequalibus hispidis. *Achyrophorus.* LINN. Sp. 2. Rhagadiolus. HALL. G. 422. n. 3. In Creta. A. *seriola* *seriola athenensis* Linn. p. 2. *Es. 2.*

Semine foliis coronato.

CATANANCHE. LINN. 824.

H. Catananche squamis calycinis inferioribus lanceolatis. *lutea.* LINN. Sp. 2. Chondrilla *cyanoidea lutea, coronopi folio non diviso.* BARR. ic. 1135. In Creta. A.

H. Catananche squamis calycinis inferioribus ovatis. *coerulea.* LINN. Sp. 1. Chondrilla *coerulea cyani capitulo.* C. B. Chondrilla. III. DOD. Pempt. 638. In Provincia. A.

HYOSERIS.

Semina exigua recta setis rigidis paucioribus coronata, in ambitu saepe nuda.

H. Hyoseris scapis unifloris nudis, foliis glabris lyrato-hastatis angulatis, seminibus omnibus setis coronatis. LINN. Sp. 2. ROYEN. 127. Taraxaco-

Ee 2

nastrum

nastrum *dentis leonis folio ad summitatem radiata*.
 VAILL. *Parif.* 121. p. 180. In *Gallia Narbonensi*.

(H. *Hyoseris* caule folioso ramofo, calyce fetis longissimis circumvallato, feminibus in ambitu nudis. Rhagadiolus HALL. 421. *Crepis barbata* LINN. Sp. 4. Hieracium calyce barbato. COLUMN. *Ecpbr.* II. p. 28. ic. Hier. Baeticum medio nigrum HERM. *Par.* p. 185. ic. In Italiae maritimis.)

~~*Hyoseris*~~
 (RHAGADIOLVS.)

Semina longa cylindrica, in ambitu foliis calycis obvoluta, corona ciliata recta, in medio fetis coronata.

H. Rhagadiolus capitulo subgloboso. squamis calycis feminigeris longioribus hispidis. HALL. G. 422. An *Hyoseris* ^{*hyoseris*} LINN. Sp. 6? *Squamae calycis dorsae undique scabrae. Setae feminum albae geniculo articulatae.*

H. Rhagadiolus capitulo ovato, squamis calycis feminigeris longioribus, inferne glabris, apicem versus dorso scabris. *Hyoseris* ^{*Rhagadiolus*} LINN. Sp. 7. *Lapsana* H. Kps. 6. *Squamae calycis glabrae, apicem versus serrato-denticulatae, setae feminum subfuscae capillares.*

H. Rhagadiolus capitulo depresso angulato, squamis calycis feminigeris brevioribus, glabris, gibbis nodosis. HALL. G. 422. *Lapsana* ^{*Zacantha*} LINN. Sp. 2. *Cichorium verrucarium, Zacantha*. CLVS. CXLIV.

Hyoseris minima p. 437.

LAPSA-

LAPSANA. LINN. 823.

Semina omnia nuda.

Lapsana calycibus fructus angulatis, pedunculis tenuibus ramosissimis. LINN. Sp. 1. Lapsana foliis imis pinnatis, superioribus lanceolatis serratis. HALL. Helv. 761. Lapsana. I. R. H. DODON. Pempt. 675. In sylvis caeduis & locis ruderosis.

Lapsana calycibus fructus depressis, caule nudo diviso, foliis ovatis dentatis. HALL. G. 423. *Hyoferis* LINN. Sp. 5. *Hyoferis angustifolia*. TABERN. 180. ic. Hieracium *minimum*. CLYS. CXLII. In arenosis Germanidis. *Non cedit* / *Abundat*

H. Lapsana calycibus fructus vndique patentibus, radiis subulatis, foliis lanceolatis indivisis. *stellata*. LINN. Sp. 3. Hieracium *Narbonense falcatum*. LOB. ic. 240.

H. Lapsana calycibus fructus vndique patentibus, radiis subulatis, foliis lyratis, lobis rotundatis. LINN. Sp. 4. H. Vp. 245. Rhadagiolus *Lapsanae foliis*. I. R. H.

H. *Lapsana Lavantha* p. 426.

CICHORIVM. LINN. 825.

Cichorium caule simplici, foliis dentato-sinuatis, pinnis triangularibus. LINN. Sp. 1. Cichorium *gylvestre* f. *Officinatum*. C. B. P. R. H. BLACKW. T. 183. Vbique ad oras agrorum.

H. Cichorium caule simplici, foliis integris crenatis. LINN. Sp. 2. Cichorium *latifolium* f. *Endivia vulgaris*. C. B. Intybum *sativum*. DOD. Pempt. 634. A.

H. Cichorium caule dichotomo spinoso. ^{spinosum} LINN.
 Sp. 3. Cichorium ^{spinosum} C. B. Prodr. p. 62. ii.
 Chondrilla genus elegans, coeruleo flore. CLVI,
 CXLV. Ad litora maris Siciliae. A.

SCOLYMVS. LINN. 826.

H. Scolymus foliis margine attenuatis. ^{maculatus} LINN.
 Sp. 1. Scolymus Theophrasti, Narbonensis. CLVI.
 CLIII. In Europa australi.

APPENDIX.

OMISSA, ET ADDENDA.

Pag. 11. post. l. 25.

CERATONIA. LINN. 983.

H. Ceratonia. ^{filiqua} LINN. Sp. p. 1026. Mat. med. 455.
 Siliqua edulis. C. B. Carobe filiqua. BLACKW. T.
 209. In Europa australi. Planta diphyta, fr. legu-
 men magnum compressum.

Pag. 25. post. l. 16.

PETIVERIA. LINN. 417.

Stamina sex, styli quatuor.

H. Petiveria ^{alliacca} LINN. Sp. p. 343. AB,
 Stockh. 1744. p. 287. T. 7. Petiveria Solani foliis,
 loculis spinosis. PLUM. Gen. 50. In Jamaica.

Pag.

Pag. 26. post. l. 15.

RIVINIA. LINN. 150.

H. Rivinia ^{humilis} LINN. *Spec. p.* 121. Rivinia *humilis racemosa*, baccis puniceis. PLUM. *Gen.* 48. Solanoides Amer. *Circææ foliis*. TOURN. *Art. Par.* 1706. *Lin. Sp. pl.* 2 p. 177.

P. 83. post. l. 10.

PHYTOLACCA. LINN. 521.

H. Phytolacca baccis petiolatis, decem striis notatis. HALL. *G.* 211. ^{decandrum} *Virgaria*, DILL. *H. Elth. T.* 239. f. 309, Virginiae indigena, nunc sponte in Italia provenit. P.

H. Phytolacca baccis sessilibus, octo striis notatis. HALL. *G.* 212. DILL. *Elth. T.* 308. *Flores subvirides*. Phytolacca *foliis integerrimis*. LINN. *Sp. p.* 441. utramque iungit. P.

P. 90. post. l. 7.

CANNA. LINN. 1. Cannacorus. T.

Stamen unicum & stylus unicus figura laciniae corollae.

H. Canna foliis ovatis utrinque acuminatis nervosis, flore luteo punctato. LINN. *Sp.* 1. qui hanc & sequentem iungit. I. R. H. *Riv. T.* III. P.

H. Canna foliis ovatis utrinque acuminatis nervosis, flore coccineo splendente. HALL. *G.* 66. I. R. H. *Riv. T.* 110. *Commer. Litt. Nor.* 1743. p. 290. *Tab. II. f. 3. sqq.* P.

11

Ec 4

Pag.

H. *Cichorium caule dichotomo spinoso*. ^{*spinosum*} LINN.
Sp. 3. Cichorium spinosum. C. B. Prodr. p. 62. ic.
Chondrillae genus elegans, coeruleo flore. CLVS.
 CXLV. Ad litora maris *Siciliae. A.*

SCOLYMVS. LINN. 826.

H. *Scolymus foliis margine attenuatis*. ^{*maculatus*} LINN.
Sp. 1. Scolymus Theophrasti, Narbonensis. CLVS.
 CLIII. In *Europa australi.*

APPENDIX.

OMISSA, ET ADDENDA.

Pag. 11. post. l. 25.

CERATONIA. LINN. 983.

H. *Ceratonia*. ^{*aliqua*} LINN. *Sp. p. 1026. Mat. med. 455.*
Siliqua edulis. C. B. Carobe siliqua. BLACKW. T.
 209. In *Europa australi. Planta diphyta, fr. legu-*
men magnum compressum.

Pag. 25. post. l. 16.

PETIVERIA. LINN. 417.

Stamina sex, styli quatuor.

H. *Petiveria alliacea. LINN. Sp. p. 342. AB.*
Stockb. 1744. p. 287. T. 7. Petiveria Solani foliis,
locutis spinosis. PLUM. Gen. 50. In Jamaica.

Pag.

an ledifolius?

An *Cistus salicifolius*. LINN. *Sp.* 19. ? *Cistus an-*
-uus I. CLVS. 76. Icon sequenti magis convenit. A.

H. *Cistus* herbaceus parulus villosus stipulatus,
floribus solitariis alaribus & terminalibus, stipulis
minimis. *Priori ex habitu similis, sed in eodem li-*
-et solo multoties minor. Stipulae angustissimae.
Flores alterni, plerique tamen cauliculi unisflori. A.

Pag. 275. post. l. 17.

FRAXINVS. LINN. 1026.

Fraxinus foliolis ferratis, floribus apetalis. *exce-*
-pt- LINN. *Sp.* 1. HALL. *H.* 167. C. R. I. R. H.
In sylvis & sepibus.

P. 312. post. l. 16.

GLECHOMA. LINN. 643. *Chamaeclema*.

T. HALL.

Glechoma (*Chamaeclema*) caulibus procumben-
tibus radicans, foliis reniformibus crenatis. HALL.
H. 1. LINN. *Sp.* 2. *Hedera terrestris vulgaris.*
C. B. BLACKW. T. 225. Ad sepes & sub arboribus.

β. *Chamaeclema flore magno.* HALL. *H.* 2. *He-*
-dera terrestris montana. C. B. In vallo suburbano.
Num Varietas?





the first of these is the fact that the
the second is the fact that the
the third is the fact that the
the fourth is the fact that the
the fifth is the fact that the
the sixth is the fact that the
the seventh is the fact that the
the eighth is the fact that the
the ninth is the fact that the
the tenth is the fact that the
the eleventh is the fact that the
the twelfth is the fact that the
the thirteenth is the fact that the
the fourteenth is the fact that the
the fifteenth is the fact that the
the sixteenth is the fact that the
the seventeenth is the fact that the
the eighteenth is the fact that the
the nineteenth is the fact that the
the twentieth is the fact that the
the twenty-first is the fact that the
the twenty-second is the fact that the
the twenty-third is the fact that the
the twenty-fourth is the fact that the
the twenty-fifth is the fact that the
the twenty-sixth is the fact that the
the twenty-seventh is the fact that the
the twenty-eighth is the fact that the
the twenty-ninth is the fact that the
the thirtieth is the fact that the
the thirty-first is the fact that the
the thirty-second is the fact that the
the thirty-third is the fact that the
the thirty-fourth is the fact that the
the thirty-fifth is the fact that the
the thirty-sixth is the fact that the
the thirty-seventh is the fact that the
the thirty-eighth is the fact that the
the thirty-ninth is the fact that the
the fortieth is the fact that the
the forty-first is the fact that the
the forty-second is the fact that the
the forty-third is the fact that the
the forty-fourth is the fact that the
the forty-fifth is the fact that the
the forty-sixth is the fact that the
the forty-seventh is the fact that the
the forty-eighth is the fact that the
the forty-ninth is the fact that the
the fiftieth is the fact that the
the fifty-first is the fact that the
the fifty-second is the fact that the
the fifty-third is the fact that the
the fifty-fourth is the fact that the
the fifty-fifth is the fact that the
the fifty-sixth is the fact that the
the fifty-seventh is the fact that the
the fifty-eighth is the fact that the
the fifty-ninth is the fact that the
the sixtieth is the fact that the
the sixty-first is the fact that the
the sixty-second is the fact that the
the sixty-third is the fact that the
the sixty-fourth is the fact that the
the sixty-fifth is the fact that the
the sixty-sixth is the fact that the
the sixty-seventh is the fact that the
the sixty-eighth is the fact that the
the sixty-ninth is the fact that the
the seventieth is the fact that the
the seventy-first is the fact that the
the seventy-second is the fact that the
the seventy-third is the fact that the
the seventy-fourth is the fact that the
the seventy-fifth is the fact that the
the seventy-sixth is the fact that the
the seventy-seventh is the fact that the
the seventy-eighth is the fact that the
the seventy-ninth is the fact that the
the eightieth is the fact that the
the eighty-first is the fact that the
the eighty-second is the fact that the
the eighty-third is the fact that the
the eighty-fourth is the fact that the
the eighty-fifth is the fact that the
the eighty-sixth is the fact that the
the eighty-seventh is the fact that the
the eighty-eighth is the fact that the
the eighty-ninth is the fact that the
the ninetieth is the fact that the
the ninety-first is the fact that the
the ninety-second is the fact that the
the ninety-third is the fact that the
the ninety-fourth is the fact that the
the ninety-fifth is the fact that the
the ninety-sixth is the fact that the
the ninety-seventh is the fact that the
the ninety-eighth is the fact that the
the ninety-ninth is the fact that the
the hundredth is the fact that the

INDEX GENERVM

<i>Draba</i>	322	<i>Festuca</i>	66
<i>Dracocephalum</i>	309	<i>Ficaria</i>	131
<i>Drosera</i>	182	<i>Ficoidea</i>	166
E.		<i>Ficoidea</i>	163
<i>Ebulus</i>	209	<i>Ficus</i>	22
<i>Echinops</i>	384	<i>Filago</i>	400
<i>Echium</i>	255	<i>Filipendula</i>	119
<i>Elaterium</i>	246	<i>Foeniculum</i>	213
<i>Elatine</i>	282	<i>Foenum graecum</i>	364
<i>Empetrum</i>	15	<i>Fragaria</i>	138
<i>Ephedra</i>	2	<i>Frangula</i>	16
<i>Ephemerum</i>	90	<i>Fraxinella</i>	120
<i>Epilobium</i>	199	<i>Fraxinus</i>	441
<i>Epimedium</i>	273	<i>Fritillaria</i>	101
<i>Epipactis</i>	85	<i>Fumaria</i>	374
<i>Erica</i>	202	G.	
<i>Erigeron</i>	403	<i>Galanthus</i>	94
<i>Erioccephalus</i>	416	<i>Gale</i>	11
<i>Eriophorum</i>	79	<i>Galega</i>	350
<i>Eruca</i>	331	<i>Galeopsis</i>	302
<i>Erucago</i>	320	<i>Galium</i>	231
<i>Ervum</i>	358	<i>Garidella</i>	125
<i>Eryngium</i>	228	<i>Genista</i>	360
<i>Eryfinum</i>	328	<i>Gentiana</i>	245
<i>Erythrina</i>	361	<i>Geranium</i>	173
<i>Erythronium</i>	102	<i>Geum</i>	171 124
<i>Esculus</i>	204	<i>Gladiolus</i>	91
<i>Evonymus</i>	208	<i>Glaucium</i>	116
<i>Eupatorium</i>	405	<i>Glaux</i>	252
<i>Euphorbia</i>	46	<i>Glechoma</i>	441
<i>Euphrasia</i>	289	<i>Gleditsia</i>	374
F.		<i>Glycine</i>	340
<i>Faba</i>	375	<i>Glycirrhiza</i>	351
<i>Fabago</i>	183	<i>Gnaphalium</i>	400
<i>Fagonia</i>	182	<i>Gnaphalodes</i>	15
<i>Fagopyrum</i>	42	<i>Gomphrena</i>	34
<i>Fagus</i>	10	<i>Gossypium</i>	158
<i>Ferrum equinum</i>	346	<i>Gramen</i>	58
		<i>Granadilla</i>	248

INDEX GENERVM.

Gratiola	282	Hyoferis	435
Guaiava	144	Hypocoum	271
Gypsophila	193	Hypericum	149
H.		Hypochaeris	434
Haemanthus	91	Hypopitys	184
Harmala	183	Hyssopus	308
Hedera	23	Hysterophorus	14
Hedysarum	248		
Helenia	417	I.	
Helianthus	408	Jalappa	241
Helianthemum	440	Jasione	244
Heliotropium	257	Jasminum	274
Helleborine	86	Jatropha	51
Helleborus	125	Iberis	325
Helminthotheca	430	Illecebrum	44
Helxine	43	Impatiens	273
Hemerocallis	100	Imperatoria	222
Hepatica	132	Indigofera	350
Heracleum	223	Intybus	429
Hermannia	158	Inula	417
Herniaria	43	Ipomaea	239
Hesperis	337	Iris	92
Heuchera	198	Isatis	320
Hibiscus	151	Isopyrum	125
Hieracium	426	Iva	14
Hippocastanum	204	Juglans	9
Hippocrepis	346	Juiuba	17
Hippophae	12	Juncus	80
Hordeum	71	Juniperus	2
Horminum	311	Justicia	285
Horminum	296	Ixia	93
Hortonia	253	K.	
Humulus	23	Kali	37
Hyacinthus	102	Ketmia	151
Hydrocharis	113	Kleinia	405
Hydrocotyle	213	Knautia	383
Hyoscyamus	267	Knawel	18

L.

INDEX GENERVM.

L.		Linum	180
Lactuca	431	Liquidambar	11
Lagoecia	377	Liriodendrum	127
Lagurus	63	Lithospermum	257
Lamium	302	Lobelia	244
Lampfuna	437	Lolium	67
Lantana	277	Lonicera	210
Lapathum	38	Lorus	344
Lappa	386	Lunaria	323
Lapsana	437	Lupinus	362
Larix	3	Lupulus	23
Laserpitium	222	Luteola	123
Lathraea	277	Lychnidea	253
Lathyrus	352	Lychnis	192
Lavatera	154	Lycium	254
Lavendula	315	Lycopsis	260
Laurocerasus	144	Lycopus	317
Laurus	21	Lyfimachia	250
Lens	358	Lythrum	197
Lentiscus	11		
Leontodon	425	M.	
Leonurus	300	Maiorana	316
Lepidium	328	Malva	154
Lerchia	30	Malus	143
Leucoium	95	Mariscus	79
Libanotis	226	Marrubium	306
Ligusticum	218	Matricaria	413
Ligustrum	275	Maurocenia	210
Lilac	274	Mays	73
Lilioasphodelus	100	Medicago	365
Lilium	101	Melampyrum	288
Limnopeuce	55	Melia	182
Limon	146	Meianthus	271
Limonium	376	Melica	64
Limosella	276	Melilotus	368
Linagrostis	79	Melissa	312
Linaria	282	Melittis	311
Linofyris	18	Melo	247

INDEX GENERVM.

<i>Glypeola</i>	322	<i>Croton</i>	
<i>Cneorum</i>	50	<i>Crucianella</i>	
<i>Cnicus</i>	390	<i>Cucubalus</i>	
<i>Cochlearia</i>	324	<i>Cucumis</i>	
<i>Coffea</i>	254	<i>Cucurbita</i>	
<i>Coix</i>	73	<i>Cuminoides</i>	
<i>Colchicum</i>	94	<i>Cunila</i>	
<i>Colocynthis</i>	247	<i>Cupressus</i>	
<i>Colutea</i>	350	<i>Cuscuta</i>	
<i>Comarum</i>	135	<i>Cyanus</i>	
<i>Commelina</i>	90	<i>Cyclamen</i>	
<i>Conium</i>	217	<i>Cydonia</i>	
<i>Convolvularia</i>	103	<i>Cynapium</i>	
<i>Convolvulus</i>	238	<i>Cynara</i>	
<i>Conyza</i>	404	<i>Cynoglossum</i>	
<i>Cornalodendron</i>	361	<i>Cynofurus</i>	
<i>Corchorus</i>	147	<i>Cyperoides</i>	
<i>Coreopsis</i>	408	<i>Cyperus</i>	
<i>Coriandrum</i>	216	<i>Cypripedium</i>	
<i>Corispermum</i>	36	<i>Cytisus</i>	
<i>Cornus</i>	234		D.
<i>Corona imperialis</i>	101	<i>Damasonium</i>	
<i>Coronilla</i>	349	<i>Daphne</i>	
<i>Coronopus</i>	325	<i>Datura</i>	
<i>Corrigiola</i>	45	<i>Daucus</i>	
<i>Corylus</i>	10	<i>Delphinium</i>	
<i>Cotinus</i>	208	<i>Dens Canis</i>	
<i>Cotula</i>	399	<i>Dentaria</i>	
<i>Cotyledon</i>	169	<i>Diastamus</i>	
<i>Crambe</i>	320	<i>Dianthus</i>	
<i>Crassula</i>	167	<i>Diervilla</i>	
<i>Crataegus</i>	142	<i>Digitalis</i>	
<i>Crepis</i>	428	<i>Dipsacus</i>	
<i>Crinum</i>	111	<i>Dodartia</i>	
<i>Critimum</i>	220	<i>Dolichos</i>	
<i>Crocus</i>	94	<i>Doronicum</i>	
<i>Crotalaria</i>	361	<i>Dorycnium</i>	

INDEX GENERVM.

P.			
<i>Padus</i>	144	Phleum	60
<i>Paeonia</i>	126	Phlomis	301
<i>Pafiarus</i>	16	Phlox	253
<i>Palma</i>	83. 84	Phytica	50
<i>Panicastrella</i>	73	Phyllis	212
<i>Panicum</i>	61	Phytalis	264
<i>Papaver</i>	114	Phyteuma	244
<i>Parietaria</i>	30	Phytolacca	439
<i>Paris</i>	184	Pictis	429
<i>Parkinsonia</i>	373	Pimpinella	215
<i>Parnassia</i>	124	Pinus	3
<i>Parthenium</i>	14	Pistacia	11
<i>Passiflora</i>	248	Pisum	352
<i>Pastinaca</i>	214	<i>Plantaginella</i>	276
<i>Pavia</i>	204	<i>Plantago</i>	236
<i>Pedicularis</i>	287	Platanus	10
<i>Peganum</i>	183	Plumbago	231
<i>Peireskia</i>	161	Poa	64
<i>Pelecinus</i>	340	<i>Podagraria</i>	214
<i>Pentapetes</i>	158	Polemonium	241
<i>Pentaphyllum</i>	135	Polyanthes	103
<i>Pentapteris</i>	57	Polygala	375
<i>Peplis</i>	197	Polygonatum	103
<i>Percbepier</i>	139	Polygonum	43
<i>Periploca</i>	269	<i>Populago</i>	127
<i>Perfica</i>	145	Populus	7
<i>Perficaria</i>	41	Portulaca	167
<i>Pervinca</i>	268	Potamogeton	56
<i>Petasitis</i>	404	Potentilla	135
<i>Petiveria</i>	438	<i>Poterium</i>	235
<i>Peucedanum</i>	220	Prasium	311
<i>Phaca</i>	342	Prenanthes	431
<i>Phalaris</i>	59	<i>Pumula</i>	253
<i>Rharnaceum</i>	45	<i>Prunus</i>	144
<i>Phaseolus</i>	362	Psidium	144
<i>Phellandrium</i>	216	Pforalea	343
<i>Philadelphus</i>	147	<i>Psyllium</i>	236
		<i>Releae similis</i>	207

F f 5

Pulegium

INDEX · GENERVM.

Gratiola	282	Hyoferis
<i>Guaicua</i>	144	Hypocoum
Gypsophila	193	Hypericum
H.		Hypochaeris
Haemanthus	91	<i>Hypopitys</i>
<i>Harmala</i>	183	Hyslopous
Hedera	23	Hysterophorus
Hedysarum	48	
Helenia	417	I.
Helianthus	408	<i>Jalappa</i>
<i>Helianthemum</i>	440	Jasione
Heliotropium	257	Jasminum
<i>Helleborine</i>	86	Jatropha
Helleborus	125	Iberis
Helminthotheca	430	Illecebrum
<i>Helxine</i>	42	Impatiens
Hemerocallis	100	Imperatoria
Hepatica	132	Indigofera
<i>Heracleum</i>	223	Intybus
Hermannia	158	Inula
Herniaria	43	Ipomaea
Hesperis	337	Iris
Heuchera	198	Itatis
Hibiscus	151	<i>Isopyrum</i>
Hieracium	426	Iva
<i>Hippocastanum</i>	204	Juglans
Hippocrepis	346	<i>Juiuba</i>
Hippophae	12	Juncus
Hordeum	71	Juniperus
Horminum	311	Justicia
<i>Horminum</i>	296	Ixia
Hortonia	253	K.
Humulus	23	<i>Kali</i>
Hyacinthus	102	<i>Ketmia</i>
Hydrocharis	113	<i>Kleinia</i>
Hydrocotyle	213	Knautia
Hyoicyamus	267	<i>Knawel</i>

INDEX GENERVM.

L.			
ca	431	Linum	180
ecia	377	Liquidambar	11
us	63	<i>Liriodendrum</i>	127
um	302	Lithospermum	257
<i>ysana</i>	437	Lobelia	244
ina	277	Lolium	67
thum	38	<i>Lonicera</i>	210
a	386	Lorus	344
na	437	Lunaria	323
	3	Lupinus	362
pitium	222	<i>Lupulus</i>	23
aea	277	<i>Luteola</i>	123
rus	352	<i>Lychnidea</i>	253
tera	154	Lychnis	192
ndula	315	Lycium	254
ocerasus	144	Lycopsis	260
is	21	Lycopus	317
	358	Lyfimachia	250
<i>fcus</i>	11	Lythrum	197
<i>oden</i>	425		
urus	300	M.	
dium	328	Maiorana	316
ia	30	Malva	154
oium	95	<i>Malus</i>	143
otis	226	Mariscus	79
ficum	218	Marrubium	306
hrum	275	Matricaria	413
	274	<i>Maurocenia</i>	210
<i>asphodelus</i>	100	<i>Mays</i>	73
m	101	Medicago	365
opeuce	55	Melampyrum	288
n	146	Melia	182
nium	376	Meianthus	271
ofella	276	Melica	64
groftis	79	Melilotus	368
ria	282	Melissa	312
<i>lyris</i>	18	Melittis	311
		<i>Melo</i>	247

Ef 4

Melo-

1844. Zuerich. 11. Jan. 1. 31.

Leider, in selbst in dem Leipz. Z. p. 326.

63



Johns : Burnett. Jan. 15, 1891.

Revised in report in Conn. Geol. 2. p. 326.

63



INDEX GENERVM.

Trapa	207	Veronica	177
Tribulus	182	Viburnum	209
<i>Tribulus aquaticus</i>	207	Vicia	355
<i>Triclis</i>	44	Vinca	268
Trientalis	252	Viola	272
Trifolium	369	<i>Virga aurea</i>	420
Triglochin	81	Viscago	183
Trigonella	364	Viscum	16
<i>Trinitas</i>	132	Vitex	277
Triticum	66	Vitis	23
<i>Trollius</i>	125	<i>Vlmaria</i>	119
Tropaeolum	204	Vlmus	20
<i>Tuberosa</i>	103	Vnifolium	104
Tulipa	102	Vrena	157
<i>Tulipifera</i>	127	<i>Vrtica</i>	31
<i>Tuna</i>	161	<i>Vua rafi</i>	201
<i>Tunica</i>	195	Vulneraria	343
Turritis	335		
Tussilago	404	X.	
Typha	82	Xanthium	13
<i>Typhoides</i>	60	Xeranthemum	406
		<i>Xiphium</i>	92
V.		Y.	
Vaccinium	200	Yucca	100
<i>Vaillantia</i>	231		
Valeriana	377	Z.	
<i>Valerianella</i>	378	<i>Zacintba</i>	436
Vella	321	Zanichellia	55
Veratrum	440	Zea	73
Verbascum	265	Ziziphora	299
Verbena	318	<i>Ziziphus</i>	16
Verbefina	407	Zygophyllum	183



1. Holstein: Jly.

Obbromt. Gm.

Trunk Bergsteig n. östl. o. Ende.

höchstgeleg. Flora Einsamkeit. J. 1912

←

1. Stahl . Zurück . 2. 11. 1941 . 1. 31.

Review is set in Am. Leg. v. 326.

63

